

(19) World Intellectual Property  
Organization  
International Bureau



(43) International Publication Date  
11 March 2004 (11.03.2004)

PCT

(10) International Publication Number  
**WO 2004/020637 A1**

(51) International Patent Classification: C12N 15/53

(21) International Application Number:  
PCT/AU2003/001111

(22) International Filing Date: 29 August 2003 (29.08.2003)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
2002951088 30 August 2002 (30.08.2002) AU  
2002952835 16 September 2002 (16.09.2002) AU

(71) Applicant (for all designated States except US): INTERNATIONAL FLOWER DEVELOPMENTS PTY. LTD. [AU/AU]; 16 Gipps Street, Collingwood, Victoria 3066 (AU).

(72) Inventors; and

(75) Inventors/Applicants (for US only): BRUGLIERA, Filippa [AU/AU]; 11 Kalimna Street, Preston, Victoria 3072 (AU). TANAKA, Yoshikazu [JP/JP]; 2-7-4 Ohginosato, Otsu, Shiga 520-0246 (JP). MASON, John [AU/AU]; 9/999 Rathdowne Street, Carlton North, Victoria 3054 (AU).

(74) Agents: HUGHES, E, John, L et al.; DAVIES COLLISON CAVE, 1 Little Collins Street, Melbourne, Victoria 3000 (AU).

(81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

**Published:**

- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: FLAVANOID 3',5'-HYDROXYLASE GENE SEQUENCES AND USES THEREFOR

(57) Abstract: The present invention relates generally to a genetic sequence encoding a polypeptide having flavonoid 3',5'-hydroxylase (F3'5'H) activity and to the use of the genetic sequence and/or its corresponding polypeptide thereof *inter alia* to manipulate color in flowers or parts thereof or in other plant tissue. More particularly, the F3'5'H has the ability to modulate dihydrokaempferol (DHK) metabolism as well as the metabolism of other substrates such as dihydroquercetin (DHQ), naringenin and eriodictyol. Even more particularly, the present invention provides a genetic sequence encoding a polypeptide having F3'5'H activity when expressed in rose or gerbera or botanically related plants. The instant invention further relates to antisense and sense molecules or RNAi-inducing molecules corresponding to all or part of the subject genetic sequence or a transcript thereof. The present invention further relates to promoters which operate efficiently in plants such as rose, gerbera or botanically related plants.

WO 2004/020637 A1

# FLAVONOID 3',5' HYDROXYLASE GENE SEQUENCES AND USES THEREFOR BACKGROUND OF THE INVENTION

## 5 FIELD OF THE INVENTION

10 The present invention relates generally to a genetic sequence encoding a polypeptide having flavonoid 3', 5'-hydroxylase (F3'5'H) activity and to the use of the genetic sequence and/or its corresponding polypeptide thereof *inter alia* to manipulate color in flowers or parts thereof or in other plant tissue. More particularly, the F3'5'H has the ability to modulate dihydrokaempferol (DHK) metabolism as well as the metabolism of other substrates such as dihydroquercetin (DHQ), naringenin and eriodictyol. Even more particularly, the present invention provides a genetic sequence encoding a polypeptide having F3'5'H activity when expressed in rose or gerbera or botanically related plants. The instant invention further relates to antisense and sense molecules or RNAi-inducing molecules corresponding to all or part of the subject genetic sequence or a transcript thereof as well as to genetically modified plants as well as cut flowers, parts and reproductive tissue from such plants. The present invention further relates to promoters which operate efficiently in plants such as rose, gerbera or botanically related plants.

## 20 DESCRIPTION OF PRIOR ART

Reference to any prior art in this specification is not, and should not be taken as, an acknowledgment or any form of suggestion that this prior art forms part of the common general knowledge in any country.

Bibliographic details of references provided in the subject specification are listed at the end of the specification.

- 2 -

The flower or ornamental plant industry strives to develop new and different varieties of flowers and/or plants. An effective way to create such novel varieties is through the manipulation of flower color. Classical breeding techniques have been used with some success to produce a wide range of colors for almost all of the commercial varieties of flowers and/or plants available today. This approach has been limited, however, by the constraints of a particular species' gene pool and for this reason it is rare for a single species to have the full spectrum of colored varieties. For example, the development of novel colored varieties of plants or plant parts such as flowers, foliage and stems would offer a significant opportunity in both the cut flower and ornamental markets. In the flower or ornamental plant industry, the development of novel colored varieties of major flowering species such as rose, chrysanthemum, tulip, lily, carnation, gerbera, orchid, lisianthus, begonia, torenia, geranium, petunia, nierenbergia, pelargonium, iris, impatiens and cyclamen would be of great interest. A more specific example would be the development of a blue rose or gerbera for the cut flower market.

In addition, the development of novel colored varieties of plant parts such as vegetables, fruits and seeds would offer significant opportunities in agriculture. For example, novel colored seeds would be useful as proprietary tags for plants. Furthermore modifications to flavonoids common to berries or fruits including grapes and apples and their juices including wine have the potential to impart altered style characteristics of value to such fruit and byproduct industries.

Flower color is predominantly due to three types of pigment: flavonoids, carotenoids and betalains. Of the three, the flavonoids are the most common and contribute a range of colors from yellow to red to blue. The flavonoid molecules that make the major contribution to flower color are the anthocyanins, which are glycosylated derivatives of cyanidin and its methylated derivative peonidin, *delphinidin* or *delphinidin-based molecules* and its methylated derivatives petunidin and malvidin and pelargonidin. Anthocyanins are localised in the vacuole of the epidermal cells of petals or the vacuole of the sub epidermal cells of leaves.

The flavonoid pigments are secondary metabolites of the phenylpropanoid pathway. The biosynthetic pathway for the flavonoid pigments (flavonoid pathway) is well established, (Hollon and Cornish, *Plant Cell* 7: 1071-1083, 1995; Mol *et al*, *Trends Plant Sci* 3: 212-217, 1998; Winkel-Shirley, *Plant Physiol* 126: 485-493, 2001a; and Winkel-Shirley, *Plant Physiol* 127: 1399-1404, 2001b) and is shown in Figures 1A and B. Three reactions and enzymes are involved in the conversion of phenylalanine to *p*-coumaroyl-CoA, one of the first key substrates in the flavonoid pathway. The enzymes are phenylalanine ammonia-lyase (PAL), cinnamate 4-hydroxylase (C4H) and 4-coumarate: CoA ligase (4CL). The first committed step in the pathway involves the condensation of three molecules of malonyl-CoA (provided by the action of acetyl CoA carboxylase (ACC) on acetyl CoA and CO<sub>2</sub>) with one molecule of *p*-coumaroyl-CoA. This reaction is catalysed by the enzyme chalcone synthase (CHS). The product of this reaction, 2',4',4'',6', tetrahydroxy-chalcone, is normally rapidly isomerized by the enzyme chalcone flavanone isomerase (CHI) to produce naringenin. Naringenin is subsequently hydroxylated at the 3 position of the central ring by flavanone 3-hydroxylase (F3H) to produce dihydrokaempferol (DHK). 5

The pattern of hydroxylation of the B-ring of dihydrokaempferol (DHK) plays a key role in determining petal color. The B-ring can be hydroxylated at either the 3', or both the 3' and 5' positions, to produce dihydroquercetin (DHQ) or dihydromyricetin (DHM), respectively. Two key enzymes involved in this part of the pathway are flavonoid 3'-hydroxylase and flavonoid 3', 5'-hydroxylase, both of the cytochrome P450 class of enzymes. Cytochrome P450 enzymes are widespread in nature and genes have been isolated and sequenced from vertebrates, insects, yeasts, fungi, bacteria and plants. 10

Flavonoid 3'-hydroxylase (F3'H) is a key enzyme in the flavonoid pathway leading to the cyanidin-based pigments which, in many plant species (for example *Rosa* spp., *Dianthus* spp., *Petunia* spp., begonia, cyclamen, impatiens, morning glory and chrysanthemum), contribute to red and pink flower color. 15

Flavonoid 3', 5'-hydroxylase (F3',5'H) is a key enzyme in the flavonoid pathway leading to the delphinidin-based pigments which, in many plant species (for example, *Petunia* spp., 20

30



- 4 -

*Viola spp.*, *Lisianthus spp.*, *Gentiana spp.*, *Solhya spp.*, *Salvia spp.*, *Clitoria spp.*, *Kennedia spp.*, *Campanula spp.*, *Lavandula spp.*, *Verbena spp.*, *Torenia spp.*, *Delphinium spp.*, *Solanum spp.*, *Cineraria spp.*, *Vitis spp.*, *Babiana stricta*, *Pinus spp.*, *Picea spp.*, *Larix spp.*, *Phaseolus spp.*, *Vaccinium spp.*, *Cyclamen spp.*, *Iris spp.*, *Pelargonium spp.*,  
 5 *Liparicaceae*, *Geranium spp.*, *Pisum spp.*, *Lathyrus spp.*, *Catharanthus spp.*, *Malva spp.*, *Mucuna spp.*, *Vicia spp.*, *Saltpaulia spp.*, *Lagerstroemia spp.*, *Bouchea spp.*, *Plumbago spp.*, *Hypocalyptus spp.*, *Rhododendron spp.*, *Linum spp.*, *Macropitium spp.*, *Hibiscus spp.*, *Hydrangea spp.*, *Cymbidium spp.*, *Millettia spp.*, *Hedysarum spp.*, *Lespedeza spp.*,  
 10 *Asparagus spp.*, *Antigonon spp.*, *Pisum spp.*, *Freesia spp.*, *Brunella spp.*, *Clarkia spp.*, etc.), contribute to purple and blue flower color. Many plant species such as roses, gerberas, chrysanthemums and carnations, do not produce delphinidin-based pigments because they lack a F3'5'H activity.

The next step in the pathway, leading to the production of the colored anthocyanins from  
 15 the dihydroflavonols (DHK, DHQ, DHM), involves dihydroflavonol-4-reductase (DFR) leading to the production of the leucoanthocyanidins. The leucoanthocyanidins are subsequently converted to the anthocyanidins, pelargonidin, cyanidin and *delphinidin* or *delphinidin-based molecules*. These flavonoid molecules are unstable under normal physiological conditions and glycosylation at the 3-position, through the action of  
 20 glycosyltransferases, stabilizes the anthocyanidin molecule thus allowing accumulation of the anthocyanins. In general, the glycosyltransferases transfer the sugar moieties from UDP sugars to the flavonoid molecules and show high specificities for the position of glycosylation and relatively low specificities for the acceptor substrates (Seitz and Hinderer, Anthocyanins. In: *Cell Culture and Somatic Cell Genetics of Plants*. Constabel,  
 25 P. and Vasil, LK. (eds.), Academic Press, New York, USA, 5: 49-76, 1988). Anthocyanins can occur as 3-monosides, 3-biosides and 3-triosides as well as 3, 5-diglycosides and 3, 7-diglycosides associated with the sugars glucose, galactose, rhamnose, arabinose and xylose (Strack and Wray, In: *The Flavonoids - Advances in Research since 1986*. Harborne, J.B. (ed), Chapman and Hall, London, UK, 1-22, 1993).

Glycosyltransferases involved in the stabilisation of the anthocyanidin molecule include UDP glucose: flavonoid 3-glucosyltransferase (3GT), which transfers a glucose moiety from UDP glucose to the 3-O-position of the anthocyanidin molecule to produce anthocyanidin 3-O-glucoside.

In petunia and pansy (amongst others), anthocyanidin 3-O-glucoside are generally glycosylated by another glycosyltransferase, UDP rhamnose: anthocyanidin 3-glucoside rhamnosyltransferase (3RT), which adds a rhamnose group to the 3-O-bound glucose of the anthocyanin molecule to produce the anthocyanidin 3-rutinosides, and once acylated, can be further modified by UDP: glucose anthocyanin 5-glucosyltransferase (5GT). However, in roses (amongst others), the anthocyanidin 3-O-glucosides are generally glycosylated by another glycosyltransferase, UDP: glucose anthocyanin 5-glucosyltransferase (5GT) to produce anthocyanidin 3, 5 diglucosides.

Many anthocyanidin glycosides exist in the form of acylated derivatives. The acyl groups that modify the anthocyanidin glycosides can be divided into two major classes based upon their structure. The aliphatic acyl groups include malonic acid or succinic acid and the aromatic class include the hydroxy cinnamic acids such as *p*-coumaric acid, caffeic acid and ferulic acid and the benzoic acids such as *p*-hydroxybenzoic acid.

Methylation at the 3' and 5' positions of the B-ring of anthocyanidin glycosides can also occur. Methylation of cyanidin-based pigments leads to the production of peonidin. Methylation of the 3' position of delphinidin-based pigments results in the production of petunidin, whilst methylation of the 3' and 5' positions results in malvidin production. Methylation of malvidin can also occur at the 5-O and 7-O positions to produce capensathin (5-O-methyl malvidin) and 5, 7-di-O-methyl malvidin.

In addition to the above modifications, pH of the vacuole or compartment where pigments are localised and copigmentation with other flavonoids such as flavonols and flavones can affect petal color. Flavonols and flavones can also be aromatically acylated (Brouillard and

- 6 -

Dangles, In: *The Flavonoids -Advances in Research since 1986*. Harborne, J.B. (ed), Chapman and Hall, London, UK, 1-22, 1993).

- 5 The ability to control F3'5'H activity, or other enzymes involved in the flavonoid pathway, in flowering plants would provide a means of manipulating the color of plant parts such as petals, fruit, leaves, sepals, seeds etc. Different colored versions of a single cultivar could thereby be generated and in some instances a single species would be able to produce a broader spectrum of colors.
- 10 Two nucleotide sequences (referred to herein as SEQ ID NO:1 and SEQ ID NO:3) encoding petunia F3'5'Hs have been cloned (see International Patent Application No. PCT/AU92/00334 and Holton *et al.*, *Nature*, 366: 276-279, 1993a). These sequences were efficient in modulating 3', 5' hydroxylation of flavonoids in petunia (see International Patent Application No. PCT/AU92/00334 incorporated herein by reference and Holton *et*
- 15 *al.*, 1993a, *supra*), tobacco (see International Patent Application No. PCT/AU92/00334 incorporated herein by reference) and carnations (see International Patent Application No. PCT/AU96/00296 incorporated herein by reference). Surprisingly, however, inclusion of these sequences in standard expression cassettes, did not lead to the production of intact or full-length transcripts as detectable by RNA or Northern blot analysis and consequently 3',
- 20 5'-hydroxylated flavonoids were not produced in roses. There is a need, therefore, to identify further genetic sequences encoding F3'5'Hs which efficiently accumulate and are then able to modulate 3', 5' hydroxylation of flavonoids such as anthocyanins in roses and other key commercial plant species.

- 7 -

## SUMMARY OF THE INVENTION

Throughout this specification, unless the context requires otherwise, the word "comprise", or variations such as "comprises" or "comprising", will be understood to imply the inclusion of a stated element or group of elements or integers but not the exclusion of any other element or integer or group of elements or integers.

Nucleotide and amino acid sequences are referred to by a sequence identifier number (SEQ ID NO:). The SEQ ID NOS: correspond numerically to the sequence identifiers <400>1 (SEQ ID NO:1), <400>2 (SEQ ID NO:2), etc.

Genetic sequences encoding a F3'5'H have been identified and cloned from a number of plant species. The F3'5'H genetic sequences when expressed in rose petal tissue results in a detectable level of delphinidin or delphinidin-based molecules as determined by a chromatographic technique such as thin layer chromatography (TLC) or high performance liquid chromatography (HPLC). Alternatively, or in addition, expression of the genetic sequences in rose petal tissue results in a sufficient level and length of transcript which is capable of being translated to F3'5'H. This is conveniently measured as delphinidin or delphinidin-based molecules, detectable using a chromatographic technique such as TLC or HPLC. The genetic sequences of the present invention permit the modulation of expression of genes encoding this enzyme by, for example, *de novo* expression, over-expression, suppression, antisense inhibition, ribozyme activity, RNAi-induction or methylation-induction. The ability to control F3'5'H synthesis in plants and more specifically in roses or gerberas permits modulation of the composition of individual anthocyanins as well as alteration of relative levels of flavonols and anthocyanins, thereby enabling the manipulation of color of tissues and/or organs of plants such as petals, leaves, seeds, sepals, fruits etc.

Accordingly, one aspect of the present invention provides an isolated nucleic acid molecule comprising a sequence of nucleotides encoding or complementary to a sequence encoding a flavonoid 3', 5' hydroxylase (F3'5'H) or a polypeptide having F3'5'H activity

- 8 -

wherein expression of said nucleic acid molecule in a rose petal tissue results in detectable levels of delphinidin or delphinidin-based molecules as measured by a chromatographic technique.

- 5 Another aspect of the present invention is directed to an isolated nucleic acid molecule comprising a sequence of nucleotides encoding or complementary to a sequence encoding a F3'5'H or a polypeptide having F3'5'H activity wherein expression of said nucleic acid molecule in a rose petal tissue results in a sufficient level and length of transcript which is translated to said F3'5'H as determined by detectable levels of delphinidin or delphinidin-  
10 based molecules as measured by a chromatographic technique.

The isolated nucleic acid molecule of the present invention, therefore, encodes a F3'5'H which is capable of more efficient conversion of DHK to DHM in roses than is the F3'5'H encoded by the nucleotide sequence set forth in SEQ ID NO:1 and SEQ ID NO:3 as  
15 measured by delphinidin production in rose petals.

Efficiency as used herein relates to the capability of the F3'5'H enzyme to convert its substrate DHK or DHQ into DHM in a rose cell (or any cell of a commercially important plant such as gerbera). This conversion provides the plant with a substrate (DHM) for other  
20 enzymes of the flavonoid pathway which are present in the plant to further modify the substrate. This modification may include for example, glycosylation, acylation, rhamnosylation and/or methylation, to produce various anthocyanins which contribute to the production of a range of colors. The modulation of 3',5'-hydroxylated anthocyanins in rose is thereby enabled. Efficiency is conveniently assessed by one or more parameters  
25 selected from: extent of F3'5'H transcription, as determined by the amount of intact F3'5'H mRNA produced (as detected by Northern blot analysis); extent of translation of the F3'5'H mRNA, as determined by the amount of translation product produced; extent of F3'5'H enzyme activity as determined by the production of anthocyanin derivatives of DHQ or DHM including delphinidin or delphinidin-based pigments (as detected by TLC or HPLC);  
30 the extent of effect on flower color.

It has also been surprisingly determined that certain combinations of promoter and *F3'5'H* gene sequences that were functional in carnation and petunia were not functional in rose. Surprisingly, only a particular subset of promoter and *F3'5'H* gene sequence combinations resulted in 3', 5'-hydroxylated flavonoids in rose flowers. These included *F3'5'H* sequences isolated from *Viola spp.*, *Salvia spp.*, *Lavandula spp.* and *Solima spp.* Furthermore, the *Viola F3'5'H* (or pansy *F3'5'H*) sequences were found to result in the highest accumulation of 3', 5'-hydroxylated flavonoids in rose. The novel promoter and *F3'5'H* gene sequence combinations can be employed *inter alia* to modulate the color or flavour or other characteristics of plants or plant parts such as but not limited to flowers, fruits, nuts, roots, stems, leaves or seeds. Thus, the present invention represents a new approach to developing plant varieties having altered color characteristics. Other uses include, for example, the production of novel extracts of *F3'5'H* transformed plants wherein the extract has use, for example, as a flavouring or food additive or health product or beverage or juice or coloring. Beverages may include but are not limited to wines, spirits, teas, coffees, milk and dairy products.

15 In a preferred embodiment, therefore, the present invention provides an isolated nucleic acid molecule comprising a sequence of nucleotides encoding, or complementary to a sequence encoding pansy *F3'5'H*, *salvia F3'5'H*, *lavender F3'5'H*, *kennedia F3'5'H* or *solima F3'5'H* or a functional derivative of the enzyme.

20 The nucleotide sequences encoding the pansy *F3'5'H* (SEQ ID NOS:9 and 11), *salvia F3'5'H* (SEQ ID NOS:13 and 15), *solima F3'5'H* (SEQ ID NOS:17), *lavender F3'5'H* (SEQ ID NOS:31) and *kennedia F3'5'H* (SEQ ID NOS:26) are defined by sequence identifiers indicated in parentheses. A summary of the sequence identifiers is shown in Table 1.

25 Accordingly, another aspect of the present invention provides a nucleic acid molecule comprising a nucleotide sequence or complementary nucleotide sequence substantially as set forth in SEQ ID NOS:9 (pansy) or SEQ ID NOS:11 (pansy) or SEQ ID NOS:13 (*salvia*) or SEQ ID NOS:15 (*salvia*) or SEQ ID NOS:17 (*solima*) or SEQ ID NOS:31 (*lavender*) or SEQ ID NOS:26 (*kennedia*) or having at least about 50% similarity thereto or capable of

30

- 10 -

hybridizing to the sequence set forth in SEQ ID NO:9 or SEQ ID NO:11 or SEQ ID NO:13 or SEQ ID NO:15 or SEQ ID NO:17 or SEQ ID NO:31 or SEQ ID NO:26 under low stringency conditions.

- 5 The amino acid sequences of the preferred F3'5'H enzymes are set forth in SEQ ID NO:10 (pansy) or SEQ ID NO:12 (pansy) or SEQ ID NO:14 (salvia) or SEQ ID NO:16 (salvia) or SEQ ID NO:18 (sollya) or SEQ ID NO:32 (lavender) or SEQ ID NO:27 (kennedia).

- 10 A further aspect of the present invention provides a method for producing a transgenic flowering plant capable of synthesizing a F3'5'H said method comprising stably transforming a cell of a suitable plant with a nucleic acid sequence which comprises a sequence of nucleotides encoding said F3'5'H under conditions permitting the eventual expression of said nucleic acid sequence, regenerating a transgenic plant from the cell and growing said transgenic plant for a time and under conditions sufficient to permit the
- 15 expression of the nucleic acid sequence. The expression of the nucleic acid sequence generally results in a transcription of sufficient level and length to encode a F3'5'H. This is conveniently determined by detectable levels of delphinidin or delphinidin-based molecules as measured by chromatographic techniques such as TLC or HPLC. The transgenic plant may thereby produce a non-indigenous F3'5'H at elevated levels relative to
- 20 the amount expressed in a comparable non-transgenic plant. This generally results in a visually detectable color change in the plant or plant part or preferably in the inflorescence or flowers of said plant.

- 25 Another aspect of the present invention contemplates a method for producing a transgenic plant with reduced F3'5'H activity, said method comprising stably transforming a cell of a suitable plant with a nucleic acid molecule which comprises a sequence of nucleotides encoding or complementary to a sequence encoding a F3'5'H activity, regenerating a transgenic plant from the cell and where necessary growing said transgenic plant under conditions sufficient to permit the expression of the nucleic acid.

30

Yet another aspect of the present invention contemplates a method for producing a genetically modified plant with reduced *R3'5'H* activity, said method comprising altering the *R3'5'H* gene through modification of the indigenous sequences via homologous recombination from an appropriately altered *R3'5'H* gene or derivative or part thereof introduced into the plant cell, and regenerating the genetically modified plant from the cell.

Still another aspect of the present invention contemplates a method for producing a transgenic flowering plant exhibiting altered floral or inflorescence properties, said method comprising stably transforming a cell of a suitable plant with a nucleic acid sequence of the present invention, regenerating a transgenic plant from the cell and growing said transgenic plant for a time and under conditions sufficient to permit the expression of the nucleic acid sequence.

Still a further aspect of the present invention contemplates a method for producing a transgenic flowering plant exhibiting altered floral or inflorescence properties, said method comprising alteration of the *R3'5'H* gene through modification of nucleotide sequences *via* homologous recombination from an appropriately altered *R3'5'H* gene or derivative or part thereof introduced into the plant cell, and regenerating the genetically modified plant from the cell.

Even yet another aspect of the present invention extends to a method for producing a transgenic plant capable of expressing a recombinant gene encoding a *R3'5'H* or part thereof or which carries a nucleic acid sequence which is substantially complementary to all or a part of a mRNA molecule encoding said *R3'5'H*, said method comprising stably transforming a cell of a suitable plant with the isolated nucleic acid molecule comprising a sequence of nucleotides encoding, or complementing to a sequence encoding, a *R3'5'H*, where necessary under conditions permitting the eventual expression of said isolated nucleic acid molecule, and regenerating a transgenic plant from the cell.



- 12 -

Even still another aspect of the present invention extends to all transgenic plants or parts of transgenic plants or progeny of the transgenic plants containing all or part of the nucleic acid sequences of the present invention, or antisense forms thereof and/or any homologs or related forms thereof and, in particular, those transgenic plants which exhibit altered floral or inflorescence properties.

Even still another aspect of the present invention extends to all transgenic plants or parts of transgenic plants or progeny of the transgenic plants containing all or part of the nucleic acid sequences of the present invention, or antisense forms thereof and/or any homologs or related forms thereof and, in particular, those transgenic plants which exhibit altered aerial parts of the plant such as fruit, berries, sepal, bract, petiole, peduncle, ovaries, anthers or stem properties.

Another aspect of the present invention contemplates the use of the extracts from transgenic plants or plant parts transgenic plants or progeny of the transgenic plants containing all or part of the nucleic acid sequences of the present invention and, in particular, the extracts from those transgenic plants when used as a flavouring or food additive or health product or beverage or juice or coloring.

A further aspect of the present invention is directed to recombinant forms of F3'5'H.

Another aspect of the present invention contemplates the use of the genetic sequences described herein in the manufacture of a genetic construct capable of expressing a F3'5'H or down-regulating an indigenous F3'5'H enzyme in a plant.

Yet another aspect of the present invention is directed to a prokaryotic or eukaryotic organism carrying a genetic sequence encoding a F3'5'H extrachromasomally in plasmid form.

Still another aspect of the present invention extends to a recombinant polypeptide comprising a sequence of amino acids substantially as set forth in SEQ ID NO:10 or SEQ ID NO:12 or SEQ ID NO:14 or SEQ ID NO:16 or SEQ ID NO:18 or SEQ ID NO:32 or SBQ ID NO:27 or an amino acid sequence having at least about 50% similarity to SBQ ID NO:10 or SEQ ID NO:12 or SEQ ID NO:14 or SEQ ID NO:16 or SBQ ID NO:18 or SBQ ID NO:32 or SBQ ID NO:27 or a derivative of said polypeptide.

The present invention further provides promoters which operate efficiently in plants such as rose and gerbera or botanically related plants. Such promoters include a rose CHS promoter, chrysanthemum CHS promoter and a CaMV 35S promoter.

A summary of sequence identifiers used throughout the subject specification is provided in Table 1:

TABLE 1

Summary of sequence identifiers

SEQ ID	NAME	SPECIES	TYPE	DESCRIPTION
1	pet171.mt	Petunia hybrida	nucleotide	F3'5'H cDNA
2	pet171.aa	Petunia hybrida	amino acid	translation of F3'5'H cDNA
3	pet172.mt	Petunia hybrida	nucleotide	F3'5'H cDNA
4	pet172.aa	Petunia hybrida	amino acid	translation of F3'5'H cDNA
5	RoseCHS promoter	Rosa hybrida	nucleotide	promoter fragment
6	D8 oligo#2	Petunia hybrida	nucleotide	oligonucleotide
7	D8 oligo #4	Petunia hybrida	nucleotide	oligonucleotide
8	chrysanthCHSATG	chrysanthemum	nucleotide	oligonucleotide
9	BP#18.mt	Viola spp.	nucleotide	F3'5'H cDNA
10	BP#18.aa	Viola spp.	amino acid	translation of F3'5'H cDNA
11	BP#40.mt	Viola spp.	nucleotide	F3'5'H cDNA
12	BP#40.aa	Viola spp.	amino acid	translation of F3'5'H cDNA
13	Sal#2.mt	Salvia spp.	nucleotide	F3'5'H cDNA

- 14 -

SEQ ID NO:	NAME	SPECIES	TYPE OF SEQ	DESCRIPTION
14	Sal#2.aa	Salvia spp.	amino acid	translation of F3'5'H cDNA
15	Sal#47.nt	Salvia spp.	nucleotide	F3'5'H cDNA
16	Sal#47.aa	Salvia spp.	amino acid	translation of F3'5'H cDNA
17	Soll#5.nt	Solhya spp.	nucleotide	F3'5'H cDNA
18	Soll#5.aa	Solhya spp.	amino acid	translation of F3'5'H cDNA
19	FLS-Nco	Petunia hybrida	nucleotide	oligonucleotide
20	BpeaHF2.nt	Clitoria ternatea	nucleotide	F3'5'H cDNA
21	BpeaHF2.aa	Clitoria ternatea	amino acid	translation of F3'5'H cDNA
22	Gen#48.nt	Gentiana triflora	nucleotide	F3'5'H cDNA
23	Gen#48.aa	Gentiana triflora	amino acid	translation of F3'5'H cDNA
24	PaiD8 5'	Petunia hybrida	nucleotide	oligonucleotide
25	Bpea primer	Clitoria ternatea	nucleotide	oligonucleotide
26	Kenn#31.nt	Kennedia spp.	nucleotide	F3'5'H cDNA
27	Kenn#31.aa	Kennedia spp.	amino acid	translation of F3'5'H cDNA
28	chrysCHS.nt	chrysanthemum	nucleotide	CHS cDNA
29	chrysCHS.aa	chrysanthemum	amino acid	translation of CHS cDNA
30	chrysCHS promoter	chrysanthemum	nucleotide	promoter fragment
31	LBG.nt	Lavandula nil	nucleotide	F3'5'H cDNA
32	LBG.aa	Lavandula nil	amino acid	translation of F3'5'H cDNA

## BRIEF DESCRIPTION OF THE FIGURES

Figures 1A and 1B are schematic representations of the biosynthesis pathway for the flavonoid pigments. Figure 1A illustrates the general production of the anthocyanidin 3-glucosides that occur in most plants that produce anthocyanins. Figure 1B represents further modifications of anthocyanins that occur in pecunia. Enzymes involved in the pathway have been indicated as follows: PAL = Phenylalanine ammonia-lyase; C4H = Cinnamate 4-hydroxylase; 4CL = 4-coumarate: CoA-ligase; CHS = Chalcone synthase; CHI = Chalcone flavanone isomerase; F3H = Flavanone 3-hydroxylase; DFR = Dihydroflavonol-4-reductase; ANS = Anthocyanidin synthase; 3GT = UDP-glucose: Flavonoid 3-O-glucosyltransferase; 3RT = UDP rhamnose: anthocyanidin 3-glucoside rhamnosyltransferase; AR-AT = Anthocyanidin-rutinoside acyltransferase, 5GT = Anthocyanin 5-glucosyltransferase; 3' OMT = Anthocyanin 3'-O-methyltransferase, 3'5' OMT = Anthocyanin 3', 5' O-methyltransferase. Other abbreviations include: DHK = dihydrokaempferol, DHQ = dihydroquercetin, DHM = dihydromyricetin,

TABLE 2: Descriptions of the abbreviations used in Figures 2 to 52

ABBREVIATION	DESCRIPTION
Amp	ampicillin resistance gene which confers resistance to the antibiotic ampicillin
ColE1ori	plasmid origin of replication
fl ori (+)	fl filamentous phage origin of replication
GenR	gentamycin resistance gene which confers resistance to the antibiotic gentamycin
LB	left border of the T-DNA
npIII	the neomycin phosphotransferase III gene which confers resistance to the antibiotic kanamycin
ori PR1	plasmid origin of replication
ori 322	plasmid origin of replication
PACYC ori	modified replicon from PACYC184 from <i>E. coli</i>

- 16 -

pVS1	a broad host range origin of replication from a plasmid from <i>Pseudomonas aeruginosa</i>
rev	approximate location of the M13 reverse primer site used in sequence analysis
RB	right border of the T-DNA
TetR	tetracycline resistance gene which confers resistance to the antibiotic tetracycline
-20	approximate location of the M13 -20 primer site used in sequence analysis
RK2	broad host range Gram-negative plasmid RK2 origin

Figure 2 is a diagrammatic representation of the plasmid pCGP602, pCGP601 and pCGP176 containing petunia *F3'5'H petHf1* cDNA clones from *P. hybrida* cv. *OGB*. The petunia *F3'5'H petHf1* fragment was used in the preparation of constructs containing the petunia *F3'5'H* cDNA clone. <sup>32</sup>P-labelled fragments of the 1.6 kb *BspHI/FspI* fragment were used to probe petal cDNA libraries. Selected restriction endonuclease sites are marked. Refer to Table 2 and Table 4 for a description of the abbreviations.

Figure 3 is a diagrammatic representation of the plasmid pCGP175 containing the petunia *F3'5'H petHf2* cDNA clone from *P. hybrida* cv. *OGB*. Selected restriction endonuclease sites are marked. Refer to Table 2 and Table 4 for a description of the abbreviations.

Figure 4 is a diagrammatic representation of the plasmid pCGP1303 containing a subclone of the petunia *F3'5'H petHf1* cDNA clone from pCGP601. The construction of pCGP1303 is described in Example 4. Selected restriction endonuclease sites are marked. Refer to Table 2 and Table 4 for a description of the abbreviations.

Figure 5 is a diagrammatic representation of the binary plasmid pCGP1452. The *AmCHS* 5': *petHf1*: *petD8* 3' gene from pCGP485 was cloned into the binary vector pWTT2132 (DNAP) in a tandem orientation with the chimaeric *SuRB* gene. The construction of pCGP1452 is described in Example 4. Selected restriction endonuclease sites are marked. Refer to Table 2 and Table 4 for a description of the abbreviations.

Figure 6 is a diagrammatic representation of the binary plasmid pWTT2132 (DNAP) containing the 35S 5' : SURB selectable marker gene and a multi-cloning site. A description of pWTT2132 is given in Example 4. Selected restriction endonuclease sites are marked. Refer to Table 2 and Table 4 for a description of the abbreviations.

Figure 7 is a diagrammatic representation of the plasmid pCGP725. The *AmCHS 5' : petH1 : petD 3'* gene from pCGP485 was cloned into pBluescript II (KS (+) vector. The construction of pCGP725 is described in Example 4. Selected restriction endonuclease sites are marked. Refer to Table 2 and Table 4 for a description of the abbreviations.

Figure 8 is a diagrammatic representation of the binary plasmid pCGP1453. The *Mac : petH1 : mas 3'* gene from pCGP628 was cloned into the binary vector pWTT2132 (DNAP) in a tandem orientation with the chimeric *SURB* gene. The construction of pCGP1453 is described in Example 4. Selected restriction endonuclease sites are marked. Refer to Table 2 and Table 4 for a description of the abbreviations.

Figure 9 is a diagrammatic representation of the binary plasmid pCGP1457. The *petD 5' : petH1 : petD 3'* gene from pCGP1107 was cloned into the binary vector pWTT2132 (DNAP) in a tandem orientation with the chimeric *SURB* gene. The construction of pCGP1457 is described in Example 4. Selected restriction endonuclease sites are marked. Refer to Table 2 and Table 4 for a description of the abbreviations.

Figure 10 is a diagrammatic representation of the binary plasmid pCGP1461. The *shortpetLS 5' : petH1 : petLS 3'* gene from pCGP497 was cloned into the binary vector pWTT2132 (DNAP) in a tandem orientation with the chimeric *SURB* gene. The construction of pCGP1461 is described in Example 4. Selected restriction endonuclease sites are marked. Refer to Table 2 and Table 4 for a description of the abbreviations.

Figure 11 is a diagrammatic representation of the binary plasmid pCGP1616. The *petRT 5' : petH1 : nos 3'* gene from pCGP846 was cloned into the binary vector pWTT2132 (DNAP) in a tandem orientation with the chimeric *SURB* gene. The construction of

pCGP1616 is described in Example 4. Selected restriction endonuclease sites are marked. Refer to Table 2 and Table 4 for a description of the abbreviations.

Figure 12 is a diagrammatic representation of the binary plasmid pCGP1623. The *mas/35S: petHf1: ocs* 3' gene from pCGP1619 was cloned into the binary vector pWTT2132 (DNAP) in a tandem orientation with the chimaeric *SuRB* gene. The construction of pCGP1623 is described in Example 4. Selected restriction endonuclease sites are marked. Refer to Table 2 and Table 4 for a description of the abbreviations.

Figure 13 is a diagrammatic representation of the binary plasmid pCGP1638. The *CaMV 35S: petHf1: nos* 3' gene from pCGP1636 was cloned into the binary vector pWTT2132 (DNAP) in a tandem orientation with the chimaeric *SuRB* gene. The construction of pCGP1636 is described in Example 4. Selected restriction endonuclease sites are marked. Refer to Table 2 and Table 4 for a description of the abbreviations.

Figure 14 is a diagrammatic representation of the binary plasmid pCGP1860. The *RoseCHS 5': petHf1: nos* 3' gene from pCGP200 was cloned into the binary vector pWTT2132 (DNAP) in a tandem orientation with the chimaeric *SuRB* gene. The construction of pCGP1860 is described in Example 4. Selected restriction endonuclease sites are marked. Refer to Table 2 and Table 4 for a description of the abbreviations.

Figure 15 is a diagrammatic representation of the binary plasmid pCGP2123. The *CaMV35S: petHf2: ocs* 3' gene from pCGP2109 was cloned into the binary vector pCGP1988 in a tandem orientation with the chimaeric *SuRB* gene. The construction of pCGP2123 is described in Example 4. Selected restriction endonuclease sites are marked. Refer to Table 2 and Table 4 for a description of the abbreviations.

Figure 16 is a diagrammatic representation of the binary plasmid pCGP1988. The multi-cloning site of the binary vector pWTT2132 (DNAP) was replaced with the multi-cloning site from pNEB193 (New England Biolabs). The construction of pCGP1988 is described in

Example 4. Selected restriction endonuclease sites are marked. Refer to Table 2 and Table 4 for a description of the abbreviations.

Figure 17 is a diagrammatic representation of the plasmid pCGP2105. The 35S 5' : oc 3' expression cassette with multiple restriction endonuclease sites between the promoter and terminator fragments is in a pBluescript SK (+) vector backbone. The construction of pCGP2105 is described in Example 4. Selected restriction endonuclease sites are marked. Refer to Table 2 and Table 4 for a description of the abbreviations.

Figure 18 is a diagrammatic representation of the binary plasmid pCGP1307. The *petD* 5' : *GUS* : *petD* 3' gene from pCGP1106 was cloned into the binary vector pCGN1548 in a tandem orientation to the chimeric *npII* selectable marker gene. The construction of pCGP1307 is described in Example 6. Selected restriction endonuclease sites are marked. Refer to Table 2 and Table 4 for a description of the abbreviations.

Figure 19 is a diagrammatic representation of the binary plasmid pCGP1506. The *longpetLS* 5' : *GUS* : *petLS* 3' gene from pCGP496 was cloned into the binary vector pBIN19 in a tandem orientation to the chimeric *npII* selectable marker gene. The construction of pCGP1506 is described in Example 6. Refer to Table 2 and Table 4 for a description of the abbreviations.

Figure 20 is a diagrammatic representation of the binary plasmid pCGP1626. The *ChrysCHS* 5' : *GUS* : *nos* 3' gene from pCGP1622 was cloned into the binary vector pWTT2132 in a tandem orientation with the chimeric *SkrB* gene. The construction of pCGP1626 is described in Example 6. Selected restriction endonuclease sites are marked. Refer to Table 2 and Table 4 for a description of the abbreviations.

Figure 21 is a diagrammatic representation of the binary plasmid pCGP1641. The *peRT* 5' : *GUS* : *peRT* 3' gene from pCGP1628 was cloned into the binary vector pWTT2132 in a tandem orientation with the chimeric *SkrB* gene. The construction of pCGP1641 is



- 20 -

described in Example 6. Selected restriction endonuclease sites are marked. Refer to Table 2 and Table 4 for a description of the abbreviations.

5 Figure 22 is a diagrammatic representation of the binary plasmid pCGP1861. The *RoseCHS* 5': *GUS: nos* 3' gene from pCGP197 was cloned into the binary vector pWTT2132 in a tandem orientation with the chimaeric *SuRB* gene. The construction of pCGP1861 is described in Example 6. Refer to Table 2 and Table 4 for a description of the abbreviations.

10 Figure 23 is a diagrammatic representation of the binary plasmid pCGP1953. The *AmCHS* 5': *GUS: petD8* 3' gene from pCGP1952 was cloned into the binary vector pWTT2132 in a tandem orientation with the chimaeric *SuRB* gene. The construction of pCGP1953 is described in Example 6. Refer to Table 2 and Table 4 for a description of the abbreviations.

15 Figure 24 is a diagrammatic representation of the binary plasmid pWTT2084 (DNAP) containing a 35S 5': *GUS: ocs* 3' gene in a convergent orientation to the chimaeric *SuRB* selectable marker gene. A description of pWTT2084 is given in Example 6. Refer to Table 2 and Table 4 for a description of the abbreviations.

20 Figure 25 is a diagrammatic representation of the plasmid pCGP1959 containing the *F3'5'H BP#18* cDNA clone from *Viola spp.* cv Black Pansy in a pBluescript SK II (+) backbone. A description of pCGP1959 is given in Example 7. Selected restriction endonuclease sites are marked. Refer to Table 2 and Table 4 for a description of the abbreviations.

25 Figure 26 is a diagrammatic representation of the plasmid pCGP1961 containing the *F3'5'H BP#40* cDNA clone from *Viola spp.* cv Black Pansy in a pBluescript SK II (+) backbone. A description of pCGP1961 is given in Example 7. Selected restriction endonuclease sites are marked. Refer to Table 2 and Table 4 for a description of the abbreviations.

30

Figure 27 is a diagrammatic representation of the binary plasmid pCGP1972. The *AmCHS* 5' : BP#18: *petD8* 3' gene from pCGP1970 was cloned into the binary vector pWTT2132 (DNAP) in a tandem orientation with the chimeric *SuRB* gene. The construction of pCGP1972 is described in Example 7. Selected restriction endonuclease sites are marked. Refer to Table 2 and Table 4 for a description of the abbreviations.

Figure 28 is a diagrammatic representation of the binary plasmid pCGP1973. The *AmCHS* 5' : BP#40: *petD8* 3' gene from pCGP1971 was cloned into the binary vector pWTT2132 (DNAP) in a tandem orientation with the chimeric *SuRB* gene. The construction of pCGP1973 is described in Example 7. Selected restriction endonuclease sites are marked. Refer to Table 2 and Table 4 for a description of the abbreviations.

Figure 29 is a diagrammatic representation of the binary plasmid pCGP1967. The *CamV* 35S: BP#18: *ocs* 3' gene from pCGP1965 was cloned into the binary vector pWTT2132 (DNAP) in a tandem orientation with the chimeric *SuRB* gene. The construction of pCGP1967 is described in Example 7. Selected restriction endonuclease sites are marked. Refer to Table 2 and Table 4 for a description of the abbreviations.

Figure 30 is a diagrammatic representation of the binary plasmid pCGP1969. The *CamV* 35S: BP#40: *ocs* 3' gene from pCGP1966 was cloned into the binary vector pWTT2132 (DNAP) in a tandem orientation with the chimeric *SuRB* gene. The construction of pCGP1969 is described in Example 7. Selected restriction endonuclease sites are marked. Refer to Table 2 and Table 4 for a description of the abbreviations.

Figure 31 is a diagrammatic representation of the plasmid pCGP1995 containing the 3' 5' *H Sal#2* cDNA clone from *Salvia* spp. in a pBluescript SK II (+) backbone. A description of pCGP1995 is given in Example 7. Selected restriction endonuclease sites are marked. Refer to Table 2 and Table 4 for a description of the abbreviations.

- 22 -

Figure 32 is a diagrammatic representation of the plasmid pCGP1999 containing the *F3'5'H Sal#47* cDNA clone from *Salvia spp* in a pBluescript SK II (+) backbone. A description of pCGP1999 is given in Example 7. Selected restriction endonuclease sites are marked. Refer to Table 2 and Table 4 for a description of the abbreviations.

Figure 33 is a diagrammatic representation of the binary plasmid pCGP2121. The *AmCHS* 5': *Sal#2: petD8* 3' gene from pCGP2116 was cloned into the binary vector pCGP1988 in a tandem orientation with the chimaeric *SuRB* gene. The construction of pCGP2121 is described in Example 7. Selected restriction endonuclease sites are marked. Refer to Table 2 and Table 4 for a description of the abbreviations.

Figure 34 is a diagrammatic representation of the binary plasmid pCGP2122. The *AmCHS* 5': *Sal#47: petD8* 3' gene from pCGP2117 was cloned into the binary vector pCGP1988 in a tandem orientation with the chimaeric *SuRB* gene. The construction of pCGP2122 is described in Example 7. Selected restriction endonuclease sites are marked. Refer to Table 2 and Table 4 for a description of the abbreviations.

Figure 35 is a diagrammatic representation of the binary plasmid pCGP2120. The *CaMV* 35S:*Sal#2:ocs* 3' gene from pCGP2112 was cloned into the binary vector pCGP1988 in a tandem orientation with the chimaeric *SuRB* gene. The construction of pCGP2120 is described in Example 7. Selected restriction endonuclease sites are marked. Refer to Table 2 and Table 4 for a description of the abbreviations.

Figure 36 is a diagrammatic representation of the binary plasmid pCGP2119. The *CaMV* 35S:*Sal#47:ocs* 3' gene from pCGP2111 was cloned into the binary vector pCGP1988 in a tandem orientation with the chimaeric *SuRB* gene. The construction of pCGP2119 is described in Example 7. Selected restriction endonuclease sites are marked. Refer to Table 2 and Table 4 for a description of the abbreviations.

Figure 37 is a diagrammatic representation of the plasmid pCGP2110 containing the *R3.5.H.Soli#5* cDNA clone from *Solva* spp. in a pBluescript SK II (+) backbone. A description of pCGP2110 is given in Example 7. Selected restriction endonuclease sites are marked. Refer to Table 2 and Table 4 for a description of the abbreviations.

Figure 38 is a diagrammatic representation of the binary plasmid pCGP2130. The *AmCHS* 5' : *Soli#5:peid8* 3' gene from pCGP2128 was cloned into the binary vector pCGP1988 in a tandem orientation with the chimaeric *SuRB* gene. The construction of pCGP2130 is described in Example 7. Refer to Table 2 and Table 4 for a description of the abbreviations.

Figure 39 is a diagrammatic representation of the binary plasmid pCGP2131. The *CamV* 35S: *Soli#5:ocv* 3' gene from pCGP2129 was cloned into the binary vector pCGP1988 in a tandem orientation with the chimaeric *SuRB* gene. The construction of pCGP2131 is described in Example 7. Refer to Table 2 and Table 4 for a description of the abbreviations.

Figure 40 is a diagrammatic representation of the plasmid pCGP2231 containing the *R3.5.H.Kenn#31* cDNA clone from *Kennedia* spp. in a pBluescript SK II (+) backbone. A description of pCGP2231 is given in Example 7. Selected restriction endonuclease sites are marked. Refer to Table 2 and Table 4 for a description of the abbreviations.

Figure 41 is a diagrammatic representation of the binary plasmid pCGP2256. The *AmCHS* 5' : *Kenn#31:peid8* 3' gene from pCGP2242 was cloned into the binary vector pCGP1988 in a tandem orientation with the chimaeric *SuRB* gene. The construction of pCGP2256 is described in Example 7. Refer to Table 2 and Table 4 for a description of the abbreviations.

- 24 -

Figure 42 is a diagrammatic representation of the binary plasmid pCGP2252. The *CaMV* 35S: *Kenn#31:ocs* 3' gene from pCGP2236 was cloned into the binary vector pCGP1988 in a tandem orientation with the chimaeric *SuRB* gene. The construction of pCGP2252 is described in Example 7. Refer to Table 2 and Table 4 for a description of the abbreviations.

Figure 43 is a diagrammatic representation of the plasmid pBHF2F containing the full-length *F3'5'H BpeaHF2* cDNA clone from *Clitoria ternatea* in a pBluescript SK II (+) backbone. A description of pBHF2F is given in Example 7. Selected restriction endonuclease sites are marked. Refer to Table 2 and Table 4 for a description of the abbreviations.

Figure 44 is a diagrammatic representation of the binary plasmid pCGP2135. The *AmCHS* 5': *BpeaHF2: pstD8* 3' gene from pCGP2133 was cloned into the binary vector pCGP1988 in a tandem orientation with the chimaeric *SuRB* gene. The construction of pCGP2135 is described in Example 7. Selected restriction endonuclease sites are marked. Refer to Table 2 and Table 4 for a description of the abbreviations.

Figure 45 is a diagrammatic representation of the binary plasmid pBEBF5. The *eCaMV* 35S: *BpeaHF2: nos* 3' gene was constructed by replacing the *GUS* fragment from pBE2113-GUSs with the *Clitoria F3'5'H BpeaHF2* cDNA clone from pBHF2F. The construction of pBEBF5 is described in Example 7. Refer to Table 2 and Table 4 for a description of the abbreviations.

Figure 46 is a diagrammatic representation of the binary plasmid pCGP2134. The *CaMV* 35S: *BpeaHF2: ocs* 3' gene from pCGP2132 was cloned into the binary vector pCGP1988 in a tandem orientation with the chimaeric *SuRB* gene. The construction of pCGP2134 is described in Example 7. Refer to Table 2 and Table 4 for a description of the abbreviations.

30

Figure 47 is a diagrammatic representation of the plasmid pG48 containing the *F3'5'H* *Gem#48* cDNA clone from *Gentiana triflora* in a pBluescript SK II (+) backbone. A description of pG48 is given in Example 7. Selected restriction endonuclease sites are marked. Refer to Table 2 and Table 4 for a description of the abbreviations.

Figure 48 is a diagrammatic representation of the binary plasmid pCGP1498. The *AmCHS* 5' : *Gem#48: petD8* 3' gene from pCGP1496 was cloned into the binary vector pCGP1988 in a tandem orientation with the chimeric *SuRB* gene. The construction of pCGP1498 is described in Example 7. Selected restriction endonuclease sites are marked. Refer to Table 2 and Table 4 for a description of the abbreviations.

Figure 49 is a diagrammatic representation of the binary plasmid pBHGHRF48. The *eCAMV* 35S: *Gem#48: nos* 3' gene was constructed by replacing the *GUS* fragment from pBE2113-35S with the *Gentiana F3'5'H Gem#48* cDNA clone from pG48. The construction of pBHGHRF48 is described in Example 7. Selected restriction endonuclease sites are marked. Refer to Table 2 and Table 4 for a description of the abbreviations.

Figure 50 is a diagrammatic representation of the binary plasmid pCGP1982. The *CAMV* 35S: *Gem#48:ocs* 3' gene from pCGP1981 was cloned into the binary vector pWTT2132 (DNAP) in a tandem orientation with the chimeric *SuRB* gene. The construction of pCGP1982 is described in Example 7. Refer to Table 2 and Table 4 for a description of the abbreviations.

Figure 51 is a diagrammatic representation of the plasmid pLRFH8 containing the *F3'5'H* *LBG* cDNA clone from *Lavandula nil* in a pBluescript SK II (+) backbone. A description of pLRFH8 is given in Example 7. Selected restriction endonuclease sites are marked. Refer to Table 2 and Table 4 for a description of the abbreviations.

- 26 -

Figure 52 is a diagrammatic representation of the binary plasmid pBELF8. The *sCaMV* 35S: LBG: *nos* 3' gene was constructed by replacing the *GUS* fragment from pBE2113-GUS<sub>s</sub> with the *Lavandula F3'5'H LBG* cDNA clone from pLHF8. The construction of pBELF8 is described in Example 7. Refer to Table 2 and Table 4 for a description of the  
5 abbreviations.

## DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

5 In accordance with the present invention, genetic sequences encoding polypeptides having R3'5'H activity have been identified, cloned and assessed. The recombinant genetic sequences of the present invention permit the modulation of expression of genes encoding this enzyme by, for example, *de novo* expression, over-expression, sense suppression, antisense inhibition, ribozyme, minizyme and DNAzyme activity, RNAi-induction or methylation-induction or other transcriptional or post-transcriptional silencing activities. RNAi-induction includes genetic molecules such as hairpin, short double stranded DNA or RNA, and partially double stranded DNAs or RNAs with one or two single stranded nucleotide overhangs. The ability to control R3'5'H synthesis in plants permits modulation of the composition of individual anthocyanins as well as alteration of relative levels of flavonols and anthocyanins, thereby enabling the manipulation of petal color. Moreover, the present invention extends to plants and reproductive or vegetative parts thereof including flowers, fruits, seeds, vegetables, leaves, stems and the like. The term "transgenic" also includes progeny plants and plants from subsequent genetics and/or crosses thereof from the primary transgenic plants.

20 Accordingly, one aspect of the present invention provides an isolated nucleic acid molecule comprising a sequence of nucleotides encoding or complementary to a sequence encoding a R3'5'H or a polypeptide having R3'5'H activity wherein expression of said nucleic acid molecule in a rose petal tissue results in detectable levels of delphinidin or

25 delphinidin-based molecules as measured by a chromatographic technique.

Another aspect of the present invention is directed to an isolated nucleic acid molecule comprising a sequence of nucleotides encoding or complementary to a sequence encoding a R3'5'H or a polypeptide having R3'5'H activity wherein expression of said nucleic acid molecule in a rose petal tissue results in a sufficient level and length of transcript which is translated to said R3'5'H as determined by detectable levels of delphinidin or delphinidin-based molecules as measured by a chromatographic technique.



- 28 -

A further aspect of the present invention is directed to an isolated nucleic acid molecule comprising a sequence of nucleotides encoding or complementary to a sequence encoding a F3'5'H or a polypeptide having F3'5'H activity wherein expression of said nucleic acid molecule in a rose petal tissue results a full-length transcript which is detectable by Northern blot analysis of total RNA isolated from rose petals.

The present invention is described and exemplified herein by reference to the identification, cloning and manipulation of genetic sequences encoding a F3'5'H which acts on DHK as well as DHQ. Preferably, the F3'5'H enzyme is a pansy, salvia, sollya lavender or kennedia F3'5'H. The F3'5'H enzyme may also be considered to include a polypeptide or protein having a F3'5'H activity or F3'5'H-like activity. The latter encompasses derivatives having altered F3'5'H activities.

A preferred aspect of the present invention, therefore, is directed to an isolated nucleic acid molecule comprising a sequence of nucleotides encoding, or complementary to a sequence encoding a F3'5'H or a functional mutant, derivative, part, fragment, homolog or analog thereof wherein the nucleic acid molecule is characterized by the following:

- (i) the F3'5'H transcript in rose petal tissue is of sufficient level and size to encode a F3'5'H resulting in detectable delphinidin or delphinidin-based molecules in the rose petal tissue as measured by a chromatographic procedure (eg. TLC or HPLC);
- (ii) the F3'5'H transcript in rose petal tissue is full-length and detected by Northern blot analysis of total RNA isolated from rose petal tissue
- (iii) the F3'5'H in rose petal tissue results in detectable delphinidin or delphinidin-based molecules as measured by a chromatographic procedure (eg. TLC or HPLC); and/or
- (iv) the F3'5'H results in a visual color change in rose petal tissue.

The term delphinidin-based pigments includes the anthocyanidin, delphinidin or any derivatives thereof including but not limited to glycosylated, acylated, methylated or other modified forms. Methylated forms of delphinidin include but are not limited to the anthocyanidin petunidin (methylated at the 3'-position), malvidin (methylated at the 3' and 5' position), 5-O methyl malvidin (methylated at the 5, 3' and 5' positions), 5, 7-O dimethyl malvidin (methylated at the 5, 7, 3' and 5' positions). The methylated anthocyanidins can also be modified by glycosylation and acylation. The term anthocyanins defines glycosylated forms of the respective anthocyanidins.

10 By the term "nucleic acid molecule" is meant a genetic sequence in a non-naturally occurring condition. Generally, this means isolated away from its natural state or synthesized or derived in a non-naturally-occurring environment. More specifically, it includes nucleic acid molecules formed or maintained *in vitro*, including genomic DNA fragments recombinant or synthetic molecules and nucleic acids in combination with 15 heterologous nucleic acids. It also extends to the genomic DNA or cDNA or part thereof encoding R3'SH or a part thereof in reverse orientation relative to its own or another promoter. It further extends to naturally occurring sequences following at least a partial purification relative to other nucleic acid sequences.

20 The term "genetic sequences" is used herein in its most general sense and encompasses any contiguous series of nucleotide bases specifying directly, or via a complementary series of bases, a sequence of amino acids in a R3'SH enzyme. Such a sequence of amino acids may constitute a full-length R3'SH such as is set forth in SEQ ID NO: 10 (pansy) or 25 SEQ ID NO: 12 (pansy) or SEQ ID NO: 14 (salvia) or SEQ ID NO: 16 (salvia) or SEQ ID NO: 18 (soya) or SEQ ID NO: 32 (lavender) or SEQ ID NO: 27 (kennedia) or an active truncated form thereof or may correspond to a particular region such as an N-terminal, C-terminal or internal portion of the enzyme. A genetic sequence may also be referred to as a sequence of nucleotides or a nucleotide sequence and includes a recombinant fusion of two or more sequences.

- 30 -

In accordance with the above aspects of the present invention there is provided a nucleic acid molecule comprising a nucleotide sequence or complementary nucleotide sequence substantially as set forth in SEQ ID NO:9 (pansy) or SEQ ID NO:11 (pansy) or SEQ ID NO:13 (salvia) or SEQ ID NO:15 (salvia) or SEQ ID NO:17 (sollya) or SEQ ID NO:31 (lavender) or SEQ ID NO:26 (kenmedia) or having at least about 50% similarity thereto or capable of hybridizing to the sequence set forth in SEQ ID NO:9 or SEQ ID NO:11 or SEQ ID NO:13 or SEQ ID NO:15 or SEQ ID NO:17 or SEQ ID NO:31 or SEQ ID NO:26 under low stringency conditions.

Table 1 provides a summary of the sequence identifiers.

Alternative percentage similarities and identities (at the nucleotide or amino acid level) encompassed by the present invention include at least about 60% or at least about 65% or at least about 70% or at least about 75% or at least about 80% or at least about 85% or at least about 90% or above, such as about 95% or about 96% or about 97% or about 98% or about 99%, such as at least about 60%, 61%, 62%, 63%, 64%, 65%, 66%, 67%, 68%, 69%, 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, 100%.

In a particularly preferred embodiment, there is provided an isolated nucleic acid molecule comprising a nucleotide sequence or complementary nucleotide sequence substantially as set forth in SEQ ID NO:9 (pansy) or SEQ ID NO:11 (pansy) or SEQ ID NO:13 (salvia) or SEQ ID NO:15 (salvia) or SEQ ID NO:17 (sollya) or SEQ ID NO:31 (lavender) or SEQ ID NO:26 (kenmedia) or having at least about 50% similarity thereto or capable of hybridizing to the sequence set forth in SEQ ID NO:1 (petunia) or SEQ ID NO:3 (petunia) or complementary strands of either under low stringency conditions, wherein said nucleotide sequence encodes a polypeptide having a R3'5'H activity.

For the purposes of determining the level of stringency to define nucleic acid molecules capable of hybridizing to SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:9 or SEQ ID NO:11

or SEQ ID NO:13 or SEQ ID NO:15 or SEQ ID NO:17 or SEQ ID NO:31 or SEQ ID NO:26 reference herein to a low stringency includes and encompasses from at least about 0% to at least about 15% v/v formamide and from at least about 1M to at least about 2 M salt for hybridization, and at least about 1 M to at least about 2 M salt for washing conditions. Generally, low stringency is from about 25-30°C to about 42°C. The temperature may be altered and higher temperatures used to replace the inclusion of formamide and/or to give alternative stringency conditions. Alternative stringency conditions may be applied where necessary, such as medium stringency, which includes and encompasses from at least about 16% v/v to at least about 30% v/v formamide and from at least about 0.5 M to at least about 0.9 M salt for hybridization, and at least about 0.5 M to at least about 0.9 M salt for washing conditions, or high stringency, which includes and encompasses from at least about 31% v/v to at least about 50% v/v formamide and from at least about 0.01 M to at least about 0.15 M salt for hybridization, and at least about 0.01 M to at least about 0.15 M salt for washing conditions. In general, washing is carried out  $T_m = 69.3 + 0.41 (G+C)\%$  (Marmur and Doty, *J. Mol. Biol.* 5: 109, 1962). However, the  $T_m$  of a duplex DNA decreases by 1°C with every increase of 1% in the number of mismatch base pairs (Borner and Lakey, *Eur. J. Biochem.* 46: 83, 1974). Formamide is optional in these hybridization conditions. Accordingly, particularly preferred levels of stringency are defined as follows: low stringency is 6 x SSC buffer, 1.0% w/v SDS at 25-42°C; a moderate stringency is 2 x SSC buffer, 1.0% w/v SDS at a temperature in the range 20°C to 65°C; high stringency is 0.1 x SSC buffer, 0.1% w/v SDS at a temperature of at least 65°C.

Another aspect of the present invention provides a nucleic acid molecule comprising a sequence of nucleotides encoding or complementary to a sequence encoding an amino acid sequence substantially as set forth in SEQ ID NO:10 or SEQ ID NO:12 or SEQ ID NO:14 or SEQ ID NO:16 or SEQ ID NO:18 or SEQ ID NO:32 or SEQ ID NO:27 or an amino acid sequence having at least about 50% similarity thereto.

The term similarity as used herein includes exact identity between compared sequences at the nucleotide or amino acid level. Where there is non-identity at the nucleotide level,

- 32 -

similarity includes differences between sequences which result in different amino acids that are nevertheless related to each other at the structural, functional, biochemical and/or conformational levels. Where there is non-identity at the amino acid level, similarity includes amino acids that are nevertheless related to each other at the structural, functional, biochemical and/or conformational levels. In a particularly preferred embodiment, nucleotide and sequence comparisons are made at the level of identity rather than similarity.

Terms used to describe sequence relationships between two or more polynucleotides or polypeptides include "reference sequence", "comparison window", "sequence similarity", "sequence identity", "percentage of sequence similarity", "percentage of sequence identity", "substantially similar" and "substantial identity". A "reference sequence" is at least 12 but frequently 15 to 18 and often at least 25 or above, such as 30 monomer units, inclusive of nucleotides and amino acid residues, in length. Because two polynucleotides may each comprise (1) a sequence (i.e. only a portion of the complete polynucleotide sequence) that is similar between the two polynucleotides, and (2) a sequence that is divergent between the two polynucleotides, sequence comparisons between two (or more) polynucleotides are typically performed by comparing sequences of the two polynucleotides over a "comparison window" to identify and compare local regions of sequence similarity. A "comparison window" refers to a conceptual segment of typically 12 contiguous residues that is compared to a reference sequence. The comparison window may comprise additions or deletions (i.e. gaps) of about 20% or less as compared to the reference sequence (which does not comprise additions or deletions) for optimal alignment of the two sequences. Optimal alignment of sequences for aligning a comparison window may be conducted by computerized implementations of algorithms (GAP, BESTFIT, FASTA, and TFASTA in the Wisconsin Genetics Software Package Release 7.0, Genetics Computer Group, 575 Science Drive Madison, WI, USA) or by inspection and the best alignment (i.e. resulting in the highest percentage homology over the comparison window) generated by any of the various methods selected. Reference also may be made to the BLAST family of programs as, for example, disclosed by Altschul *et al.* (*Nucl. Acids Res.* 25: 3389-3402, 1997). A detailed discussion of sequence analysis can be found in Unit

19.3 of Ausubel *et al.* ("Current Protocols in Molecular Biology" John Wiley & Sons Inc, 1994-1998, Chapter 15, 1998).

5 The terms "sequence similarity" and "sequence identity" as used herein refers to the extent that sequences are identical or functionally or structurally similar on a nucleotide-by-nucleotide basis or an amino acid-by-amino acid basis over a window of comparison. Thus, a "percentage of sequence identity", for example, is calculated by comparing two optimally aligned sequences over the window of comparison, determining the number of positions at which the identical nucleic acid base (e.g. A, T, C, G, U) or the identical amino acid residue (e.g. Ala, Pro, Ser, Thr, Gly, Val, Leu, Ile, Phe, Tyr, Trp, Lys, Arg, His, Asp, Glu, Asn, Gln, Cys and Met) occurs in both sequences to yield the number of matched positions, dividing the number of matched positions by the total number of positions in the window of comparison (i.e., the window size), and multiplying the result by 100 to yield the percentage of sequence identity. For the purposes of the present invention, "sequence identity" will be understood to mean the "match percentage" calculated by the DNASIS computer program (Version 2.5 for windows; available from Hitachi Software engineering Co., Ltd., South San Francisco, California, USA) using standard defaults as used in the reference manual accompanying the software. Similar comments apply in relation to sequence similarity.

20 The nucleic acid molecules of the present invention may be further characterized by having, or previously having, prior to derivatization on overall lower A:T content (or higher GC content) compared to a nucleic acid molecule which encodes a R3'5'H but which does not result in detectable intact transcript in rose petal tissue or, when expressed, does not result in detectable delphinidin or delphinidin-based molecules, as measured by a chromatographic procedure such as TLC or HPLC. Furthermore, the % of A's or T's in the third position of a codon is also lower than other R4'5'H enzymes. Reference herein to a chromatographic procedure includes a related procedure. By "related" means a technically related procedure or a procedure which provides a similar result. Examples of related procedures include other forms of chromatography (eg. gas chromatography).

- 34 -

In addition, nucleotide sequences which do not express well in rose tissue may be modified such as in reducing overall % AT or at least reduce the levels of % AT in the third position of a codon. Such time expression in rose tissue is elevated.

- 5 The nucleic acid sequences contemplated herein also encompass oligonucleotides useful as genetic probes for amplification reactions or as antisense or sense molecules capable of regulating expression of the corresponding gene in a plant. Sense molecules include hairpin constructs, short double stranded DNAs and RNAs and partially double stranded DNAs and RNAs which one or more single stranded nucleotide over hangs. An antisense
- 10 molecule as used herein may also encompass a genetic construct comprising the structural genomic or cDNA gene or part thereof in reverse orientation relative to its or another promoter. It may also encompass a homologous genetic sequence. An antisense or sense molecule may also be directed to terminal or internal portions of the gene encoding a polypeptide having a F3'5'H activity or to combinations of the above such that the
- 15 expression of the gene is reduced or eliminated.

- With respect to this aspect of the invention, there is provided an oligonucleotide of 5-50 nucleotides such as 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48,
- 20 49, 50, 51, 52, 53, 54, 55 having substantial similarity to a part or region of a molecule with a nucleotide sequence set forth in SEQ ID NO:9 or SEQ ID NO:11 or SEQ ID NO:13 or SEQ ID NO:15 or SEQ ID NO:17 or SEQ ID NO:31 or SEQ ID NO:26. By substantial similarity or complementarity in this context is meant a hybridizable similarity under low, alternatively and preferably medium and alternatively and most preferably high stringency
- 25 conditions specific for oligonucleotide hybridization (Sambrook *et al.*, *Molecular Cloning: A Laboratory Manual*, 2<sup>nd</sup> edition, Cold Spring Harbor Laboratories, Cold Spring Harbor, NY, USA, 1989). Such an oligonucleotide is useful, for example, in screening for F3'5'H genetic sequences from various sources or for monitoring an introduced genetic sequence in a transgenic plant. The preferred oligonucleotide is directed to a conserved F3'5'H
- 30 genetic sequence or a sequence conserved within a plant genus, plant species and/or plant variety.

In one aspect of the present invention, the oligonucleotide corresponds to the 5' or the 3' end of the R3'5'H genetic sequences. For convenience, the 5' end is considered herein to define a region substantially between the start codon of the structural gene to a centre portion of the gene, and the 3' end is considered herein to define a region substantially between the centre portion of the gene and the terminating codon of the structural gene. It is clear, therefore, that oligonucleotides or probes may hybridize to the 5' end or the 3' end or to a region common to both the 5' and the 3' ends. The present invention extends to all such probes.

In one embodiment, the nucleic acid sequence encoding a R3'5'H or various functional derivatives thereof is used to reduce the level of an endogenous a R3'5'H (e.g. *via co-suppression or antisense-mediated suppression*) or other post-transcriptional gene silencing (PTGS) processes including RNAi or alternatively the nucleic acid sequence encoding this enzyme or various derivatives or parts thereof is used in the sense or antisense orientation to reduce the level of a R3'5'H. The use of sense strands, double or partially single stranded such as constructs with hairpin loops is particularly useful in inducing a PTGS response. In a further alternative, ribozymes, minizymes or DNAszymes could be used to inactivate target nucleic acid sequences.

Still a further embodiment encompasses post-transcriptional inhibition to reduce translation into polypeptide material. Still yet another embodiment involves specifically inducing or removing methylation.

Reference herein to the altering of a R3'5'H activity relates to an elevation or reduction in activity of up to 30% or more preferably of 30-50%, or even more preferably 50-75% or still more preferably 75% or greater above or below the normal endogenous or existing levels of activity. Such elevation or reduction may be referred to as modulation of a R3'5'H enzyme activity. Generally, modulation is at the level of transcription or translation of R3'5'H genetic sequences.



- 36 -

The nucleic acids of the present invention may be a ribonucleic acid or deoxyribonucleic acids, single or double stranded and linear or covalently closed circular molecules. Preferably, the nucleic acid molecule is cDNA. The present invention also extends to other nucleic acid molecules which hybridize under low, preferably under medium and most preferably under high stringency conditions with the nucleic acid molecules of the present invention and in particular to the sequence of nucleotides set forth in SEQ ID NO:9 or SEQ ID NO:11 or SEQ ID NO:13 or SEQ ID NO:15 or SEQ ID NO:17 or SEQ ID NO:31 or SEQ ID NO:26 or a part or region thereof. In its most preferred embodiment, the present invention extends to a nucleic acid molecule having a nucleotide sequence set forth in SEQ ID NO:9 or SEQ ID NO:11 or SEQ ID NO:13 or SEQ ID NO:15 or SEQ ID NO:17 or SEQ ID NO:31 or SEQ ID NO:26 or to a molecule having at least 40%, more preferably at least 45%, even more preferably at least 55%, still more preferably at least 65%-70%, and yet even more preferably greater than 85% similarity at the level of nucleotide or amino acid sequence to at least one or more regions of the sequence set forth in SEQ ID NO:9 or SEQ ID NO:11 or SEQ ID NO:13 or SEQ ID NO:15 or SEQ ID NO:17 or SEQ ID NO:31 or SEQ ID NO:26 and wherein the nucleic acid encodes or is complementary to a sequence which encodes an enzyme having a F3'5'H activity. It should be noted, however, that nucleotide or amino acid sequences may have similarities below the above given percentages and yet still encode a F3'5'H activity and such molecules may still be considered in the scope of the present invention where they have regions of sequence conservation. The present invention further extends to nucleic acid molecules in the form of oligonucleotide primers or probes capable of hybridizing to a portion of the nucleic acid molecules contemplated above, and in particular those set forth in SEQ ID NO:9 or SEQ ID NO:11 or SEQ ID NO:13 or SEQ ID NO:15 or SEQ ID NO:17 or SEQ ID NO:31 or SEQ ID NO:26, under low, preferably under medium and most preferably under high stringency conditions. Preferably the portion corresponds to the 5' or the 3' end of the gene. For convenience the 5' end is considered herein to define a region substantially between the start codon of the structural genetic sequence to a centre portion of the gene, and the 3' end is considered herein to define a region substantially between the centre portion of the gene and the terminating codon of the structural genetic sequence. It is clear, therefore, that

oligonucleotides or probes may hybridize to the 5' end or the 3' end or to a region common to both the 5' and the 3' ends. The present invention extends to all such probes.

The term gene is used in its broadest sense and includes cDNA corresponding to the exons of a gene. Accordingly, reference herein to a gene is to be taken to include:-

(i) a classical genomic gene consisting of transcriptional and/or translational regulatory sequences and/or a coding region and/or non-translated sequences (i.e. introns, 5'- and 3'- untranslated sequences); or

(ii) mRNA or cDNA corresponding to the coding regions (i.e. exons) and 5'- and 3'- untranslated sequences of the gene.

The term gene is also used to describe synthetic or fusion molecules encoding all or part of an expression product. In particular embodiments, the term nucleic acid molecule and gene may be used interchangeably.

The nucleic acid or its complementary form may encode the full-length enzyme or a part or derivative thereof. By "derivative" is meant any single or multiple amino acid substitutions, deletions, and/or additions relative to the naturally occurring enzyme and which retains a 3'5'TH activity. In this regard, the nucleic acid includes the naturally occurring nucleotide sequence encoding a 3'5'TH or may contain single or multiple nucleotide substitutions, deletions and/or additions to said naturally occurring sequence. The nucleic acid of the present invention or its complementary form may also encode a "part" of the 3'5'TH, whether active or inactive, and such a nucleic acid molecule may be useful as an oligonucleotide probe, primer for polymerase chain reactions or in various mutagenic techniques, or for the generation of antisense molecules.

Reference herein to a "part" of a nucleic acid molecule, nucleotide sequence or amino acid sequence, preferably relates to a molecule which contains at least about 10 contiguous nucleotides or five contiguous amino acids, as appropriate.

Amino acid insertional derivatives of the F3'5'H of the present invention include amino and/or carboxyl terminal fusions as well as intra-sequence insertions of single or multiple amino acids. Insertional amino acid sequence variants are those in which one or more amino acid residues are introduced into a predetermined site in the protein although random insertion is also possible with suitable screening of the resulting product. Deletional variants are characterized by the removal of one or more amino acids from the sequence. Substitutional amino acid variants are those in which at least one residue in the sequence has been removed and a different residue inserted in its place. Typical substitutions are those made in accordance with Table 3.

**TABLE 3** Suitable residues for amino acid substitutions

Original residue	Exemplary substitutions
Ala	Ser
Arg	Lys
Asn	Gln; His
Asp	Glu
Cys	Ser
Gln	Asn; Glu
Glu	Asp
Gly	Pro
His	Asn; Gln
Ile	Leu; Val
Leu	Ile; Val
Lys	Arg; Gln; Glu
Met	Leu; Ile; Val
Phe	Met; Leu; Tyr
Ser	Thr
Thr	Ser

25 The terms "analog" and "derivatives" also extend to any functional chemical equivalent of a R3'5'H and also to any amino acid derivative described above. For convenience, reference to R3'5'H herein includes reference to any functional mutant, derivative, part, fragment, homolog or analog thereof.

20 Other examples of recombinant or synthetic mutants and derivatives of the R3'5'H enzyme of the present invention include single or multiple substitutions, deletions and/or additions of any molecule associated with the enzyme such as carbohydrates, lipids and/or proteins or polypeptides.

15 example, in Sambrook *et al.* (1989, *supra*). Techniques for making substitution mutations at predetermined sites in DNA having known or partially known sequence are well known and include, for example, M13 mutagenesis. The manipulation of DNA sequence to produce variant proteins which manifest as substitutional, insertional or deletional variants are conveniently described, for The amino acid variants referred to above may readily be made using peptide synthetic techniques well known in the art, such as solid phase peptide synthesis (McIntire, *J. Am. Chem. Soc.* 85: 2149, 1964) and the like, or by recombinant DNA manipulations.

10 Where the R3'5'H is derivatized by amino acid substitution, the amino acids are generally replaced by other amino acids having like properties, such as hydrophobicity, hydrophilicity, electropositivity, bulky side chains and the like. Amino acid substitutions are typically of single residues. Amino acid insertions will usually be in the order of about 1-10 amino acid residues and deletions will range from about 1-20 residues. Preferably, deletions or insertions are made in adjacent pairs, i.e. a deletion of two residues or insertion of two residues.

Original residue	Exemplary substitutions
Tyr	Tyr
Tyr	Trp; Phe
Val	Ile; Leu; Met

- 40 -

The present invention is exemplified using nucleic acid sequences derived from pansy, salvia, sollya or lavender or kennedia since this represents the most convenient and preferred source of material to date. However, one skilled in the art will immediately appreciate that similar sequences can be isolated from any number of sources such as other plants or certain microorganisms. All such nucleic acid sequences encoding directly or indirectly a F3'5'H are encompassed by the present invention regardless of their source. Examples of other suitable sources of genes encoding F3'5'H include, but are not limited to *Vitis spp.*, *Babiana stricta*, *Pinus spp.*, *Picea spp.*, *Larix spp.*, *Phaseolus spp.*, *Vaccinium spp.*, *Cyclamen spp.*, *Iris spp.*, *Pelargonium spp.*, *Liparidae*, *Geranium spp.*, *Plum spp.*, *Lathyrus spp.*, *Clitoria spp.*, *Catharanthus spp.*, *Malva spp.*, *Mucuna spp.*, *Vicia spp.*, *Saintpaulia spp.*, *Lagerstroemia spp.*, *bouchina spp.*, *Plumbago spp.*, *Hypocalyptus spp.*, *Rhododendron spp.*, *Linum spp.*, *Macroptilium spp.*, *Hibiscus spp.*, *Hydrangea spp.*, *Cymbidium spp.*, *Millettia spp.*, *Hedysarum spp.*, *Lespedeza spp.*, *Asparagus spp.*, *Antigonon spp.*, *Freesia spp.*, *Brunella spp.*, *Clarkia spp.*, etc.

In accordance with the present invention, a nucleic acid sequence encoding a F3'5'H may be introduced into and expressed in a transgenic plant in either orientation thereby providing a means either to convert suitable substrates, if synthesized in the plant cell, ultimately into DHM, or alternatively to inhibit such conversion of metabolites by reducing or eliminating endogenous or existing F3'5'H activity. The production of these 3', 5'-hydroxylated substrates will subsequently be converted to delphinidin-based pigments that will modify petal color and may contribute to the production of a bluer color. Expression of the nucleic acid sequence in the plant may be constitutive, inducible or developmental and may also be tissue-specific. The word "expression" is used in its broadest sense to include production of RNA or of both RNA and protein. It also extends to partial expression of a nucleic acid molecule.

According to this aspect of the present invention, there is provided a method for producing a transgenic flowering plant capable of synthesizing a F3'5'H, said method comprising stably transforming a cell of a suitable plant with a nucleic acid sequence which comprises a sequence of nucleotides encoding said F3'5'H under conditions permitting the eventual

expression of said nucleic acid sequence, regenerating a transgenic plant from the cell and growing said transgenic plant for a time and under conditions sufficient to permit the transgenic plant to produce non-expression of the nucleic acid sequence. The transgenic plant may thereby produce non-indigenous F3'5'H at elevated levels relative to the amount expressed in a comparable non-transgenic plant.

Another aspect of the present invention contemplates a method for producing a transgenic plant with reduced indigenous or existing flavonoid 3', 5'-hydroxylase activity, said method comprising stably transforming a cell of a suitable plant with a nucleic acid molecule which comprises a sequence of nucleotides encoding or complementary to a sequence encoding a F3'5'H activity, regenerating a transgenic plant from the cell and where necessary growing said transgenic plant under conditions sufficient to permit the expression of the nucleic acid.

Yet another aspect of the present invention contemplates a method for producing a genetically modified plant with reduced indigenous or existing F3'5'H activity, said method comprising altering the F3'5'H gene through modification of the indigenous sequences via homologous recombination from an appropriately altered F3'5'H gene or derivative or part thereof introduced into the plant cell, and regenerating the genetically modified plant from the cell.

As used herein an "indigenous" enzyme is one, which is native to or naturally expressed in a particular cell. A "non-indigenous" enzyme is an enzyme not native to the cell but expressed through the introduction of genetic material into a plant cell, for example, through a transgene. An "endogenous" enzyme is an enzyme produced by a cell but which may or may not be indigenous to that cell.

In a preferred embodiment, the present invention contemplates a method for producing a transgenic flowering plant exhibiting altered floral or inflorescence properties, said method comprising stably transforming a cell of a suitable plant with a nucleic acid sequence of the present invention, regenerating a transgenic plant from the cell and growing said transgenic

- 42 -

plant for a time and under conditions sufficient to permit the expression of the nucleic acid sequence.

Alternatively, said method may comprise stably transforming a cell of a suitable plant with  
5 a nucleic acid sequence of the present invention or its complementary sequence, regenerating a transgenic plant from the cell and growing said transgenic plant for a time and under conditions sufficient to alter the level of activity of the indigenous or existing F3'5'H. Preferably the altered level would be less than the indigenous or existing level of F3'5'H activity in a comparable non-transgenic plant. Without wishing to limit the present  
10 invention, one theory of mode of action is that reduction of the indigenous F3'5'H activity requires the expression of the introduced nucleic acid sequence or its complementary sequence. However, expression of the introduced genetic sequence or its complement may not be required to achieve the desired effect: namely, a flowering plant exhibiting altered floral or inflorescence properties.

15 In a related embodiment, the present invention contemplates a method for producing a flowering plant exhibiting altered floral or inflorescence properties, said method comprising alteration of the flavonoid 3', 5'-hydroxylase gene through modification of the indigenous sequences via homologous recombination from an appropriately altered F3'5'H  
20 gene or derivative or part thereof introduced into the plant cell, and regenerating the genetically modified plant from the cell.

Preferably, the altered floral or inflorescence includes the production of different shades of blue or purple or red flowers or other colors, depending on the genotype and physiological  
25 conditions of the recipient plant.

Accordingly, the present invention extends to a method for producing a transgenic plant capable of expressing a recombinant gene encoding a F3'5'H or part thereof or which carries a nucleic acid sequence which is substantially complementary to all or a part of a  
30 mRNA molecule encoding the F3'5'H, said method comprising stably transforming a cell of a suitable plant with the isolated nucleic acid molecule comprising a sequence of

nucleotides encoding, or complementary to a sequence encoding, a R3'5'H, where necessary under conditions permitting the eventual expression of said isolated nucleic acid molecule, and regenerating a transgenic plant from the cell. By suitable plant is meant a plant capable of producing DHK and possessing the appropriate physiological properties required for the development of the color desired.

One skilled in the art will immediately recognise the variations applicable to the methods of the present invention, such as increasing or decreasing the expression of the enzyme naturally present in a target plant leading to differing shades of colors such as different shades of blue, purple or red.

The present invention, therefore, extends to all transgenic plants or parts or cells therefrom of transgenic plants or progeny of the transgenic plants containing all or part of the nucleic acid sequences of the present invention, or antisense forms thereof and/or any homologs or related forms thereof and, in particular, those transgenic plants which exhibit altered floral or inflorescence properties. The transgenic plants may contain an introduced nucleic acid molecule comprising a nucleotide sequence encoding or complementary to a sequence encoding a R3'5'H. Generally, the nucleic acid would be stably introduced into the plant genome, although the present invention also extends to the introduction of a R3'5'H nucleotide sequence within an autonomously-replicating nucleic acid sequence such as a DNA or RNA virus capable of replicating within the plant cell. The invention also extends to seeds from such transgenic plants. Such seeds, especially if colored, are useful as proprietary tags for plants. Any and all methods for introducing genetic material into plant cells including but not limited to *Agrobacterium*-mediated transformation, biolistic particle bombardment etc. are encompassed by the present invention.

Another aspect of the present invention contemplates the use of the extracts from transgenic plants or plant parts or cells therefrom of transgenic plants or progeny of the transgenic plants containing all or part of the nucleic acid sequences of the present invention and, in particular, the extracts from those transgenic plants which used as a flavouring or food additive or health product or beverage or juice or coloring.



- 44 -

Plant parts contemplated by the present invention includes, but is not limited to flowers, fruits, vegetables, nuts, roots, stems, leaves or seeds.

- 5 The extracts of the present invention may be derived from the plants or plant part or cells therefrom in a number of different ways including but not limited to chemical extraction or heat extraction or filtration or squeezing or pulverization.

- 10 The plant, plant part or cells therefrom or extract can be utilized in any number of different ways such as for the production of a flavouring (e.g. a food essence), a food additive (e.g. a stabilizer, a colorant) a health product (e.g. an antioxidant, a tablet) a beverage (e.g. wine, spirit, tea) or a juice (e.g. fruit juice) or coloring (e.g. food coloring, fabric coloring, dye, paint, tint).

- 15 A further aspect of the present invention is directed to recombinant forms of F3'5'H. The recombinant forms of the enzyme will provide a source of material for research, for example, more active enzymes and may be useful in developing *in vitro* systems for production of colored compounds.

- 20 Still a further aspect of the present invention contemplates the use of the genetic sequences described herein in the manufacture of a genetic construct capable of expressing a F3'5'H or down-regulating an indigenous F3'5'H enzyme in a plant.

- 25 The term genetic construct has been used interchangeably throughout the specification and claims with the terms "fusion molecule", "recombinant molecule", "recombinant nucleotide sequence". A genetic construct may include a single nucleic acid molecule comprising a nucleotide sequence encoding a single protein or may contain multiple open reading frames encoding 2 or more proteins. It may also contain a promoter operably linked to 1 or more of the open reading frames.

30

- 45 -

Another aspect of the present invention is directed to a prokaryotic or eukaryotic organism carrying a genetic sequence encoding a R3'5'H extrachromasomally in plasmid form.

The present invention further extends to a recombinant polypeptide comprising a sequence of amino acids substantially as set forth in SEQ ID NO:10 or SEQ ID NO:12 or SEQ ID NO:14 or SEQ ID NO:16 or SEQ ID NO:18 or SEQ ID NO:32 or SEQ ID NO:27 or an amino acid sequence having at least about 50% similarity to SEQ ID NO:10 or SEQ ID NO:12 or SEQ ID NO:14 or SEQ ID NO:16 or SEQ ID NO:18 or SEQ ID NO:32 or SEQ ID NO:27 or a derivative of said polypeptide.

A "recombinant polypeptide" means a polypeptide encoded by a nucleotide sequence introduced into a cell directly or indirectly by human intervention or into a parent or other relative or precursor of the cell. A recombinant polypeptide may also be made using cell-free, *in vitro* transcription systems. The term "recombinant polypeptide" includes an isolated polypeptide or when present in a cell or cell preparation. It may also be in a plant or parts of a plant regenerated from a cell which produces said polypeptide.

A "polypeptide" includes a peptide or protein and is encompassed by the term "enzyme". The recombinant polypeptide may also be a fusion molecule comprising two or more heterologous amino acid sequences.

The present invention is further described by the following non-limiting examples.

- 46 -

**EXAMPLE 1*****General methods***

In general, the methods followed were as described in Sambrook *et al.* (1989, *supra*) or  
 5 Sambrook and Russell, *Molecular Cloning: A Laboratory Manual* 3<sup>rd</sup> edition, Cold Spring  
 Harbor Laboratories, Cold Spring Harbor, NY, USA, 2001 or *Plant Molecular Biology*  
*Manual* (2<sup>nd</sup> edition), Gelvin and Schilperoot (eds), Kluwer Academic Publisher, The  
 Netherlands, 1994 or *Plant Molecular Biology Labfax*, Croy (ed), Bios scientific  
 Publishers, Oxford, UK, 1993.

10

The cloning vectors pBluescript and PCR script were obtained from Stratagene, USA.  
 pCR7 2.1 was obtained from Invitrogen, USA.

***E. coli transformation***

15 The *Escherichia coli* strains used were:

**DH5 $\alpha$** 

supB44,  $\Delta$  (lacZYA-ArgF)U169, ( $\phi$ 80lacZAM15), hsdR17( $r_K^-$ ,  $m_K^+$ ),  
 recA1, endA1, gyrA96, thi-1, relA1, deoR. (Hanahan, *J. Mol. Biol.* 166: 557, 1983)

20

**XLI-Blue**

supB44, hsdR17( $r_K^-$ ,  $m_K^+$ ), recA1, endA1, gyrA96, thi-1, relA1.  
 lac<sup>+</sup>, [F<sup>+</sup>proAB, lacI<sup>q</sup>, lacZAM15, Tn10(tet<sup>R</sup>)] (Bullock *et al.*, *Biotechniques* 5: 376, 1987).

25 BL21-CodonPlus-RIL strain

*ompT hsdS(Rb- mB-) dcm+ Tet<sup>r</sup> gal endA Hte [argU ileY leuW Cam<sup>r</sup>]*

M15 *E. coli* is derived from *E. coli* K12 and has the phenotype NaI<sup>s</sup>, Str<sup>s</sup>, Rif<sup>s</sup>, Thi<sup>-</sup>, Ara<sup>+</sup>,  
 Gal<sup>+</sup>, Mil<sup>+</sup>, F<sup>-</sup>, RecA<sup>+</sup>, Uvr<sup>+</sup>, Lon<sup>+</sup>.

30 Transformation of the *E. coli* strains was performed according to the method of Inoue *et al.*, (*Gene* 96: 23-28, 1990).

30 Repair of overhanging ends after restriction endonuclease digestion  
Overhanging 5' ends were repaired using DNA polymerase (Klenow fragment) according to standard protocols (Sambrook *et al.*, 1989 *supra*). Overhanging 3' ends were repaired using T4 DNA polymerase according to standard protocols (Sambrook *et al.*, 1989 *supra*).

25 Isolation and purification of DNA fragments  
Fragments were generally isolated on a 1% (w/v) agarose gel and purified using the QIAEX II Gel Extraction kit (Qiagen) or Brecsaclean Kit (Bresatec, Australia) following procedures recommended by the manufacturer.

20 DNA ligations  
DNA ligations were carried out using the Amersham Ligation Kit or Promega Ligation Kit according to procedures recommended by the manufacturer.

5 Plasmid DNA was introduced into the *Agrobacterium tumefaciens* strain AGLO by adding 5 µg of plasmid DNA to 100 µL of competent AGLO cells prepared by inoculating a 50 mL LB culture (Sambrook *et al.*, 1989, *supra*) and incubation for 16 hours with shaking at 28°C. The cells were then pelleted and resuspended in 0.5 mL of 85% (v/v) 100 mM CaCl<sub>2</sub>/15% (v/v) glycerol. The DNA-*Agrobacterium* mixture was frozen by incubation in liquid N<sub>2</sub> for 2 minutes and then allowed to thaw by incubation at 37°C for 5 minutes. The DNA/bacterial mix was then placed on ice for a further 10 minutes. The cells were then mixed with 1 mL of LB (Sambrook *et al.*, 1989 *supra*) media and incubated with shaking for 16 hours at 28°C. Cells of *A. tumefaciens* carrying the plasmid were selected on LB agar plates containing appropriate antibiotics such as 50 µg/mL tetracycline or 100 µg/mL gentamycin. The confirmation of the plasmid in *A. tumefaciens* was done by restriction endonuclease mapping of DNA isolated from the antibiotic-resistant transformants.

The disarmed *Agrobacterium tumefaciens* strain used was AGLO (Lazo *et al.* *Bio/technology* 9: 963-967, 1991).

*Agrobacterium tumefaciens* strains and transformations

- 48 -

**Removal of phosphoryl groups from nucleic acids**

Shrimp alkaline phosphatase (SAP) (USB) was typically used to remove phosphoryl groups from cloning vectors to prevent re-circularization according to the manufacturer's recommendations.

**Polymerase Chain Reaction (PCR)**

Unless otherwise specified, PCR conditions using plasmid DNA as template included using 2 ng of plasmid DNA, 100 ng of each primer, 2  $\mu$ L 10 mM dNTP mix, 5  $\mu$ L 10 x Taq DNA polymerase buffer, 0.5  $\mu$ L Taq DNA Polymerase in a total volume of 50  $\mu$ L. Cycling conditions comprised an initial denaturation step of 5 minutes at 94°C, followed by 35 cycles of 94°C for 20 sec, 50°C for 30 sec and 72°C for 1 minute with a final treatment at 72°C for 10 minutes before storage at 4°C.

PCRs were performed in a Perkin Elmer GeneAmp PCR System 9600.

**<sup>32</sup>P-Labelling of DNA Probes**

DNA fragments (50 to 100 ng) were radioactively labelled with 50  $\mu$ Ci of [ $\alpha$ -<sup>32</sup>P]-dCTP using a Gigaprime kit (Geneworks). Unincorporated [ $\alpha$ -<sup>32</sup>P]-dCTP was removed by chromatography on Sephadex G-50 (Fine) columns or Microbiospin P-30 Tris chromatography columns (BioRad).

**Plasmid Isolation**

Single colonies were analyzed for inserts by inoculating LB broth (Sambrook *et al.*, 1989, *supra*) with appropriate antibiotic selection (e.g. 100  $\mu$ g/mL ampicillin or 10 to 50  $\mu$ g/mL tetracycline etc.) and incubating the liquid culture at 37°C (for *E. coli*) or 29°C (for *A. tumefaciens*) for ~16 hours with shaking. Plasmid DNA was purified using the alkali-lysis procedure (Sambrook *et al.*, 1989, *supra*) or using The WizardPlus SV minipreps DNA purification system (Promega) or Qiagen Plasmid Mini Kit (Qiagen). Once the presence of an insert had been determined, larger amounts of plasmid DNA were prepared from 50 mL overnight cultures using the alkali-lysis procedure (Sambrook *et al.*, 1989, *supra*) or

- 49 -

QIAGEN Plasmid Midi kit (QIAGEN) and following conditions recommended by the manufacturer.

### DNA Sequence Analysis

5 DNA sequencing was performed using the PRISM (trademark) Ready Reaction Dye Primer Cycle Sequencing Kits from Applied Biosystems. The protocols supplied by the manufacturer were followed. The cycle sequencing reactions were performed using a Perkin Elmer PCR machine (GeneAmp PCR System 9600). Sequencing runs were generally performed by the Australian Genome Research Facility at The Walter and Eliza Hall Institute of Medical Research (Melbourne, Australia) or in-house on an automated 373A DNA sequencer (Applied Biosystems).

Sequences were analysed using a MacVector™ application (version 6.5.3) (Oxford Molecular Ltd, Oxford, England).

15 Homology searches against Genbank, SWISS-PROT and EMBL databases were performed using the RASTA and TFASTA programs (Pearson and Lipman, *Proc. Natl. Acad. Sci. USA* 85(8): 2444-2448, 1988) or BLAST programs (Altschul *et al.*, *J. Mol. Biol.* 215(3): 403-410, 1990). Percentage sequence similarities were obtained using LALIGN program (Burg and Miller, *Adv. Appl. Math.* 12: 373-381, 1991) or ClustalW program (Thompson *et al.*, *Nucleic Acids Research* 22: 4673-4680, 1994) within the MacVector™ application (Oxford Molecular Ltd, England) using default settings.

25 Multiple sequence alignments were produced using ClustalW (Thompson *et al.*, 1994, *supra*) using default settings.

- 50 -

**EXAMPLE 2*****Plant transformations******Petunia hybrida* transformations (Sw63 x Skr4)**

- 5 As described in Holton *et al.* (1993a, *supra*) by any other method well known in the art.

***Rosa hybrida* transformations**

- As described in U.S. Patent Application No. 542,841 (PCT/US91/04412) or Robinson and Firoozabady (*Scientia Horticulturae*, 55: 83-99, 1993), Rout *et al.* (*Scientia Horticulturae*,  
10 81: 201-238, 1999) or Marchant *et al.* (*Molecular Breeding* 4: 187-194, 1998) or by any other method well known in the art.

Cuttings of *Rosa hybrida* were generally obtained from Van Wyk and Son Flower Supply, Victoria.

15

***Dianthus caryophyllus* transformations**

- International Patent Application No. PCT/US92/02612 (carnation transformation). As described in International Patent Application No. PCT/AU96/00296 (Violet carnation), Lu  
20 *et al.* (*Bio/Technology* 9: 864-868, 1991), Robinson and Firoozabady (1993, *supra*) or by any other method well known in the art.

Cuttings of *Dianthus caryophyllus* cv. Kortina Chandel or Monte Lisa were obtained from Van Wyk and Son Flower Supply, Victoria.

25

- 51 -

## EXAMPLE 3

*Transgenic Analysis**Color coding*

5 The Royal Horticultural Society's Color Chart (Kew, UK) was used to provide a description of observed color. They provide an alternative means by which to describe the color phenotypes observed. The designated numbers, however, should be taken only as a guide to the perceived colors and should not be regarded as limiting the possible colors which may be obtained.

*Chromatographic analysis*

Thin Layer Chromatography (TLC) and High Performance Liquid Chromatography (HPLC) analysis was performed generally as described in Bruguiera *et al.* (*Plant J.* 5, 81-92, 1994).

*Extraction of anthocyanidins*

Prior to HPLC analysis, the anthocyanin and flavonol molecules present in petal and stamen extracts were acid hydrolysed to remove glycosyl moieties from the anthocyanidin or flavonol core. Anthocyanidin and flavonol standards were used to help identify the compounds present in the floral extracts.

Anthocyanidins in the reaction mixture were analysed by HPLC via gradient elution using gradient conditions of 50%B to 60%B over 10 minutes, then 60% B for 10 minutes and finally 60% B to 100% B over 5 minutes where solvent A consisted of TFA: H<sub>2</sub>O (5:995) and solvent B consisted of acetonitrile: TFA: H<sub>2</sub>O (500:5:495). An Asahi Pac ODP-50 cartridge column (250 mm x 4.6 mm ID) was used for the reversed phase chromatographic separations. The flow rate was 1 mL/min and the temperature was 40°C. The detection of the anthocyanidin compounds was carried out using a Shimadzu SPD-M6A three dimensional detector at 400-650 nm.

30

25

20

15

10

5



- 52 -

The anthocyanidin peaks were identified by reference to known standards, viz delphinidin or delphinidin-based molecules, petunidin, malvidin, cyanidin and peonidin

**Stages of flower development**

5

***Petunia***

*Petunia hybrida* cv. Skr4 x Sw63 flowers were harvested at developmental stages defined as follows:

- 10    Stage 1:        Unpigmented, closed bud.  
      Stage 2:        Pigmented, closed bud.  
      Stage 3:        Pigmented bud with emerging corolla  
      Stage 4:        Pigmented, opened flower with anther intact (pre-dehiscence)  
      Stage 5:        Fully opened flower with all anthers dehisced.

15

For TLC or HPLC analysis, petals were collected from stage 4 flowers at the stage of maximum pigment accumulation.

- For Northern blot analysis, petals were collected from stages 2 to 3 flowers at the stage of  
20    maximal expression of flavonoid pathway genes.

***Carnation***

*Dianthus caryophyllus* flowers were harvested at developmental stages defined as follows:

- 25    Stage 1:        Closed bud, petals not visible.  
      Stage 2:        Flower buds opening: tips of petals visible.  
      Stage 3:        Tips of nearly all petals exposed. "Paint-brush stage".  
      Stage 4:        Outer petals at 45° angle to stem.  
      Stage 5:        Flower fully open.

30

For TLC or HPLC analysis, petals were collected from stage 4 flowers at the stage of maximum pigment accumulation.

For Northern blot analysis, petals were collected from stage 3 flowers at the stage of maximal expression of flavonoid pathway genes.

# Rose

Stages of *Rosa hybrida* flower development were defined as follows:

- 10 Stages 1: Unpigmented, tightly closed bud
- Stage 2: Pigmmented, tightly closed bud
- Stage 3: Pigmmented, closed bud; sepals just beginning to open.
- Stage 4: Flower bud beginning to open; petals heavily pigmented; sepals have separated.
- 15 Stage 5: Sepals completely unfolded; some curling. Petals are heavily pigmented and unfolding.

For TLC or HPLC analysis, petals were collected from stage 4 flowers at the stage of maximum pigment accumulation.

For Northern blot analysis, petals were collected from stage 3 to 4 flowers at the stage of maximal expression of flavonoid pathway genes (Tanaka *et al*, *Plant Cell Physiol.*, 36(6): 1023-1031, 1995).

## Anthocyanin/Flavonol measurements by spectrophotometric measurements

Approximately 200mg of fresh petal tissue was added to 2 mL of methanol/1% (v/v) HCl and incubated for ~16 hours at 4°C. A 1 in 20 dilution (e.g. 50 µL made to 1000 µL) was then made and the absorbance at 350 nm and 530 nm was recorded.

The approximate flavonols and anthocyanin amounts (mmoles/gram) were then calculated according to the following formulae:

- 54 -

*Anthocyanin content*

$$\frac{(A_{530} / 34,000) \times \text{volume of extraction buffer (mL)} \times \text{dilution factor} \times 10^6}{\text{mass of petal tissue (grams)}}$$

5

*Flavonol content*

$$\frac{(A_{350} / 14,300) \times \text{volume of extraction buffer (mL)} \times \text{dilution factor} \times 10^6}{\text{mass of petal tissue (grams)}}$$

10 *Northern/RNA blot analysis*

Transcription of a transferred gene was monitored by isolating RNA and estimating the quantity and size of the expected transcript. Northern blot analysis was used to monitor the steady-state level of particular transcripts in petals. A transcript was determined to be intact or full-length based on the estimated size expected from the gene used. In general

15 when cDNAs were used as coding sequences the size of the transcript expected would be the size of the cDNA plus any 5' untranslated component of the fused promoter fragment plus any 3' untranslated sequence from the fused terminator fragment. In some cases where a cDNA region contained a putative polyadenylation site and the terminator region contained a putative polyadenylation site, 2 transcripts would be detected. One would be of

20 a size consistent with polyadenylation occurring just downstream from the polyadenylation site within the cDNA sequence. The second transcript would be larger and consistent with the transcript being polyadenylated after the polyadenylation site within the terminator fragment.

25 Total RNA was isolated from petals or leaves using a Plant RNAeasy kit (QIAGEN) following procedures recommended by the manufacturer. For rose samples 1% (w/v) PVP was added to the extraction buffer.

RNA samples (5 µg) were electrophoresed through 2.2 M formaldehyde/1.2% w/v agarose gels using running buffer containing 40 mM morpholinopropanesulphonic acid (pH 7.0), 5 mM sodium acetate, 0.1 mM EDTA (pH 8.0). The RNA was stained with ethidium bromide and visualised under UV-light. The ribosomal RNA was generally used as a guide in confirming that the RNA had not been degraded by intra- or extra- cellular ribonucleases. The RNA was transferred to Hybond-N membrane filters (Amersham) and treated as described by the manufacturer.

Control samples were included on RNA gels as a measure of the integrity of the radiolabelled probe and as guides to expected transcript sizes. Controls for *petH1* and *petH2* genes included RNA isolated from petunia OGB petals (stages 3 to 4) or from flowers of transgenic carnations shown previously to accumulate *petH1* transcripts. Controls for other *F3'5'H* genes generally included RNA isolated from petals of the same species from which the *F3'5'H* sequence had been isolated.

RNA blots were probed with <sup>32</sup>P-labelled fragments. Prehybridization (1 hour at 42°C) and hybridization (16 hours at 42°C) of the membrane filters were carried out in 50% v/v formamide, 1 M NaCl, 1% w/v SDS, 10% w/v dextran sulphate. The membrane filters were generally washed in 2 x SSC, 1% w/v SDS at 65°C for between 1 to 2 hours and then 0.2 x SSC, 1% w/v SDS at 65°C for between 0.5 to 1 hour. Membrane filters were generally exposed to Kodak XAR film with an intensifying screen at -70°C for 16 to 72 hours.

#### EXAMPLE 4

##### *Introduction of chimeric petunia F3'5'H genes into rose*

As described in the introduction, the pattern of hydroxylation of the B-ring of the anthocyanidin molecule plays a key role in determining petal color. The production of the dihydroflavonol DHM, leads to the production of the purple/blue delphinidin-based pigments in plants such as petunia. The absence of the F3'5'H activity has been correlated

- 56 -

with the absence of blue flowers in many plant species such as *Rosa*, *Gerbera*, *Antirrhinum*, *Dianthus* and *Dendranthema*.

Based on success in producing delphinidin-based pigments in a mutant petunia line (Holton *et al.*, 1993a, *supra* and International Patent Application No. PCT/AU92/00334), in tobacco flowers (International Patent Application No. PCT/AU92/00334) and in carnation flowers (International Patent Application No. PCT/AU96/00296), similar chimeric petunia *F3'5'H* genes were also introduced into roses in order to produce novel delphinidin-based pigments and modify flower color.

#### Preparation of chimeric petunia *F3'5'H* gene constructs

A summary of promoter, terminator and coding fragments used in the preparation of constructs and the respective abbreviations is listed in Table 4.

TABLE 4 Abbreviations used in construct preparations

ABBREVIATION	DESCRIPTION
<i>AmCHS 5'</i>	1.2 kb promoter fragment from the <i>Antirrhinum majus</i> chalcone synthase ( <i>CHS</i> ) gene (Sommer and Saedler, <i>Mol Gen. Gen.</i> , 202: 429-434, 1986)
<i>CaMV 35S</i>	~0.2 kb incorporating <i>Bgl</i> III fragment containing the promoter region from the Cauliflower Mosaic Virus 35S ( <i>CaMV 35S</i> ) gene - (Franck <i>et al.</i> , <i>Cell</i> 21: 285-294, 1980, Guilley <i>et al.</i> , <i>Cell</i> , 30: 763-773, 1982)
<i>35S 5'</i>	promoter fragment from <i>CaMV 35S</i> gene (Franck <i>et al.</i> , 1980, <i>supra</i> ) with an ~60bp 5' untranslated leader sequence from the petunia chlorophyll a/b binding protein gene ( <i>Cab 22</i> gene) (Harpster <i>et al.</i> , <i>MGG</i> , 212: 182-190, 1988)
<i>chrysCHS 5'</i>	promoter region from a <i>CHS</i> gene from chrysanthemum (SEQ ID NO: 30)

ABBREVIATION	DESCRIPTION
<i>aCAMV 35S</i>	enhanced <i>aCAMV 35S</i> promoter as described in Mitsuhashi <i>et al.</i> , <i>Plant Cell Physiol.</i> 37: 49-59, 1996
<i>GUS</i>	$\beta$ -glucuronidase ( <i>GUS</i> ) coding sequence (Jefferson, <i>et al.</i> , <i>EMBO J.</i> 6: 3901-3907, 1987)
<i>Mac</i>	Hybrid promoter consisting of the promoter from the mannopine synthase ( <i>mas</i> ) gene and a <i>aCAMV 35S</i> enhancer region (Conrad <i>et al.</i> , <i>Plant Mol. Biol.</i> 15: 373-381, 1990)
<i>mas/35S</i>	Hybrid promoter consisting of a promoter region from <i>aCAMV 35S</i> gene with enhancer elements from a promoter fragment of mannopine synthase ( <i>mas</i> ) gene of <i>Agrobacterium tumefaciens</i> (Janssen and Gardner, <i>Plant Molecular Biology</i> , 14: 61-72, 1989)
<i>mas 5'</i>	Promoter region from the <i>mas</i> of <i>A. tumefaciens</i>
<i>mas 3'</i>	Terminator region from the <i>mas</i> gene of <i>A. tumefaciens</i>
<i>nos 5'</i>	Promoter region from the nopaline synthase ( <i>nos</i> ) gene of <i>A. tumefaciens</i> (DePickett <i>et al.</i> , <i>J Mol and Appl Genetics</i> 1: 561-573, 1982)
<i>nos 3'</i>	Terminator region from the <i>nos</i> gene of <i>A. tumefaciens</i> (DePickett <i>et al.</i> , 1982, <i>supra</i> )
<i>npII</i>	Kanamycin-resistance gene (encodes neomycin phosphotransferase which deactivates aminoglycoside antibiotics such as kanamycin, neomycin and G418)
<i>ocs 3'</i>	~1.6kb terminator fragment from octopine synthase gene of <i>A. tumefaciens</i> (described in Janssen and Gardner, 1989, <i>supra</i> )
<i>pelD 5'</i>	~3.2kb promoter region from a phospholipid transfer protein gene (D8) of <i>Petunia hybrida</i> (Holton, Isolation and characterization of petal specific genes from <i>Petunia hybrida</i> . PhD thesis, University of Melbourne, Australia, 1992) (SEQ ID NO: 24)
<i>pelD 3'</i>	~0.7kb terminator region from a phospholipid transfer protein gene (D8) of <i>Petunia hybrida</i> cv. OGB (Holton, 1992, <i>supra</i> )

ABBREVIATION	DESCRIPTION
<i>long petFLS 5'</i>	~4.0kb fragment containing the promoter region from a flavonol synthase ( <i>FLS</i> ) gene of <i>P. hybrida</i>
<i>short petFLS 5'</i>	~2.2kb fragment containing the promoter region from <i>FLS</i> gene of <i>P. hybrida</i>
<i>petFLS 3'</i>	~0.95kb fragment containing the terminator region from <i>FLS</i> gene of <i>P. hybrida</i>
<i>petHf1</i>	Petunia <i>F3'5'H Hf1</i> cDNA clone (Holton <i>et al.</i> , 1993a, <i>supra</i> ) (SEQ ID NO: 1)
<i>petHf2</i>	Petunia <i>F3'5'H Hf2</i> cDNA clone (Holton <i>et al.</i> , 1993a, <i>supra</i> ) (SEQ ID NO: 3)
<i>petRT 5'</i>	Promoter region of an anthocyanidin-3- glucoside rhamnosyltransferase ( <i>3RT</i> ) gene from <i>P. hybrida</i> (Brugliera, Characterization of floral specific genes isolated from <i>Petunia hybrida</i> . RMIT, Australia. PhD thesis, 1994)
<i>petRT 3'</i>	Terminator region of a <i>3RT</i> gene from <i>P. hybrida</i> (Brugliera, 1994, <i>supra</i> )
<i>RoseCHS 5'</i>	~2.8kb fragment containing the promoter region from a <i>CHS</i> gene of <i>Rosa hybrida</i> (SEQ ID: 5)
<i>SuRB</i>	Chlorsulfuron-resistance gene (encodes Acetolactate Synthase) with its own terminator from <i>Nicotiana tabacum</i> (Lee <i>et al.</i> , <i>EMBO J.</i> 7: 1241-1248, 1988)

In order to produce delphinidin or delphinidin-based molecules in rose petals, a number of binary vector constructs were prepared utilising the petunia *F3'5'H* cDNA fragments and various promoter and terminator fragments. The chimaeric petunia *F3'5'H* genes had proved successful in carnation and petunia leading to detectable intact *F3'5'H* transcripts (as detected by Northern blot analysis) and to the production of delphinidin or delphinidin-based molecules pigments. Table 5 summarises the list of binary vector constructs containing petunia *F3'5'H* cDNA fragments.

**TABLE 5** Summary of chimaeric petunia *F3'5'H* gene expression cassettes contained in binary vector constructs used in the transformation of roses (see Table 4 for an explanation of abbreviations).

PLASMID	<i>F3'5'H</i> GENE	SELECTABLE GENE	MARKER
PCGP1452	<i>AmCHS 5': petH1: peid8 3'</i>	35S 5': <i>SURB</i>	
PCGP1453	<i>Mac: petH1: mas 3'</i>	35S 5': <i>SURB</i>	
PCGP1457	<i>peid8 5': petH1: peid8 3'</i>	35S 5': <i>SURB</i>	
PCGP1461	<i>short petLS 5': petH1: petLS 3'</i>	35S 3': <i>SURB</i>	
PCGP1616	<i>petRT 5': petH1: nos 3'</i>	35S 5': <i>SURB</i>	
PCGP1638	<i>CaMV 35S: petH1: ocs 3'</i>	35S 5': <i>SURB</i>	
PCGP1623	<i>mas 35S: petH1: ocs 3'</i>	35S 5': <i>SURB</i>	
PCGP1860	<i>RoseCHS 5': petH1: nos 3'</i>	35S 5': <i>SURB</i>	
PCGP2123	<i>CaMV 35S: petH2: ocs 3'</i>	35S 5': <i>SURB</i>	

*Isolation of petunia F3'5'H cDNA clones (petH1 and petH2)*

The isolation and characterization of cDNA clones of petunia *F3'5'H* (*petH1* and *petH2*) contained in PCGP602 (Figure 2) and PCGP175 (Figure 3) respectively (SEQ ID NO:1 and SEQ ID NO:3, respectively) have been described in International Patent Application No. PCT/AU92/00334 and Holton *et al.* (1993a, *supra*).

The plasmids PCGP601 (Figure 2), PCGP602 (Figure 2), PCGP176 (Figure 2) contain homologs of the petunia *petH1 F3'5'H* cDNA clone. The plasmid PCGP601 contains a petunia *F3'5'H petH1* homolog that includes 52bp of 5' untranslated sequence. The plasmid PCGP602 contains a petunia *F3'5'H petH1* homolog that includes 125bp of 5' untranslated sequence (SEQ ID NO:1). The plasmid PCGP176 (described in Holton *et al.*, 1993a *supra*) contains a petunia *F3'5'H petH1* homolog that includes 27bp of 5' untranslated sequence and a further ~127bp of 3' untranslated sequence over the petunia *F3'5'H petH1* cDNA clone in PCGP602.



- 60 -

**Construction of pCGP1303 (*petHf1* in pUC19 backbone)**

The petunia *F3'5'H* cDNA clone contained in the plasmid pCGP601 (described above) (Figure 2) included 52 bp of 5' untranslated sequence and 141 bp of 3' untranslated sequence including 16 bp of the poly A tail. The plasmid pCGP601 (Figure 2) was firstly  
5 linearized by digestion with the restriction endonuclease *BspHI*. The ends were repaired and the petunia *F3'5'H petHf1* cDNA clone was released upon digestion with the restriction endonuclease *FspI*. The *BspHI* recognition sequence encompasses the putative translation initiating codon and the *FspI* recognition sequence commences 2 bp downstream from the stop codon. The 1.6 kb fragment containing the petunia *F3'5'H*  
10 *petHf1* cDNA clone was purified and ligated with repaired *EcoRI* ends of pUC19 (New England Biolabs). Correct insertion of the fragment was established by restriction endonuclease analysis of DNA isolated from ampicillin-resistant transformants. The resulting plasmid was designated as pCGP1303 (Figure 4).

15 **Construction of pCGP627 (short *petHf1* in pBluescript backbone)**

The plasmid pCGP176 (Holton *et al.*, 1993a, *supra*) (Figure 2) was digested with the restriction endonuclease *SpeI* and *EcoRI*. The ends were then repaired and allowed to religate. The resulting plasmid was designated as pCGP627 and contained the identical cDNA clone as in pCGP176 except that the restriction endonuclease sites *PstI*, *BamHI* and  
20 *SmaI* were removed from the multi-cloning site of the pBluescript vector at the 5' end of the cDNA clone.

**The binary vector pCGP1452 (*AmCHS* 3'; *petHf1*; *petD8* 3')**

The plasmid pCGP1452 (Figure 5) contains a chimaeric petunia *F3'5'H (petHf1)* gene under the control of a promoter fragment from the *Antirrhinum majus* chalcone synthase gene (*CHS*) (Sommer and Saedler, 1986, *supra*) with a terminator fragment from the petunia phospholipid transfer protein (PLTP) gene (*petD8* 3') (Holton, 1992, *supra*). The chimeric petunia *F3'5'H* cassette is in a tandem orientation with respect to the 35S 5':  
25 *SuRB* gene of the binary vector, pWTT2132 (DNA Plant Technologies, USA = DNAP)  
30 (Figure 6).

### Intermediates in the preparation of the binary pCGP1452

#### The binary vector pWTT2132

The binary vector plasmid pWTT2132 (DNAP) (Figure 6) contains a chimeric gene comprised of a 35S 5' promoter sequence (Riack et al., 1980, *supra*), ligated with the coding region and terminator sequence for acetylacrylate synthase (ALS) gene from the *SuaB* locus of tobacco (Lee et al., 1988, *supra*). An ~60bp 5' untranslated leader sequence from the petunia chlorophyll *a/b* binding protein gene (Cab 22 gene) (Harper et al., 1988, 212: 182-190, 1988) is included between the 35S 5' promoter fragment and the *SuaB* sequence.

Construction of pCGP725 (AmCHS 5': petHT: petD8 3' in pBluescript)

A chimeric petunia *F3'5'H* gene under the control *Antirrhinum majus CHS* (*AmCHS 5'*) promoter with a petunia PLTP terminator (*petD8 3'*) was constructed by cloning the 1.6kb *BclI/FspI* petunia *F3'5'H* (*petHT*) fragment from pCGP602 (Holton et al., 1993a, *supra*) (Figure 2) between a 1.2 kb *Antirrhinum majus CHS* gene fragment 5' to the site of translation initiation (Sommer and Siedler, 1986, *supra*) and a 0.7 kb *SmaI/XhoI* PLTP fragment (*petD8 3'*) from pCGP13ABam (Holton, 1992, *supra*), 3' to the deduced stop codon. The resulting plasmid in a pBluescript II KS (Stratagene, USA) backbone vector was designated pCGP725 (Figure 7).

Construction of pCGP485 and pCGP1452 (AmCHS 5': petHT: petD8 3' binary vectors)

The chimeric *F3'5'H* gene from pCGP725 (Figure 7) was cloned into the binary vector pCGN1547 containing an *npII* selectable marker gene cassette (McBride and Summerfelt *Plant Molecular Biology* 14: 269-276, 1990) to create pCGP485. A 3.5 kb fragment containing the *AmCHS 5'*; *petHT*; *petD8 3'* cassette was released upon digestion of pCGP485 with the restriction endonuclease *PstI*. The overhanging ends were repaired and the purified 3.5 kb fragment was ligated with *SmaI* ends of the binary vector, pWTT2132 (DNAP). Correct insertion of the fragment in a tandem orientation with respect to the 35S 5': *SuaB* selectable marker gene cassette was established by restriction endonuclease analysis of plasmid DNA isolated from tetracycline-resistant transformants. The plasmid was designated as pCGP1452 (Figure 5).

Plant transformation with pCGP1452

The T-DNA contained in the binary vector plasmid pCGP1452 (Figure 5) was introduced into rose via *Agrobacterium*-mediated transformation.

5

The binary vector pCGP1453 (Mac: petHf1: mas 3')

The plasmid pCGP1453 (Figure 8) contains a chimeric petunia *F3'5'H* (*petHf1*) gene under the control of a *Mac* promoter (Comai *et al.*, 1990, *supra*) with a terminator fragment from the mannopine synthase gene of *Agrobacterium* (*mas 3'*). The chimeric petunia *F3'5'H* cassette is in a tandem orientation with respect to the 35S 5': *SuRB* selectable marker gene cassette of the binary vector, pWTT2132 (DNAP) (Figure 6).

10

A 3.9 kb fragment containing the *Mac: petHf1: mas 3'* gene was released from the plasmid pCGP628 (described in International Patent Application No. PCT/AU94/00265) upon digestion with the restriction endonuclease *Pst*I. The overhanging ends were repaired and the purified fragment was ligated with *Sma*I ends of pWTT2132 (DNAP). Correct insertion of the *Mac: petHf1: mas 3'* gene in a tandem orientation with respect to the 35S 5': *SuRB* selectable marker gene cassette was established by restriction endonuclease analysis of plasmid DNA isolated from tetracycline-resistant transformants. The plasmid was designated as pCGP1453 (Figure 8).

15

20

Plant transformation with pCGP1453

The T-DNA contained in the binary vector plasmid pCGP1453 (Figure 8) was introduced into rose via *Agrobacterium*-mediated transformation.

25

The binary vector pCGP1457 (petD8.5'; petH1'; pet D8 3')

The plasmid pCGP1457 (Figure 9) contains a chimeric petunia F3'5'H (petH1) gene under the control of a promoter fragment from the petunia *PLTP* gene (petD8 5') with a terminator fragment from the petunia *PLTP* gene (petD8 3'). The chimeric petunia F3'5'H cassette is in a tandem orientation with respect to the 35S 5' : 5SRRB gene of the binary vector, pWTT2132 (DNAP) (Figure 6).

Intermediates in the preparation of the binary vector pCGP1457

Isolation of petunia D8 genomic clone

Preparation of *P. hybrida* cv. OGB (Old Glory Blue) genomic library in  $\lambda$ 2001

A genomic DNA library was constructed from *Petunia hybrida* cv. OGB DNA in the vector  $\lambda$ 2001 (Kam *et al.*, *Gene* 32: 217-224, 1984) using a *Sma*3A partial digestion of the genomic DNA as described in Holton, 1992 (*supra*). Screening of the OGB genomic library for the petunia D8 gene was as described in Holton, 1992, *supra*.

Isolation of D8 genomic clone OGB2.6

PCR was performed in order to find a non-mutant genomic clone representing D8. Oligo #2 (5' to 3' GTTCTGAGGAAAGATATACAAAT) (SEQ ID NO:6) and Oligo #4 (5' to 3' CAAGATCGTAGGACTGCATG) (SEQ ID NO:7) were used to amplify D8 gene fragments, across the intron region, using 4  $\mu$ L of phage suspension from the clones isolated from the primary screening of the OGB genomic library. The reactions were carried out in a total volume of 50  $\mu$ L containing 1 x Amplification buffer (Cetus), 0.2 mM dNTP mix, <1  $\mu$ g of template DNA, 50 pmoles of each primer and 0.25  $\mu$ L of Taq polymerase (5 units/ $\mu$ L - Cetus). The reaction mixtures were overlaid with 30  $\mu$ L of mineral oil and temperature cycled using a Gene Machine (Innovonics). The reactions were cycled 30 times using the following conditions: 94°C for 1 minute, 55°C for 50 seconds, 72°C for 2 minutes. One quarter of each PCR reaction was run on an agarose gel using TAE running buffer.

Three clones,  $\lambda$ OGB-2.4,  $\lambda$ OGB-2.5, and  $\lambda$ OGB-2.6, gave fragments of approximately 1 kb whereas the mutant clone,  $\lambda$ OGB-3.2 (described in Holton, 1992, *supra*), had produced a product of 1.25 kb. The  $\lambda$ OGB-2.6 clone was chosen for further analysis.

**pCGP382**

The genomic clone,  $\lambda$ OGB-2.6, contained a single 3.9 kb *Xba*I fragment that hybridized with the D8 cDNA. This *Xba*I fragment was isolated and purified and ligated with the *Xba*I ends of pBluescriptII KS- (Stratagene, USA). Restriction mapping of this clone revealed an internal *Pst*I site 350 bp from the 3' end. However, the "mutant" genomic clone in pCGP13, had an internal *Pst*I near the putative initiating "ATG" of the coding region (approximately 1.5 kb from its 3' end). The difference in the position of the *Pst*I site in both clones suggested that the  $\lambda$ OGB-2.6 *Xba*I fragment did not contain the whole genomic sequence of D8. A Southern blot was performed on *Pst*I digested  $\lambda$ OGB-2.6 DNA, and a fragment of 2.7 kb was found to hybridize with the D8 cDNA. Restriction endonuclease mapping confirmed that this fragment contained the 3' coding region and flanking sequences.

In order to obtain a fragment containing the whole D8 genomic sequence, a number of cloning steps were undertaken. The  $\lambda$ OGB-2.6 *Pst*I fragment of 2.7 kb was purified and ligated with *Pst*I ends of pBluescriptII KS- (Stratagene, USA). The resultant clone was digested with *Xba*I to remove the 350 bp *Pst*I/*Xba*I fragment. This fragment was replaced by the 3.9 kb *Xba*I fragment from  $\lambda$ OGB-2.6 to produce the plasmid pCGP382.

A 3.2 kb fragment containing the promoter region from the D8 2.6 gene in pCGP382 was released upon digestion with the restriction endonucleases *Hind*III and *Nco*I. The fragment was purified and ligated with the 4.8 kb *Nco*I/*Hind*III fragment of pJB1 (Bodean, Molecular and genetic regulation of Bronze-2 and other maize anthocyanin genes. Dissertation, Stanford University, USA, 1994) to produce pCGP1101 containing a *petD85': GUS: nos 3'* cassette.

A 1.6 kb petunia *F3'5'H petHf1* fragment was released from the plasmid pCGP602 (Holton *et al.*, 1993a, *supra*) (SEQ ID NO:1) (Figure 2) upon digestion with the restriction endonucleases *Bsp*HI and *Bam*HI. The fragment was purified and ligated with the 6.2 kb *Nco*I/*Bam*HI fragment of pCGP1101 to produce pCGP1102 containing a *petD8 5': petHf1: nos 3'* expression cassette.

A 0.75 kb *Bam*HI *petD8* 3' fragment (Holton, 1992, *supra*) was purified from the plasmid pCGP13ABamHI and ligated with *Bam*HI/*Bgl*II ends of pCGP1102 to produce the plasmid pCGP1107 containing a *petD8* 5': *petHf1*: *petD8* 3' expression cassette.

The plasmid pCGP1107 was linearised upon digestion with the restriction endonuclease *Xba*I. The overhanging ends were repaired and then the 5.3 kb fragment containing the *petD8* 5': *petHf1*: *petD8* 3' expression cassette was released upon digestion with the restriction endonuclease *Pst*I. The fragment was purified and ligated with *Sma*I/*Pst*I ends of the binary vector pWTT2132 (DNAP) (Figure 6). Correct insertion of the *petD8* 5': *petHf1*: *petD8* 3' gene in a tandem orientation with respect to the 35S 5': *SuRB* selectable marker gene cassette was established by restriction endonuclease analysis of plasmid DNA isolated from tetracycline-resistant transformants. The plasmid was designated as pCGP1457 (Figure 9).

#### Plant transformation with pCGP1457

The T-DNA contained in the binary vector plasmid pCGP1457 (Figure 9) was introduced into rose via *Agrobacterium*-mediated transformation.

#### The binary vector pCGP1461 (short *petLS* 5': *petHf1*: *petLS* 3')

The plasmid pCGP1461 (Figure 10) contains a chimeric petunia *F3'5'H* (*petHf1*) gene under the control of a promoter fragment from the petunia flavonol synthase (*FLS*) gene (short *petLS* 5') with a terminator fragment from the petunia *FLS* gene (*petLS* 3'). The chimeric petunia *F3'5'H* gene is in a tandem orientation with respect to the 35S 5': *SuRB* gene of the binary vector, pWTT2132 (Figure 6).

*Intermediates in the preparation of the binary vector pCGP1461*

*Isolation of petunia FLS gene*

*Preparation of P. hybrida cv. Th7 genomic library*

A *P. hybrida* cv. Th7 genomic library was prepared according to Sambrook *et al.* (1989, *supra*) using a *Sau3A* partial digestion of the genomic DNA. The partially digested DNA was cloned into EMBL-3 lambda vector (Stratagene, USA).

The Th7 genomic DNA library was screened with <sup>32</sup>P-labelled fragments of a petunia *FLS* cDNA clone (Holton *et al.*, *Plant J.* 4: 1003-1010, 1993b) using high stringency conditions.

Two genomic clones (*FLS2* and *FLS3*) were chosen for further analysis and found to contain sequences upstream of the putative initiating methionine of the petunia *FLS* coding region with *FLS2* containing a longer promoter region than *FLS3*.

*pCGP486*

A 6 kb fragment was released upon digestion of the genomic clone *FLS2* with the restriction endonuclease *XhoI*. The fragment containing the short petunia *FLS* gene was purified and ligated with *XhoI* ends of pBluescript SK (Stratagene, USA). Correct insertion of the fragment was established by restriction endonuclease analysis of DNA isolated from ampicillin-resistant transformants. The resulting plasmid was designated as pCGP486.

*pCGP487*

A 9 kb fragment was released upon digestion of the genomic clone *FLS3* with the restriction endonuclease *XhoI*. The fragment containing the petunia *FLS* gene was purified and ligated with *XhoI* ends of pBluescript SK (Stratagene, USA). Correct insertion of the fragment was established by restriction endonuclease analysis of DNA isolated from ampicillin-resistant transformants. The resulting plasmid was designated as pCGP487.

**pCGP717**

A 2.2 kb petunia *FLS* promoter upstream from the putative translational initiation site was released from the plasmid pCGP487 upon digestion with the restriction endonucleases *XhoI* and *PstI*. The fragment generated was purified and ligated with *XhoI/PstI* ends of pBluescript II KS+ (Stratagene, USA). Correct insertion of the fragment was established by restriction endonuclease analysis of DNA isolated from ampicillin-resistant transformants. The resulting plasmid was designated as pCGP717.

**pCGP716**

A 0.95 kb petunia *FLS* terminator fragment downstream from the putative translational stop site was released from the plasmid pCGP487 upon digestion with the restriction endonucleases *HindIII* and *SacI*. The fragment generated was purified and ligated with *HindIII/SacI* ends of pBluescript II KS+ (Stratagene, USA). Correct insertion of the fragment was established by restriction endonuclease analysis of DNA isolated from ampicillin-resistant transformants. The resulting plasmid was designated as pCGP716.

**Construction of pCGP493 (short petFLS 5'-petFLS3' expression cassette)**

A 1.8 kb fragment containing the short petunia *FLS* promoter fragment was amplified by PCR using the plasmid pCGP717 as template and the T3 primer (Stratagene, USA) and an *FLS*-Neo primer (5' AAA ATC GAT ACC ATG GTC TTT TTT TCT TTG TCT ATA C 3') (SEQ ID NO:19). The PCR product was digested with the restriction endonucleases *XhoI* and *ClaI* and the purified fragment was ligated with *XhoI/ClaI* ends of pCGP716. Correct insertion of the fragment was established by restriction endonuclease analysis of DNA isolated from ampicillin-resistant transformants. The resulting plasmid was designated as pCGP493.

**Construction of pCGP497 (short petFLS 5'-petHT1: petFLS3' expression cassette)**

The petunia *F3'5'H* (*petHT1*) cDNA clone was released from the plasmid pCGP627 (described above) upon digestion with the restriction endonucleases *BspHI* and *RspI*. The *BspHI* recognition sequence encompasses the putative translation initiating codon and the *RspI* recognition sequence commences 2 bp downstream from the stop codon. The petunia



- 68 -

*F3'5'H petHf1* fragment generated was purified and ligated with *Clal* (repaired ends)/*NcoI* ends of the plasmid pCGP493. Correct insertion of the fragment was established by restriction endonuclease analysis of DNA isolated from ampicillin-resistant transformants. The resulting plasmid was designated as pCGP497.

5

**Construction of pCGP1461 (short *petFLS* 5': *petHf1*: *petFLS*3' binary vector)**

The plasmid pCGP497 was linearised upon digestion with the restriction endonuclease *SacI*. The overhanging ends were repaired and a 4.35 kb fragment containing the *short petFLS* 5': *petHf1*: *petFLS*3' gene expression cassette was released upon digestion with the  
10 restriction endonuclease *KpnI*. The fragment generated was purified and ligated with *PstI* (ends repaired)/*KpnI* ends of the binary vector pWTT2132 (DNAP) (Figure 6). Correct insertion of the fragment in a tandem orientation with respect to the 35S 5': *SuRB* selectable marker gene cassette of pWTT2132 was established by restriction endonuclease analysis of plasmid DNA isolated from tetracycline-resistant transformants. The resulting  
15 plasmid was designated as pCGP1461 (Figure 10).

**Plant transformation with pCGP1461**

The T-DNA contained in the binary vector plasmid pCGP1461 (Figure 10) was introduced into rose via *Agrobacterium*-mediated transformation.

20

**The binary vector pCGP1616 (*petRT* 5': *petHf1*: *nos* 3')**

The plasmid pCGP1616 (Figure 11) contains a chimeric petunia *F3'5'H* (*petHf1*) gene under the control of a promoter fragment from the *P. hybrida* 3RT gene (*petRT* 5') (Brugliera, 1994, *supra*) with a terminator fragment from the nopaline synthase gene (*nos*  
25 3') of *Agrobacterium* (Depicker, *et al.*, 1982, *supra*). The chimeric petunia *F3'5'H* cassette is in a tandem orientation with respect to the 35S 5': *SuRB* gene of the binary vector, pWTT2132 (DNAP) (Figure 6).

# **Intermediates in the preparation of the binary vector pCGP1616**

## **Isolation of petunia 3RT gene**

## **P. hybrida cv. Th7 genomic DNA library construction in EMBL3**

5 A *Petunia hybrida* cv. Th7 genomic library was prepared according to Sambrook *et al.* 1989, *supra* using a *Sau3A* partial digestion of the genomic DNA. The partially digested DNA was cloned into EMBL3-3 lambda vector (Stratagene, USA). Screening of the Th7 genomic library for the petunia 3RT gene was as described in Bruggliera, 1994, *supra*.

10 A 3 kb fragment containing the *petRT* 5': *petHT*: *nos* 3' cassette was released from the plasmid pCGP846 (described in Bruggliera, 1994, *supra*) upon digestion with the restriction endonucleases *Pst*I and *Bam*HI. The purified fragment was ligated with *Pst*I/*Bam*HI ends of pWTT2132 (DNAP) (Figure 6). Correct insertion of the fragment in a tandem orientation with respect to the 35S 5': *SuRB* selectable marker gene cassette was established by restriction endonuclease analysis of plasmid DNA isolated from tetacycline-resistant transformants. The plasmid was designated as pCGP1616 (Figure 11).

## **Plant transformation with pCGP1616**

20 The T-DNA contained in the binary vector plasmid pCGP1616 (Figure 11) was introduced into rose via *Agrobacterium*-mediated transformation.

## **The binary vector pCGP1623 (mus/35S: *petHT*: *ocs* 3')**

25 The plasmid pCGP1623 (Figure 12) contains a chimeric petunia *F3'5'H* (*petHT*) gene under the control of the expression cassette contained in pKIWI101 (Janssen and Gardner, 1989, *supra*) consisting of a promoter fragment from the cauliflower mosaic virus 35S gene (35S 5') with an enhancing sequence from the promoter of the mannopine synthase gene (*mas*) of *Agrobacterium* and a terminator fragment from the octopine synthase gene of *Agrobacterium* (*ocs* 3'). The chimeric petunia *F3'5'H* cassette is in a tandem orientation with respect to the 35S 5': *SuRB* gene of the binary vector, pWTT2132 (DNAP) (Figure 6).

- 70 -

***Intermediates in the preparation of the binary vector pCGP1623***

The ~1.6 kb fragment of the petunia *F3'5'H petHf1* cDNA clone contained in the plasmid pCGP1303 (Figure 4) was released upon digestion with the restriction endonucleases *BspHI* and *SmaI*. The petunia *F3'5'H petHf1* fragment was purified and ligated with a ~5.9 kb *NcoI/EcoRI* (repaired ends) fragment of pKIWI101 (Janssen and Gardner, 1989, *supra*) to produce the plasmid pCGP1619.

A partial digest of the plasmid pCGP1619 with the restriction endonuclease *XhoI* released a 4.9 kb fragment containing the *mas/35S: petHf1: ocs 3'* expression cassette. The fragment was purified and ligated with *SaII* ends of pWTT2132 (DNAP) (Figure 6). Correct insertion of the fragment in a tandem orientation with respect to the 35S 5': *SuRB* selectable marker gene cassette was established by restriction endonuclease analysis of plasmid DNA isolated from tetracycline-resistant transformants. The plasmid was designated as pCGP1623 (Figure 12).

***Plant transformation with pCGP1623***

The T-DNA contained in the binary vector plasmid pCGP1623 (Figure 12) was introduced into rose via *Agrobacterium*-mediated transformation.

***The binary vector pCGP1638 (35S 5': petHf1: ocs 3')***

The plasmid pCGP1638 (Figure 13) contains a chimeric petunia *F3'5'H (petHf1)* gene under the control of a *CaMV 35S* promoter (35S 5') with an octopine synthase terminator (*ocs 3'*). A ~60 bp 5' untranslated leader sequence from the petunia chlorophyll a/b binding protein gene (*Cab 22* gene) (Harpster *et al.*, 1988, *supra*) is included between the *CaMV 35S* promoter fragment and the petunia *F3'5'H petHf1* cDNA clone. The chimeric petunia *F3'5'H* cassette is in a tandem orientation with respect to the 35S 5': *SuRB* selectable marker gene cassette of the binary vector, pWTT2132 (Figure 6).

# Intermediates in the preparation of the binary vector pCGP1638

## Construction of pCGP1273

The plasmid pCGP1273 was constructed by subcloning a ~3kb *HindIII/HpaI* fragment containing 35S 5' : *GUS*: *ocs* 3' gene from the binary vector pJ13499 (Jones *et al*, *Transgenic Research*, 1: 285-297, 1992) with the *HindIII/SmaI* ends of the plasmid pBluescript KS II (+) (Stratagene, USA).

## Construction of pCGP1634

A ~3kb *HindIII/BamHI* fragment containing the 35S 5' : *GUS*: *ocs* 3' gene from pCGP1273 was then isolated and ligated with the *HindIII/BamHI* ends of the cloning vector pUC19 (New England Biolabs) to create the plasmid pCGP1634.

## Construction of pCGP1636

The *GUS* fragment from the plasmid pCGP1634 was removed by digesting pCGP1634 with the restriction endonucleases *NcoI* and *XbaI* and purifying the ~3.7kb fragment containing the 35S 5' promoter fragment, the *ocs* 3' terminator fragment and the pUC19 vector backbone.

The petunia *F3.5.H.peatH1* cDNA clone was released from pCGP1303 (figure 4) upon digestion with the restriction endonucleases *BspHI* and *XbaI*. The resulting ~1.6kb fragment was purified and ligated with the ~3.7kb *NcoI/XbaI* fragment from pCGP1634. Correct insertion of the petunia *F3.5.H.peatH1* fragment was established by restriction endonuclease analysis of plasmid DNA isolated from ampicillin-resistant transformants. The resulting plasmid containing a 35S 5' : *peatH1*: *ocs* 3' gene was designated pCGP1636.

## Construction of pCGP1638

The 35S 5' : *peatH1*: *ocs* 3' gene from the plasmid pCGP1636 was released upon digestion of pCGP1636 with the restriction endonucleases *PstI* and *EcoRI*. The ends were repaired and the ~2.6kb fragment was purified and ligated with the *Sam* ends of the binary vector, pWTT12132 (DNAP). Correct insertion of the 35S 5' : *peatH1*: *ocs* 3' gene in a tandem orientation with respect to the 35S 5' : *GUS* selectable marker gene cassette was established by restriction endonuclease analysis of plasmid DNA isolated from

- 72 -

tetracycline-resistant transformants. The plasmid was designated as pCGP1638 (Figure 13).

Plant transformation with pCGP1638

- 5 The T-DNA contained in the binary vector plasmid pCGP1638 (Figure 13) was introduced into rose via *Agrobacterium*-mediated transformation.

The binary vector pCGP1860 (*RoseCHS* 5': *petHf1*: *nos* 3')

- 10 The plasmid pCGP1860 (Figure 14) contains a chimeric petunia *F3'5'H* (*petHf1*) gene under the control of a promoter fragment from the chalcone synthase gene of *Rosa hybrida* (*RoseCHS* 5') with a terminator fragment from the nopaline synthase gene of *Agrobacterium* (*nos* 3'). The chimeric petunia *F3'5'H* cassette is in a tandem orientation with respect to the 35S 5': *SuRB* selectable marker gene cassette of the binary vector, pWTT2132 (DNAP) (Figure 6).

15

Intermediates in the preparation of the binary vector pCGP1860

Isolation of Rose *CHS* promoter

A rose genomic DNA library was prepared from genomic DNA isolated from young leaves of *Rosa hybrida* cv. Kardinal.

20

- The Kardinal genomic DNA library was screened with <sup>32</sup>P-labelled fragment of rose *CHS* cDNA clone contained in the plasmid pCGP634. The rose *CHS* cDNA clone was isolated by screening of a petal cDNA library prepared from RNA isolated from petals of *Rosa hybrida* cv Kardinal (Tanaka *et al.*, 1995, *supra*) using a petunia *CHS* cDNA fragment as probe (clone *1F11* contained in pCGP701, described in Brugliera *et al.*, 1994, *supra*).  
25 Conditions are as described in Tanaka *et al.*, 1995 (*supra*).

- A rose genomic clone (*roseCHS20λ*) was chosen for further analysis and found to contain ~6.4 kb of sequence upstream of the putative initiating methionine of the rose *CHS* coding region.  
30

An ~6.4 kb fragment upstream from the translational initiation site was cloned into pBluescript KS (-) (Stratagene) and the plasmid was designated as pCGP1114.

5 The plasmid pCGP1114 was digested with the restriction endonucleases *HindIII* and *NotI* to release a 2.7-3.0kb fragment which was purified and ligated with the *HindIII/SmaI* ends of pUC19 (New England Biolabs). Correct insertion of the rose *CHS* promoter fragment was established by restriction endonuclease analysis of DNA isolated from ampicillin-resistant transformants. The resulting plasmid was designated as pCGP1116. The DNA sequence of the rose *CHS* promoter fragment was determined using pCGP1116 as template (SEQ ID NO:5).

10 Construction of pCGP197 (RoseCHS 5': GUS : nos 3' in pUC18 backbone)  
An ~3.0 kb fragment containing the rose chalcone synthase promoter (RoseCHS 5') was released from the plasmid pCGP1116 upon digestion with the restriction endonucleases *HindIII* and *Asp718*. The fragment was purified and ligated with a *HindIII/Asp718* fragment from pJB1 (Bodeau, 1994, *supra*) containing the vector backbone, β-glucuronidase (*GUS*) and nos 3' fragments. Correct insertion of the rose *CHS* promoter fragment upstream of the *GUS* coding sequence was established by restriction endonuclease analysis of DNA isolated from ampicillin-resistant transformants. The resulting plasmid was designated as pCGP197.

25 Construction of pCGP200 (RoseCHS 5': petH1 : nos 3' in pUC18 backbone)  
A 1.8 kb fragment containing the petunia *F3'5'H (petH1)* fragment was released from the plasmid pCGP1303 (described above) (Figure 4) upon digestion with the restriction endonucleases *BspHI* and *SacI*. The petunia *F3'5'H petH1* fragment was purified and ligated with *NotI/SacI* ends of pCGP197. Correct insertion of the petunia *F3'5'H petH1* fragment between the rose *CHS* promoter and nos 3' fragments was established by restriction endonuclease analysis of DNA isolated from ampicillin-resistant transformants. The resulting plasmid was designated as pCGP200.

Construction of pCGP1860 (RoseCHS 5': petHf1: nos 3' in a binary vector)

An ~4.9 kb fragment containing the *RoseCHS* 5': *petHf1*: *nos* 3' cassette was released from the plasmid pCGP200 upon digestion with the restriction endonuclease *Bgl*II. The fragment was purified and ligated with *Bam*HI ends of the binary vector, pWTT2132 (DNAP) (Figure 6). Correct insertion of the fragment in a tandem orientation with respect to the 35S 5': *SuRB* selectable marker gene cassette of pWTT2132 was established by restriction endonuclease analysis of plasmid DNA isolated from tetracycline-resistant transformants. The resulting plasmid was designated as pCGP1860 (Figure 13).

10 Plant transformation with pCGP1860

The T-DNA contained in the binary vector plasmid pCGP1860 (Figure 14) was introduced into rose via *Agrobacterium*-mediated transformation.

The binary vector pCGP2123 (CaMV 35S: petHf2: ocs 3')

15 The plasmid pCGP2123 (Figure 15) contains a chimeric petunia *F3'5'H* (*petHf2*) gene under the control of a CaMV35S promoter with a terminator fragment from the octopine synthase gene of *Agrobacterium* (*ocs* 3'). The chimeric petunia *F3'5'H* cassette is in a tandem orientation with respect to the 35S 5': *SuRB* selectable marker gene cassette of the binary vector, pCGP1988 (Figure 16).

20

Intermediates in the preparation of the binary vector pCGP2123

Construction of pCGP1988 (a derivative of the binary vector, pWTT2132)

The binary vector pCGP1988 (Figure 16) is based on binary vector pWTT2132 (DNAP) (Figure 6) but contains the multi-cloning site from pNEB193 (New England Biolabs). The plasmid pNEB193 was firstly linearized by digestion with the restriction endonuclease *Eco*RI. The overhanging ends were repaired and the multi-cloning fragment was released upon digestion with the restriction endonuclease *Pst*I. The fragment was purified and ligated with *Sal*I (ends repaired)/*Pst*I ends of the binary vector pWTT2132 (DNAP). Correct insertion of the multi-cloning fragment into pWTT2132 was established by restriction endonuclease analysis of plasmid DNA isolated from tetracycline-resistant transformants. The resulting plasmid was designated as pCGP1988 (Figure 16).

30

- Construction of pCGP2000 (CaMV 35S promoter fragment in Bluescript)
- The plasmid pCGP2000 was an intermediate plasmid containing a cauliflower mosaic virus (CaMV) 35S promoter fragment in a pBluescript SK (Stratagene, USA) backbone.
- The CaMV 35S promoter fragment from pK1W1101 (Janssen and Gardner, 1989, *supra*) was released upon digestion with the restriction endonucleases *Xba*I and *Pst*I. The ~0.35 kb fragment generated was purified and ligated with *Xba*I/*Pst*I ends of the vector pBluescript SK. Correct insertion of the fragment was established by restriction endonuclease analysis of plasmid DNA isolated from ampicillin-resistant transformants. The plasmid was designated as pCGP2000.
- 10
- Construction of pCGP2105 (CaMV 35S 5' and *ocs* 3' fragments in pBluescript)
- The plasmid pCGP2105 (Figure 17) contained a CaMV 35S promoter fragment along with a terminator fragment from the octopine synthase gene of *Agrobacterium* (*ocs* 3') both from pK1W1101 (Janssen and Gardner, 1989, *supra*).
- 15
- The *ocs* 3' fragment from pK1W1101 (Janssen and Gardner, 1989, *supra*) was isolated by firstly digesting the plasmid pK1W1101 with the restriction endonuclease *Eco*RI, followed by repair of the overhanging ends, and finally by digestion with the restriction endonuclease *Xho*I to release a 1.6 kb fragment. This fragment was then ligated with *Hinc*II/*Xho*I ends of pCGP2000. Correct insertion of the fragment was established by restriction endonuclease analysis of plasmid DNA isolated from ampicillin-resistant transformants. The plasmid was designated pCGP2105 (Figure 17).
- 20
- Construction of pCGP2109 (CaMV 35S: *peHf2*: *ocs* 3' gene in pBluescript)
- The plasmid pCGP2109 contained the CaMV 35S: *peHf2*: *ocs* 3' expression gene cassette in a pBluescript backbone.
- 25
- The 1.8 kb petunia *F3'5'H peHf2* cDNA clone was released from pCGP175 (Holtan *et al.*, 1993a, *supra*) upon digestion with the restriction endonucleases *Xba*I and *Sap*I. The overhanging ends were repaired and the purified fragment was ligated with *Pst*I (ends
- 30



- 76 -

repaired/*EcoRV* ends of pCGP2105 (described above) (Figure 17). Correct insertion of the fragment was established by restriction endonuclease analysis of plasmid DNA isolated from ampicillin-resistant transformants. The plasmid was designated pCGP2109.

5 **Construction of pCGP2123 (*CaMV* 35S: *petHf2*: *ocs* 3' binary vector)**

The *CaMV* 35S: *petHf2*: *ocs* 3' cassette was released from pCGP2109 upon digestion with the restriction endonucleases *Asp*718 and *Xba*I. The overhanging ends were repaired and the resultant ~3.7 kb fragment containing the *CaMV* 35S: *petHf2*: *ocs* 3' gene was purified and ligated with repaired ends of *Asp*718 of the binary vector, pCGP1988 (Figure 16).

- 10 Correct insertion of the *CaMV* 35S: *petHf2*: *ocs* 3' gene in a tandem orientation with respect to the 35S 5': *SuRB* selectable marker gene cassette was established by restriction endonuclease analysis of plasmid DNA isolated from tetracycline-resistant transformants. The plasmid was designated as pCGP2123 (Figure 15).

15 **Plant transformation with pCGP2123**

The T-DNA contained in the binary vector plasmid pCGP2123 (Figure 15) was introduced into rose via *Agrobacterium*-mediated transformation.

**EXAMPLE 5**

20 **Analysis of transgenic roses**

The transgenic roses produced in the experiments described in Example 4 were grown to flowering. Flowers were collected and the colors of the petals were coded using the Royal Horticultural Society Colour Charts (RHSCC). The anthocyanins were extracted and the anthocyanidins (specifically the presence of delphinidin or delphinidin-based molecules) analysed by TLC and/or HPLC analysis. Total RNA was also isolated from petal tissue and Northern blot analysis was used to detect transcripts of petunia *F3'5'H* transgenes, endogenous rose *CHS* gene and *SuRB* transgene. The results of the transgenic analysis are summarised in Table 6.

- 77 -

Although over 250 transgenic Kardinal roses were produced (Table 6) none produced flowers with a change in color. TLC and/or HPLC analysis failed to detect accumulation of delphinidin or delphinidin-based molecules confirming the absence of efficient  $P3'5'H$  activity. Subsequent Northern analysis on total RNA isolated from petal tissue of these transgenic roses revealed either no detectable intact petunia  $P3'5'H$  (*peHf1*) or *peHf2* transcripts, or in some cases (see footnotes), degraded transcripts. Hybridization of the same membranes with the selectable marker gene (*SuaB*) or with an endogenous rose *CHS* cDNA probe revealed discrete hybridizing transcripts indicating that the total RNA isolated was not degraded. The detection of the *SuaB* transgene transcripts confirmed that the roses were transgenic.

TABLE 6 Results of transgenic analysis of rose petals transformed with the T-DNA from various petunia  $P3'5'H$  (*peHf1* or *peHf2*) gene expression cassettes.

PLASMID	$P3'5'H$ GENE	EVENTS	DEL	RNA
pCGP1452	<i>AmCHS 5': peHf1: petD8 3'</i>	34	0/28	0/34
pCGP1453	<i>Mac: peHf1: mas 3'</i>	16	0/14	0/13
pCGP1457	<i>petD8 5': peHf1: petD8 3'</i>	11	0/11	0/11
pCGP1461	<i>short petD8 5': peHf1: petD8 3'</i>	11	0/11	0/11
pCGP1616	<i>peHf1 5': peHf1: nos 3'</i>	4	0/4	0/4
pCGP1623	<i>mas/35S: peHf1: nos 3'</i>	27	0/20	0/12
pCGP1638	<i>CaMV 35S: peHf1: nos 3'</i>	22	0/14	0/14
pCGP1860	<i>RoseCHS 5': peHf1: nos 3'</i>	15	0/13	0/13
pCGP2123	<i>CaMV 35S: peHf2: nos 3'</i>	40	0/26	0/10

EVENTS = number of independent transgenic events produced  
 DEL = number of transgenic events in which delphinidin or delphinidin-based molecules was detected (by TLC or HPLC) in petals over the total number of events analyzed

15

- 78 -

RNA = number of transgenic events in which intact *F3'5'H* (*petHf1* or *petHf2*) transcripts were detected by Northern blot analysis in total RNA isolated from rose petals over the total number of events analyzed

- 1 = Degraded transcripts were detected in 5 of the 34 analyzed
- 5 2 = Degraded transcripts were detected in 8 of the 13 analyzed
- 3 = Degraded transcripts were detected in 8 of the 12 analyzed

The fact that no intact petunia *F3'5'H* (*petHf1* or *petHf2*) transcripts were ever detected in transgenic rose petals transformed with the T-DNAs described (Table 6) suggested a number of possibilities:

1. that the RNA isolated was degraded. This was not the case as the RNA had been stained by ethidium bromide and visualised under UV-light. The intact visible ribosomal RNA bands were used as an indicator of the quality of the RNA isolated. Furthermore the detection of full-length transcripts of the endogenous rose *CHS* and *SuRB* transgenes confirmed that the RNA preparation was not degraded.
2. that there was no initiation of transcription of the chimeric *F3'5'H* genes evaluated. This was a possibility with some of the expression cassettes analysed, as no *F3'5'H* transcripts were detected by Northern analysis. However all of the petunia *F3'5'H* expression cassettes had proven to be functional (ie. result in an intact transcript and result in the production of delphinidin-based pigments) in other plants such as carnation and petunia.
3. that the petunia *F3'5'H* *petHf1* and *petHf2* mRNAs were unstable in roses. This was also a possibility as degraded petunia *F3'5'H* transcripts were detected by Northern analysis in total RNA isolated from petals of some events. However the petunia *petHf1* and *petHf2* mRNAs had been proven to be stable in other plants such as carnation and petunia. Such instability could be due to aberrant translation leading to mRNA turnover, some feature of the sequence inherently unstable in rose cells, some other factor or factors.

30

There was a need therefore to find suitable promoter fragments that would efficiently drive expression of genes in rose petals and find suitable *F3'5'H* sequences that would result in intact transcripts accumulating in rose petals leading to functional *F3'5'H* activity and to the production of delphinidin-based pigments.

# EXAMPLE 6

## *Evaluation of promoters in roses*

### *Development of GUS gene expression cassettes*

The evaluation of the promoter and terminator fragments was performed using the GUS reporter gene. Therefore, a number of promoters were linked to the  $\beta$ -glucuronidase reporter gene (*GUS*) (Jefferson *et al.*, 1987, *supra*) and introduced into roses in an attempt to identify expression cassettes that lead to effective initiation of transcription in rose flowers.

A summary of the promoters and terminator fragments evaluated is given in Table 7.

TABLE 7 List of chimaeric *GUS* gene expression cassettes evaluated in roses

PLASMID	CASSETTE	EXPRESSION	SELECTABLE	BACKBONE
pCGP1307	<i>petD8 5': GUS: petD8 3'</i>	<i>mas 5': nptII: mas 3'</i>	pCGN1548	pBIN19
pCGP1506	<i>long petALS 5': GUS: petALS 3'</i>	<i>nos 5': nptII: nos 3'</i>		pWTT2132
pCGP1626	<i>chrysCHS 5': GUS: petRT 3'</i>	<i>35S 5': SURB</i>	pWTT2132	pWTT2132
pCGP1641	<i>petRT 5': GUS: petRT 3'</i>	<i>35S 5': SURB</i>	pWTT2132	pWTT2132
pCGP1861	<i>RoseCHS 5': GUS: nos 3'</i>	<i>35S 5': SURB</i>	pWTT2132	pWTT2132
pCGP1953	<i>AmCHS 5': GUS: petD8 3'</i>	<i>35S 5': SURB</i>	pWTT2132	pWTT2132
pWTT2084	<i>35S 5': GUS: nos 3'</i>	<i>35S 5': SURB</i>	pWTT2132	pWTT2132

- 80 -

**The binary vector pCGP1307 (petD8 5': GUS: petD8 3')**

The plasmid pCGP1307 (Figure 18) contains a chimeric *GUS* gene under the control of a promoter and terminator fragment from the petunia *PLTP* gene (*petD8* 5' and *petD8* 3', respectively). The chimeric *GUS* reporter gene cassette is in a tandem orientation with respect to the *mas* 5': *nptII*: *mas* 3' selectable marker gene cassette of the binary vector pCGN1548 (McBride and Summerfelt, 1990, *supra*).

**Intermediates in the preparation of the binary vector pCGP1307**

The *nos* 3' fragment from pCGP1101 (see Example 4) was replaced with the 0.75 kb *petD8* 3' fragment (Holton, 1992, *supra*) to produce the plasmid pCGP1106 containing a *petD8* 5': *GUS*: *petD8* 3' expression cassette.

The 5.3 kb fragment containing the *petD8* 5': *GUS*: *petD8* 3' expression cassette was released from the plasmid pCGP1106 upon digestion with the restriction endonucleases *HindIII* and *PstI*. The fragment was purified and ligated with *HindIII*/*PstI* ends of the binary vector, pCGN1548 (McBride and Summerfelt, 1990, *supra*). Correct insertion of the fragment was established by restriction endonuclease analysis of DNA isolated from gentamycin-resistant transformants. The resulting plasmid was designated as pCGP1307 (Figure 18).

**Plant transformation with pCGP1307**

The T-DNA contained in the binary vector plasmid pCGP1307 (Figure 18) was introduced into rose via *Agrobacterium*-mediated transformation.

**The binary vector pCGP1506 (long petFLS 5': GUS: petFLS 3')**

The plasmid pCGP1506 (Figure 19) contains a chimeric *GUS* gene under the control of promoter and terminator fragments from the petunia flavonol synthase gene (*petFLS* 5' and *petFLS* 3', respectively). The chimeric *GUS* reporter gene cassette is in a tandem orientation with respect to the *nos* 5': *nptII*: *nos* 3' selectable marker gene cassette of the binary vector pBIN19 (Bevan, *Nucleic Acids Res* 12: 8711-8721, 1984).

# *Intermediates in the preparation of the binary vector pCGP1506*

A 4 kb long petunia *FLS* promoter fragment upstream from the putative translational initiation site was released from the plasmid pCGP486 (described in Example 4) upon digestion with the restriction endonucleases *Xho*I and *Pst*I. The fragment generated was purified and ligated with *Xho*I/*Pst*I ends of pBluescript II KS+ (Stratagene, USA). Correct insertion of the fragment was established by restriction endonuclease analysis of DNA isolated from ampicillin-resistant transformants. The resulting plasmid was designated as pCGP715.

**10** *Construction of pCGP494 (long petFLS 5':petFLS3' expression cassette)*  
A 4.0 kb fragment containing the long petunia *FLS* promoter fragment was amplified by PCR using the plasmid pCGP715 as template and the T3 primer (Stratagene, USA) and an *FLS*-*Neo* primer (5' AAA ATC GAT ACC ATG GTC TTT TTT TCT TTG TCT ATA C 3') (SEQ ID NO:19). The PCR product was digested with the restriction endonucleases *Xho*I and *Cla*I and the purified fragment was ligated with *Xho*I/*Cla*I ends of pCGP716 (described in Example 4). Correct insertion of the fragment was established by restriction endonuclease analysis of DNA isolated from ampicillin-resistant transformants. The resulting plasmid was designated as pCGP494.

**20** *Construction of pCGP496 (long petFLS 5':GUS:petFLS3' expression cassette)*  
The *GUS* coding sequence from the plasmid pJB1 (Bodan, 1994, *supra*) was released upon digestion with the restriction endonucleases *Nco*I and *Sma*I. The *GUS* fragment generated was purified and ligated with *Cla*I (repaired ends)/*Nco*I ends of the plasmid pCGP494. Correct insertion of the fragment was established by restriction endonuclease analysis of DNA isolated from ampicillin-resistant transformants. The resulting plasmid was designated as pCGP496.

**30** *Construction of pCGP1506 (long petFLS 5':GUS:petFLS3' binary vector)*  
The plasmid pCGP496 was firstly linearised upon digestion with the restriction endonuclease *Xho*I. The overhanging ends were partially repaired (using only dTTP and dCTP in the reparation reaction) and a 6.7 kb fragment containing the long petFLS 5':

- 82 -

*GUS: petFLS3'* gene expression cassette was released upon digestion with the restriction endonuclease *SacI*. The fragment generated was purified and ligated with *Bam*HI (partially repaired ends using dGTP and dATP in the reparation reaction)/*SacI* ends of the binary vector pBIN19. Correct insertion of the fragment in a tandem orientation with respect to the *nos 5': nptII: nos 3'* selectable marker gene cassette was established by restriction endonuclease analysis of plasmid DNA isolated from kanamycin-resistant transformants. The resulting plasmid was designated as pCGP1506 (Figure 19).

#### Plant transformation with pCGP1506

The T-DNA contained in the binary vector plasmid pCGP1506 (Figure 19) was introduced into rose via *Agrobacterium*-mediated transformation.

#### The binary vector pCGP1626 (*chrysCHS 5': GUS: petRT 3'*)

The plasmid pCGP1626 (Figure 20) contains a chimeric *GUS* gene under the control of promoter fragment from the chalcone synthase gene of chrysanthemum (*chrysCHS 5'*) and a terminator fragment from the 3RT gene of petunia (*petRT 3'*) (Brugliera, 1994, *supra*). The chimeric *GUS* reporter gene cassette is in a tandem orientation with respect to the 35S 5': *SuRB* selectable marker gene cassette of the binary vector pWTT2132 (DNAP) (Figure 6).

#### Intermediates in the preparation of the binary vector pCGP1626

##### Isolation of chrysanthemum *CHS* promoter

A chrysanthemum genomic DNA library was prepared from genomic DNA isolated from young leaf material of the chrysanthemum cv Hero.

The chrysanthemum genomic DNA library was screened with <sup>32</sup>P-labelled fragments of a chrysanthemum *CHS* cDNA clone (SEQ ID NO:28) (contained in the plasmid pCGP856) using high stringency conditions. The plasmid pCGP856 contains a 1.5 kb cDNA clone of *CHS* isolated from a petal cDNA library prepared from RNA isolated from the chrysanthemum cv. Dark Pink Pom Pom.

A genomic clone (*CHS5*) was chosen for further analysis and found to contain ~3 kb of sequence upstream of the putative initiating methionine of the chrysanthemum *CHS* coding region.

5 A 4 kb fragment was released upon digestion of the genomic clone *CHS5* with the restriction endonuclease *HindIII*. The fragment containing the chrysanthemum *CHS* promoter was purified and ligated with *HindIII* ends of pBluescript SK (Stratagene, USA). Correct insertion of the fragment was established by restriction endonuclease analysis of DNA isolated from ampicillin-resistant transformants. The resulting plasmid was designated as pCGP1316.

10 A 2.6 kb chrysanthemum *CHS* promoter fragment upstream from the putative translational initiation site was amplified by PCR using pCGP1316 as template and primers "chrysanthCHSATG" (5'-GTTAAGGAAGCCATGGGT-3') (SEQ ID NO:8) and the M13 reverse primer (Stratagene, USA). Primer "chrysanthCHSATG" incorporated an *NcoI* restriction endonuclease recognition sequence at the putative translation initiation point for ease of cloning. The PCR fragment was purified and ligated with *KcoRV* (dT-tailed) ends of pBluescript KS (Holton and Graham, *Nuc. Acids Res.* 19: 1156, 1990). Correct insertion of the fragment was established by restriction endonuclease analysis of DNA isolated from ampicillin-resistant transformants. The resulting plasmid was designated as pCGP1620. The nucleotide sequence of the chrysanthemum *CHS* promoter fragment contained in pCGP1620 is represented as SEQ ID NO:30.

Construction of pCGP1622 (chrysCHS 5': GUS: nos 3' in pUC backbone)

25 A ~2.5 kb fragment containing the chrysanthemum *CHS* promoter was released from the plasmid pCGP1620 upon digestion with the restriction endonucleases *NcoI* and *PstI*. The fragment was purified and ligated with a 4.8 kb *NcoI/PstI* fragment of pJBI (Bodeau, 1994, *supra*) containing the backbone vector with the *GUS* and nos 3' fragments. Correct insertion of the fragment was established by restriction endonuclease analysis of DNA isolated from ampicillin-resistant transformants. The resulting plasmid was designated as pCGP1622.



**Construction of pCGP1626 (*chrysCHS* 5': *GUS*: *nos* 3' in binary vector)**

A ~4.6 kb fragment containing the *chrysCHS* 5': *GUS*: *nos* 3' cassette was released from the plasmid pCGP1622 upon digestion with the restriction endonucleases *Pst*I and *Bgl*II.

- 5 The fragment was purified and ligated with *Pst*I/*Bam*HI ends of the binary vector pWTT2132 (DNAP) (Figure 6). Correct insertion of the cassette in a tandem orientation with respect to the 35S 5': *SuRB* selectable marker gene cassette was established by restriction endonuclease analysis of DNA isolated from tetracycline-resistant transformants. The resulting plasmid was designated as pCGP1626 (Figure 20).

10

**Plant transformation with pCGP1626**

The T-DNA contained in the binary vector plasmid pCGP1626 (Figure 20) was introduced into rose via *Agrobacterium*-mediated transformation.

- 15 **The binary vector pCGP1641 (*petRT* 5': *GUS*: *petRT* 3')**

The plasmid pCGP1641 (Figure 21) contains a chimeric *GUS* gene under the control of a petunia 3RT promoter (*petRT* 5'), covering 1.1kb upstream from the putative 3RT translation initiation codon with a petunia 3RT terminator (*petRT* 3') covering 2.5 kb downstream from the 3RT stop codon. The chimeric *GUS* cassette is in a tandem orientation with respect to the 35S 5': *SuRB* selectable marker gene cassette of the binary vector, pWTT2132 (DNAP) (Figure 6).

20

**Intermediates in the preparation of the binary vector pCGP1641**

**Isolation of petunia 3RT gene**

- 25 The isolation of the petunia 3RT gene corresponding to the *Rt* locus of *P. hybrida* has been described in Brugliera, 1994, *supra*.

Construction of pCGP1625 (CAMV 35S: GUS: pRT 3' cassette)

The intermediate plasmid pCGP1625 contains a CAMV 35S: GUS: pRT 3' cassette in a pUC backbone. The 2.5 kb fragment containing a pRT terminator sequences was released from the plasmid pCGP1610 (described in Brugliera, 1994, *supra*) upon digestion

5 with the restriction endonucleases *Bam*HI and *Sac*I. The fragment was purified and ligated with the *Bgl*III/*Sac*I 4.9kb fragment of pJB1 (Bodeau, 1994, *supra*) containing the vector backbone and the CAMV 35S promoter and GUS fragments. Correct insertion of the petunia 5RT terminator fragment downstream of the GUS fragment was established by restriction endonuclease analysis of DNA isolated from ampicillin-resistant transformants. The resulting plasmid was designated as pCGP1625.

10

Construction of pCGP1628 (pRT 5': GUS: pRT 3' cassette)

A 1.1 kb pRT promoter fragment was released from the plasmid pCGP1611 (described in Brugliera, 1994, *supra*) upon digestion with the restriction endonucleases *Nco*I and *Pst*I. The purified fragment was ligated with *Nco*I/*Pst*I ends of the 7kb fragment of pCGP1625 containing the vector backbone and the GUS and pRT 3' fragments. Correct insertion of the pRT promoter fragment upstream of the GUS fragment was established by restriction endonuclease analysis of DNA isolated from ampicillin-resistant transformants. The resulting plasmid was designated as pCGP1628.

15

Construction of pCGP1641 (pRT 5': GUS: pRT 3' binary vector)

A 5.4 kb fragment containing the pRT 5': GUS: pRT 3' cassette was released from pCGP1628 upon digestion with the restriction endonuclease *Pst*I. The fragment was purified and ligated with *Pst*I ends of the binary vector pWTT2132 (DNAP) (Figure 6). Correct insertion of the fragment in a tandem orientation with respect to the 35S 5': *SuRB* selectable marker gene cassette was established by restriction endonuclease analysis of plasmid DNA isolated from tetracycline-resistant transformants. The resulting plasmid was designated as pCGP1641 (Figure 21).

25

20

- 86 -

**Plant transformation with pCGP1641**

The T-DNA contained in the binary vector plasmid pCGP1641 (Figure 21) was introduced into rose via *Agrobacterium*-mediated transformation.

5     **The binary vector pCGP1861 (RoseCHS 5': GUS: nos 3')**

The plasmid pCGP1861 (Figure 22) contains a chimeric *GUS* gene under the control of a promoter fragment from the *CHS* gene of *R. hybrida* (*RoseCHS 5'*) with a terminator fragment from the *nos* gene of *Agrobacterium* (*nos 3'*). The chimeric *GUS* reporter gene cassette is in a tandem orientation with respect to the 35S 5': *SuRB* selectable marker gene cassette of the binary vector, pWTT2132 (Figure 6).

An ~5 kb fragment containing the *RoseCHS 5': GUS: nos 3'* cassette was released from pCGP197 (described in Example 4) upon digestion with the restriction endonuclease *Bgl*II. The fragment was purified and ligated with *Bam*HI ends of the binary vector, pWTT2132 (DNAP). Correct insertion of the fragment in a tandem orientation with respect to the 35S 5': *SuRB* selectable marker gene cassette was established by restriction endonuclease analysis of plasmid DNA isolated from tetracycline-resistant transformants. The resulting plasmid was designated as pCGP1861 (Figure 22).

20     **Plant transformation with pCGP1861**

The T-DNA contained in the binary vector plasmid pCGP1861 (Figure 22) was introduced into rose via *Agrobacterium*-mediated transformation.

**The binary vector pCGP1953 (AmCHS 5': GUS: petD8 3')**

25     The plasmid pCGP1953 (Figure 23) contains a chimeric *GUS* gene under the control of a promoter fragment from the *CHS* gene of *Antirrhinum majus* (*AmCHS 5'*) with a petunia *PLTP* terminator (*petD8 3'*). The chimeric *GUS* reporter gene cassette is in a tandem orientation with respect to the 35S 5': *SuRB* selectable marker gene cassette of the binary vector, pWTT2132 (DNAP) (Figure 6).

30

# *Intermediates in the preparation of the binary vector pCGP1953*

The plasmid pJB1 (Bodeau, 1994, *supra*) was linearised with the restriction endonuclease *NcoI*. The overhanging ends were repaired and the 1.8 kb *GUS* fragment was released upon digestion with *BamHI*. The *GUS* fragment was purified and was ligated with the 5 kb *XbaI*(ends repaired)/*BamHI* fragment of pCGP726 containing the pBluescript backbone vector and the *AmCHS* 5' and *peid8* 3' fragments (described in Example 4). Correct insertion of the *GUS* fragment between the *AmCHS* 5' and *peid8* 3' fragments was established by restriction endonuclease analysis of plasmid DNA isolated from ampicillin-resistant transformants. The plasmid was designated as pCGP1952.

A 3.8 kb fragment containing the *AmCHS* 5' *GUS*: *peid8* 3' expression cassette was released from the plasmid pCGP1952 upon digestion with the restriction endonucleases *EagI* and *PstI*. The overhanging ends were repaired and the purified fragment was ligated with the repaired ends of an *Asp718* digested pWTT2132 binary vector (Figure 6). Correct insertion of the fragment in a tandem orientation with respect to the 35S 5' *SwRB* selectable marker gene cassette was established by restriction endonuclease analysis of plasmid DNA isolated from tetracycline-resistant transformants. The plasmid was designated as pCGP1953 (Figure 23).

## *Plant transformation with pCGP1953*

The T-DNA contained in the binary vector plasmid pCGP1953 (Figure 23) was introduced into rose via *Agrobacterium*-mediated transformation.

## *The binary vector pWTT2084 (35S 5' *GUS*: *ocs* 3')*

The plasmid pWTT2084 (DNAP) (Figure 24) contains a chimeric *GUS* gene under the control of a *CaMV* 35S promoter (35S 5') with an octopine synthase terminator (*ocs* 3'). An ~60 bp 5' untranslated leader sequence from the petunia chlorophyll *a/b* binding protein gene (*Cab* 22 gene) (Harpster *et al.*, 1988, *supra*) is included between the *CaMV* 35S promoter fragment and the *GUS* clone. The chimeric *GUS* cassette is in a tandem orientation with respect to the 35S 5' *SwRB* selectable marker gene cassette of the binary vector, pWTT2084.

- 88 -

**Plant transformation with pWTT2084**

The T-DNA contained in the binary vector plasmid pWTT2084 (Figure 24) was introduced into rose via *Agrobacterium*-mediated transformation.

5

Transgenic analysis of roses transformed with GUS expression cassettes

Northern blot analysis was performed on total RNA isolated from petals of developmental stages 3 to 4 of transgenic Kardinal roses transformed with the T-DNA of various GUS expression cassettes. There was either no accumulating transcript or an intact full-length transcript of the expected size of ~1.8kb as detected by Northern blot hybridisation. The relative levels of GUS transcripts accumulating in the rose petals were recorded (see Table 8).

10

**TABLE 8** Summary of Northern analysis on transgenic Kardinal rose flowers (open bud stage) containing GUS constructs.

15

PLASMID	GUS REPORTER GENE	SELECTABLE MARKER GENE	GUS TRANSCRIPT LEVELS
pCGP1307	<i>petD8</i> 5': GUS: <i>petD8</i> 3'	<i>mas</i> 5': <i>nptII</i> : <i>mas</i> 3'	—
pCGP1506	<i>petFLS</i> 5': GUS: <i>petFLS</i> 3'	<i>nos</i> 5': <i>nptII</i> : <i>nos</i> 3'	—
pCGP1626	<i>chrysCHS</i> 5': GUS: <i>petRT</i> 3'	<i>35S</i> 5': <i>SuRB</i>	++ to +++
pCGP1641	<i>petRT</i> 5': GUS: <i>petRT</i> 3'	<i>35S</i> 5': <i>SuRB</i>	—
pCGP1861	<i>RoseCHS</i> 5': GUS: <i>nos</i> 3'	<i>35S</i> 5': <i>SuRB</i>	++++
pCGP1953	<i>AmCHS</i> 5': GUS: <i>petD8</i> 3'	<i>35S</i> 5': <i>SuRB</i>	—
pWTT2084	<i>35S</i> 5': GUS: <i>ocs</i> 3'	<i>35S</i> 5': <i>SuRB</i>	+++++

— = no transcripts detected

+ to +++++ = relative levels (low to high) of full-length GUS transcript detected by Northern blot analysis

20

Based on the above results (Table 8), the *CaMV 35S (35S 5)* and rose *CHS (RoseCHS 5)* promoters appear to drive relatively high levels of transcription in rose petals. The chrysanthemum *CHS* promoter (*chysCHS 5*) appears to also lead to high transcript levels but not as high as those obtained using *CaMV 35S* or rose *CHS* promoters. Surprisingly, anthrimum (snapdragon) *CHS (AmCHS 5)*, petunia *3RT (peIRT 5)*, petunia *FLS (petFLS 5)* and petunia *PLTP-(peID8 5)* promoters did not appear to function in rose petals as no *GUS* transcripts were detected with expression cassettes incorporating these promoters. However, these same promoters fused to *peHf1* and/or *-GUS* genes had previously been proven to function well in carnation and petunia leading to relatively high full-length transcript levels and for *peHf1* genes, the production of delphinidin or delphinidin-based molecules pigments. The result obtained with the anthrimum *CHS* promoter (*AmCHS 5*) fused with the *GUS* gene was more surprising as promoter regions from homologous genes from two other species (rose and chrysanthemum) appeared to function relatively well in roses. The anthrimum *CHS* promoter had also been successfully used in conjunction with petunia *F3'5'H (peHf1)* to produce the novel violet-colored carnations Florigene Moondust (see International Patent Application No. PCT/AU96/00296).

The evaluation of promoter and terminator fragments fused with the *GUS* gene also provided further evidence to suggest that the petunia *F3'5'H peHf1* and *peHf2* sequences were unstable in roses as constructs containing the petunia *F3'5'H* sequences ligated to the *CaMV 35S*, -rose *CHS* and chrysanthemum *CHS* promoters (which do function in rose) did not result in intact petunia *F3'5'H peHf1* or *peHf2* transcripts in roses (see Table 6).

#### EXAMPLE 7

##### Isolation of *F3'5'H* sequences from species other than petunia

Since the petunia *F3'5'H* sequences had already been proven to function in various plants such as carnation, petunia and tobacco and ultimately resulted in the production of delphinidin-based pigments, it was reasonable to assume that these sequences would also prove functional in roses. There was an assumption that the enzyme activity may vary depending on the background of the species, indeed between cultivars of a given species, that the petunia *F3'5'H* was introduced into. However, there was no expectation that full-

- 90 -

length recombinant petunia *F3'5'H* mRNA would not accumulate. Analysis of the petunia *F3'5'H* nucleotide sequences (*petHf1* and *petHf2*) did not reveal any sequences which might lead to instability and subsequent degradation (Johnson *et al.*, *In A look beyond transcription*, ASPP, USA, Bailey-Serres and Gallie, eds, 1998), intron: exon splice  
5 junctions (Brendel *et al.*, *In A look beyond transcription*, ASPP, USA, Bailey-Serres and Gallie, eds, 1998), or any autocatalytic or degradation trigger sequences reported in the scientific literature to date (*In A look beyond transcription*, ASPP, USA, Bailey-Serres and Gallie, eds, 1998). The surprising result suggested that there were factors specific to rose that resulted in petunia *F3'5'H* sequences being unstable.

10

Since it was not obvious why the petunia *F3'5'H* sequences were unstable in roses but stable in carnation, petunia or tobacco, a number of *F3'5'H* sequences were isolated across a range of families in an attempt to determine whether any *F3'5'H* sequence would be stable in rose and then identify any *F3'5'H* sequences that would lead to the synthesis of  
15 stable *F3'5'H* transcripts and *F3'5'H* activity and ultimately the production of delphinidin-based pigments in roses leading to a change in flower color.

#### Construction of petal cDNA libraries

Petal cDNA libraries were prepared from RNA isolated from petals from bud to opened  
20 flower stages from various species of plants described in Table 9. *Rosa hybrida* is classified in the family Rosaceae, Order Rosales, Subclass Rosidae and so species that produced delphinidin-based pigments and so contained a functional *F3'5'H* and belonged to the Subclass Rosidae were selected. *Petunia hybrida* is classified in the Family Solanaceae, Order Solanales, Subclass Asteridae and so species from the Subclass  
25 Asteridae that produced delphinidin-based pigments were also selected.

**TABLE 9** List of flowers from which total RNA was isolated for the preparation of petal cDNA libraries. Information obtained from National Center for Biotechnology Information (NCBI) website under Taxonomy browser (TaxBrowser) as of August 2003.

FLOWER	SPECIES	FAMILY	ORDER	SUBCLASS
gentian	<i>Gentiana</i> spp.	Gentianaceae	Gentianales	Asteridae
lavender	<i>Lawandula</i> spp.	Lamiaceae	Lamiales	Asteridae
salvia	<i>Salvia</i> spp.	Lamiaceae	Lamiales	Asteridae
sollya	<i>Sollya</i> spp.	Pinosporaceae	Apiales	Asteridae
kennedia	<i>Kennedia</i> spp.	Fabaceae	Fabales	Rosidae
butterfly pea	<i>Clitonia ternata</i>	Fabaceae	Fabales	Rosidae
pansy	<i>Viola</i> spp.	Violaceae	Malpighiales	Rosidae

Unless otherwise described, total RNA was isolated from the petal tissue of purple/blue flowers using the method of Turpen and Griffith (*BioTechniques* 4: 11-15, 1986). Poly(A)<sup>+</sup> RNA was selected from the total RNA, using oligotex-dT<sup>TM</sup> (Qiagen) or by three cycles of oligo-dT cellulose chromatography (Aviv and Leder, *Proc. Natl. Acad. Sci. USA* 69: 1408, 1972).

In general a λZAP/IV Gigapack II Cloning kit (Stratagene, USA) (Short *et al.*, *Nucl. Acids Res.* 16: 7583-7600, 1988) was used to construct directional petal cDNA libraries in λZAPII using around 5 µg of poly(A)<sup>+</sup> RNA isolated from petal as template. The total number of recombinants obtained was generally in the order of  $1 \times 10^5$  to  $1 \times 10^6$ .

After transfecting XL1-Blue MRF' cells, the packaged cDNA mixtures were plated at around 50,000 pfu per 15 cm diameter plate. The plates were incubated at 37°C for 8 hours, and the phage were eluted in 100 mM NaCl, 8 mM MgSO<sub>4</sub>, 50 mM Tris-HCl pH 8.0, 0.01% (w/v) gelatin (Phage Storage Buffer (PSB)) (Sambrook *et al.*, 1989, *supra*). Chloroform was added and the phages stored at 4°C as amplified libraries.



- 92 -

In general around 100,000 pfu of the amplified libraries were plated onto NZY plates (Sambrook *et al.*, 1989, *supra*) at a density of around 10,000 pfu per 15 cm plate after transfecting XL1-Blue MRF' cells, and incubated at 37°C for 8 hours. After incubation at 4°C overnight, duplicate lifts were taken onto Colony/Plaque Screen™ filters (DuPont) and  
5 treated as recommended by the manufacturer.

#### Plasmid Isolation

Helper phage R408 (Stratagene, USA) was used to excise pBluescript phagemids containing cDNA inserts from amplified λZAPII or λZAP cDNA libraries using methods  
10 described by the manufacturer.

#### Screening of petal cDNA Libraries

Prior to hybridization, duplicate plaque lifts were washed in prewashing solution (50 mM Tris-HCl pH7.5, 1 M NaCl, 1 mM EDTA, 0.1% (w/v) sarcosine) at 65°C for 30 minutes;  
15 followed by washing in 0.4 M sodium hydroxide at 65°C for 30 minutes; then washed in a solution of 0.2 M Tris-HCl pH 8.0, 0.1 x SSC, 0.1% (w/v) SDS at 65°C for 30 minutes and finally rinsed in 2 x SSC, 1.0% (w/v) SDS.

The membrane lifts from the petal cDNA libraries were hybridized with <sup>32</sup>P-labelled  
20 fragments of a 1.6 kb *Bsp*HI/*Fsp*I fragment from pCGP602 (Figure 2) (SEQ ID NO: 1) containing the petunia *F3'5'H petHf1* cDNA clone (Holton *et al.*, 1993a, *supra*).

Hybridization conditions included a prehybridization step in 10% v/v formamide, 1 M NaCl, 10% w/v dextran sulphate, 1% w/v SDS at 42°C for at least 1 hour. The <sup>32</sup>P-labelled  
25 fragments (each at 1x10<sup>6</sup> cpm/mL) were then added to the hybridization solution and hybridization was continued at 42°C for a further 16 hours. The filters were then washed in 2 x SSC, 1% w/v SDS at 42°C for 2 x 1 hour and exposed to Kodak XAR film with an intensifying screen at -70°C for 16 hours.

Strongly hybridizing plaques were picked into PSB (Sambrook *et al.*, 1989, *supra*) and rescreened to isolate purified plaques, using the plating and hybridization conditions as described for the initial screening of the cDNA library. The plasmids contained in the  $\lambda$ ZAP or  $\lambda$ ZAP bacteriophage vectors were rescued and sequence data was generated from the 3' and 5' ends of the cDNA inserts. New F3'5'H cDNA clones were identified based on sequence similarity to the petunia F3'5'H *peHf1* cDNA clone.

The cDNA clones isolated were given plasmid designation numbers as described in Table 10.

TABLE 10 Plasmid numbers and SEQ ID NO. of F3'5'H cDNA clones isolated from various species

SPECIES	CLONE	PLASMID NUMBER	FIGURE NUMBER	SEQ ID NO.
<i>Viola spp.</i>	BP#18	pCGP1959	25	9
<i>Viola spp.</i>	BP#40	pCGP1961	26	11
<i>Salvia spp.</i>	Sal#2	pCGP1995	31	13
<i>Salvia spp.</i>	Sal#47	pCGP1999	32	15
<i>Soliva spp.</i>	Soil#5	pCGP2110	37	17
<i>Kennedia spp.</i>	Kenn#31	pCGP2231	40	26
<i>Clinoria ternatea</i>	BpeaHR2	pBHR2P4	43	20
<i>Gentiana triflora</i>	Gent#48	pG48	47	22
<i>Lavandula nil</i>	LBG	PLHF8	51	31

15 *Viola* (pansy) F3'5'H constructs

Isolation of F3'5'H cDNA clones from petals of *Viola* spp. (pansy)

Total RNA and poly (A)<sup>+</sup> RNA was isolated from petals of young buds of *Viola* spp. cultivar black pansy as described above. A petal cDNA library was constructed using  $\lambda$ ZAPII/Gigapack II Cloning kit (Stratagene, USA) and screened as described above. Two full-length pansy F3'5'H cDNA clones (BP#18 (SEQ ID NO:9) in pCGP1959 (Figure 25)

- 94 -

and BP#40 (SEQ ID NO:11) in pCGP1961 (Figure 26)) were identified by sequence similarity to the petunia *F3'5'H petHf1* cDNA clone (SEQ ID NO:1). The BP#18 and BP#40 shared 82% identity at the nucleotide level. Comparison of the nucleotide sequence of pansy *F3'5'H* clones (BP#18 and BP#40) with that of the petunia *F3'5'H* revealed  
 5 around 60% identity to the petunia *F3'5'H petHf1* clone and 62% identity to the petunia *F3'5'H petHf2* clone.

**The binary vectors, pCGP1972 and pCGP1973 (*AmCHS* 5': BP#18 or BP#40: *petD8* 3')**

The plasmids pCGP1972 (Figure 27) and pCGP1973 (Figure 28) contain the pansy *F3'5'H* cDNA clone (BP#18 and BP#40, respectively) between an *A. majus* (snapdragon) *CHS* promoter fragment (*AmCHS* 5') and a petunia *PLTP* terminator fragment (*petD8* 3'). The chimeric *F3'5'H* genes are in tandem with respect to the 35S 5': *SuRB* selectable marker gene cassette of the binary vector, pWTT2132 (DNAP) (Figure 6).  
 10

15 The petunia *F3'5'H (petHf1)* cDNA clone in pCGP725 (described in Example 4) (Figure 7) was removed by initially digesting pCGP725 with the restriction endonuclease *Bam*HI. The ends were repaired and the linearised plasmid was further digested with the restriction endonuclease *Xba*I. The ~4.9kb fragment containing the vector with the *AmCHS* 5' and *petD8* 3' fragments was purified and ligated with the ~1.6kb *Kpn*I (ends repaired)/*Xba*I  
 20 fragment containing the pansy *F3'5'H* BP#18 or BP#40 cDNA clone from pCGP1959 or pCGP1961, respectively to produce pCGP1970 and pCGP1971, respectively. The *AmCHS* 5': pansy *F3'5'H*: *petD8* 3' cassette was then isolated from pCGP1970 or pCGP1971 by firstly digesting with the restriction endonuclease *Not*I. The ends of the linearised plasmid were repaired and then the chimeric *F3'5'H* genes were released upon digestion with the  
 25 restriction endonuclease *Eco*RV. The purified fragments were then ligated with *Asp*718 (repaired ends) of the binary vector pWTT2132 (DNAP). Correct insertion of the fragment was established by restriction endonuclease analysis of plasmid DNA isolated from tetracycline-resistant transformants. The resulting plasmids were designated pCGP1972 (Figure 27) and pCGP1973 (Figure 28), respectively.

30

Coronation and petunia transformation with pCGP1972 and 1973

The T-DNAs contained in the binary vector plasmids pCGP1972 (Figure 27) and pCGP1973 (Figure 28) were introduced separately into *Dianthus caryophyllus* cultivars Kordina Chanel and Monte Lisa and *Petunia hybrida* cv. Skry4 x Sw63 via *Agrobacterium*-mediated transformation.

The binary vectors, pCGP1967 and pCGP1969 (CaMV 35S; pansy F3'5'H; ocs 3')

The binary vectors pCGP1967 (Figure 29) and pCGP1969 (Figure 30) contain chimeric *CaMV 35S: pansy F3'5'H: ocs 3'* genes in tandem with respect to the 35S 5': *SuRB* selectable marker gene cassette of the binary vector, pWTT2132 (DNAP) (Figure 6).

Intermediates in the preparation of the binary vectors pCGP1967 and pCGP1969

The plasmids pCGP1959 (Figure 25) and pCGP1961 (Figure 26) were firstly linearized upon digestion with the restriction endonuclease *KpnI*. The overhanging *KpnI* ends were repaired and the pansy *F3'5'H* cDNA clones, BP#18 and BP#40, were released upon digestion with the restriction endonuclease *PstI*. The ~1.6 kb fragments generated were ligated with an ~5.9 kb *EcoRI* (repaired ends)/*PstI* fragment of pK1W1101 (Janssen and Gardner, 1989, *supra*). Correct insertion of each fragment was established by restriction endonuclease analysis of plasmid DNA isolated from ampicillin-resistant transformants. The resulting plasmids were designated pCGP1965 and pCGP1966, respectively.

The plasmids pCGP1965 and pCGP1966 were firstly partially digested with the restriction endonuclease *XhoI*. The resulting fragments were further digested with the restriction endonuclease *XbaI*. The overhanging ends were repaired and the 3.6kb fragments containing the *CaMV 35S: pansy F3'5'H: ocs 3'* chimeric genes were isolated and ligated with *Asp18* repaired ends of pWTT2132 (Figure 6). Correct insertion of each fragment was established by restriction endonuclease analysis of plasmid DNA isolated from tetracycline-resistant transformants. The resulting plasmids were designated pCGP1967 (Figure 29) and pCGP1969 (Figure 30), respectively.

- 96 -

**Rose transformation with pCGP1967 and pCGP1969**

The T-DNAs contained in the binary vector plasmids pCGP1967 (Figure 29) and pCGP1969 (Figure 31) were introduced separately into *Rosa hybrida* cv. Kardinal and Soft Promise via *Agrobacterium*-mediated transformation. The T-DNA contained in the binary  
 5 vector plasmids pCGP1969 (Figure 31) was also introduced into *Rosa hybrida* cv. Pamela and Medeo via *Agrobacterium*-mediated transformation.

**Salvia F3'5'H constructs****Isolation of a F3'5'H cDNA clone from petals of Salvia spp.**

10 Total RNA and poly (A)<sup>+</sup> RNA was isolated from young petal buds of *Salvia* spp. (bought from a nursery) as described above. A petal cDNA library was constructed using λZAPIII/Gigapack II Cloning kit (Stratagene, USA). Two full-length salvia F3'5'H cDNA clones (Sal#2 (SEQ ID NO:13) in pCGP1995 (Figure 31) and Sal#47 (SEQ ID NO:15) in pCGP1999 (Figure 32)) were identified by sequence similarity with the petunia F3'5'H  
 15 *petHf1* cDNA clone. The Sal#2 and Sal#47 shared 95% identity at the nucleotide level. Comparison of the nucleotide sequence of salvia F3'5'H clones (Sal#2 and Sal#47) with that of the petunia F3'5'H revealed around 57% identity to the petunia F3'5'H *petHf1* clone (SEQ ID NO:1) and 58% identity to the petunia F3'5'H *petHf2* clone (SEQ ID NO:3).

20

The binary vectors, pCGP2121 and pCGP2122

(*AmCHS* 5': *Salvia* F3'5'H #2 or #47: *petD8* 3')

The plasmids pCGP2121 (Figure 33) and pCGP2122 (Figure 34) contain the salvia F3'5'H cDNA clones (Sal#2 and Sal#47, respectively) between a snapdragon *CHS* promoter  
 25 fragment (*AmCHS* 5') and a petunia *PLTP* terminator fragment (*petD8* 3') in tandem with the 35S 5': *SuRB* selectable marker gene cassette of the binary vector pWTT2132 (DNAP) (Figure 6).

The petunia *F3'5'H* (*peH7*) cDNA clone in pCGP725 (described in Example 4) (Figure 7) was removed by initially digesting pCGP725 with the restriction endonuclease *Bam*HI. The ends were repaired and the linearised plasmid was further digested with the restriction endonuclease *Xba*I. The ~4.9kb fragment containing the vector with the *Am*CHS 5' and *petD8* 3' fragments was purified and ligated with the ~1.6kb *Xho*I/*Bam*HI (ends repaired) fragment from pCGP1995 (Figure 31) containing the salvia *F3'5'H* #2 or *Xho*I/*Eco*RI (ends repaired) fragment from pCGP1999 (Figure 32) containing the salvia *F3'5'H* #47, respectively to produce pCGP2116 and pCGP2117, respectively.

10 The *Am*CHS 5': salvia *F3'5'H*: *petD8* 3' cassette was isolated from pCGP2116 or pCGP2117 by firstly digesting with the restriction endonuclease *Nco*I. The ends of the linearized plasmid were repaired and then the chimeric *F3'5'H* gene cassettes were released upon digestion with the restriction endonuclease *Eco*RV. The ~3.6kb purified fragments were then ligated with *Asp*718 repaired ends of the binary vector pCGP1988 (Figure 16) (described in Example 4). Correct insertion of each fragment was established by restriction endonuclease analysis of plasmid DNA isolated from tetraacycline-resistant transformants. The resulting plasmids were designated pCGP2121 (Figure 33) and pCGP2122 (Figure 34), respectively.

#### 20 Carnation and petunia transformation with pCGP2121 and pCGP2122

The T-DNAs contained in the binary vector plasmids pCGP2121 (Figure 33) and pCGP2122 (Figure 34) were introduced separately into *Dianthus caryophyllus* cultivars Korina Chapel and Monte Lisa and *Petunia hybrida* cv. Sk4 x Sw63 via *Agrobacterium*-mediated transformation.

The binary vectors, pCGP2120 and pCGP2119 (CamV 35S: salvia *F3'5'H*: *ocs* 3')  
The binary vectors pCGP2120 (Figure 35) and pCGP2119 (Figure 36) contain chimeric CamV 35S: salvia *F3'5'H*: *ocs* 3' gene cassettes in tandem with the 35S 5': *SURB* selectable marker gene cassette of the binary vector pCGP1988 (Figure 16).

- 98 -

***Intermediates in the preparation of the binary vectors pCGP2120 and pCGP2119***

The plasmids pCGP1995 (Figure 31) and pCGP1999 (Figure 32) were firstly linearized upon digestion with the restriction endonuclease *Xho*I. The overhanging *Xho*I ends were repaired and then the *salvia F3'5'H* cDNA clones Sal#2 or Sal#47 were released upon  
5 digestion with the restriction endonuclease *Eco*RI. In the case of pCGP1995 a partial digest with *Eco*RI was undertaken. The ~1.7 kb fragments were ligated with the *Cla*I (repaired ends)/*Eco*RI ends of pCGP2105 (Figure 17). Correct insertion of each fragment was established by restriction endonuclease analysis of plasmid DNA isolated from ampicillin-resistant transformants. The resulting plasmids were designated pCGP2112 and  
10 pCGP2111, respectively.

The plasmids pCGP2112 and pCGP2111 were digested with the restriction endonucleases *Xho*I and *Xba*I. The resulting overhanging ends were repaired and ~3.6 kb fragments containing the *CaMV 35S: salvia F3'5'H: ocs 3'* chimeric genes were isolated and ligated  
15 with *Asp*718 repaired ends of the binary vector, pCGP1988 (described in Example 4). Correct insertion of each fragment was established by restriction endonuclease analysis of plasmid DNA isolated from tetracycline-resistant transformants. The resulting plasmids were designated pCGP2120 (Figure 35) and pCGP2119 (Figure 36), respectively.

***Rose transformation with pCGP2120 and pCGP2119***

The T-DNAs contained in the binary vector plasmids pCGP2120 (Figure 35) and pCGP2119 (Figure 36) were introduced separately into *Rosa hybrida* cv. Kardinal via *Agrobacterium*-mediated transformation.

***Solhya F3'5'H constructs******Isolation of a F3'5'H cDNA clone from petals of Solhya spp.***

Total RNA and poly (A)<sup>+</sup> RNA was isolated from young petal buds of *Solhya* spp. (bought from a nursery) as described above. A petal cDNA library was constructed using λZAPII/Gigapack II Cloning kit (Stratagene, USA). One full-length *Solhya F3'5'H* cDNA clone  
30 (Sol#5 (SEQ ID NO:17) in pCGP2110 (Figure 37)) was identified by sequence similarity to the petunia *F3'5'H petHf1* cDNA clone. Comparison of the nucleotide sequence of the

soflya *F3'5'H* clone with that of the petunia *F3'5'H* revealed around 48% identity to the petunia *F3'5'H pelH1* clone (SEQ ID NO:1) and 52% identity to the petunia *F3'5'H pelH2* clone (SEQ ID NO:3).

The binary vector pCGP2130 (*AmCHS 5': soflya F3'5'H: pelD8 3'*)

The plasmid pCGP2130 (Figure 38) contains the soflya *F3'5'H SolH#5* cDNA clone between a snapdragon *CHS* promoter fragment (*AmCHS 5'*) and a petunia *PLTP* terminator fragment (*pelD8 3'*) in a tandem orientation with respect to the 35S 5': *SuRB* selectable marker gene cassette of the binary vector pCGP1988 (Figure 16).

The petunia *F3'5'H (pelH1)* cDNA clone in pCGP725 (described in Example 4) (Figure 7) was removed by initially digesting pCGP725 with the restriction endonucleases *XbaI* and *BamHI*. The ends were repaired the ~4.9kb fragment containing the vector with the *AmCHS 5'* and *pelD8 3'* fragments was purified and ligated with the repaired ends of the ~1.6kb *Asp718/PstI* fragment from pCGP2110 containing the soflya *F3'5'H* cDNA clone to produce pCGP2128. Correct insertion of the soflya *F3'5'H* fragment in tandem with the *AmCHS 5'* and *pelD8 3'* fragments was confirmed by restriction endonuclease mapping.

The *AmCHS 5': soflya F3'5'H: pelD8 3'* gene cassette was then isolated from pCGP2128 by firstly digesting with the restriction endonuclease *NotI*. The ends of the linearized plasmid were repaired and then the chimeric *F3'5'H* gene was released upon digestion with the restriction endonuclease *RcoRV*. The ~3.5kb purified fragment was then ligated with *Asp718* (repaired ends) of the binary vector pCGP1988 (described in Example 4) (Figure 16). Correct insertion of the fragment was established by restriction endonuclease analysis of plasmid DNA isolated from tetracycline-resistant transformants. The resulting plasmid was designated pCGP2130 (Figure 38).

Correlation and petunia transformation with pCGP2130

The T-DNA contained in the binary vector plasmid pCGP2130 (Figure 38) was introduced into *Dianthus caryophyllus* cultivars Kortinga Chanel and Monte Lisa and *Petunia hybrida* cv. Skirt x Sw63 via *Agrobacterium*-mediated transformation.



- 100 -

**The binary vector pCGP2131 (CaMV 35S: sollya F3'5'H: ocs 3')**

The binary vector pCGP2131 (Figure 39) contains a chimeric *CaMV 35S: sollya F3'5'H: ocs 3'* gene in tandem with the *35S 5': SuRB* selectable marker gene cassette of the binary  
5 vector pCGP1988 (Figure 16).

**Intermediates in the preparation of the binary vector pCGP2131**

The plasmid pCGP2110 was firstly linearized upon digestion with the restriction endonuclease *Asp718*. The overhanging ends were repaired and then the *sollya F3'5'H*  
10 cDNA clone was released upon digestion with the restriction endonuclease *PstI*. The ~1.7 kb fragment was ligated with the *EcoRV/PstI* ends of pCGP2105 (Figure 17). Correct insertion of the fragment was established by restriction endonuclease analysis of plasmid DNA isolated from ampicillin-resistant transformants. The resulting plasmid was designated pCGP2129.

15 A 3.6 kb fragment containing the *CaMV 35S: sollya F3'5'H: ocs 3'* chimeric gene was released upon digestion with the restriction endonucleases *Asp718* and *XbaI*. The overhanging ends were repaired and the purified fragment was ligated with of *Asp718* repaired ends of the binary vector, pCGP1988 (Figure 16). Correct insertion of the  
20 fragment was established by restriction endonuclease analysis of plasmid DNA isolated from tetracycline-resistant transformants. The resulting plasmid was designated pCGP2131 (Figure 39).

**Rose transformation with pCGP2131**

25 The T-DNA contained in the binary vector plasmid pCGP2131 (Figure 39) was introduced into *Rosa hybrida* cv. *Kardinal* via *Agrobacterium*-mediated transformation.



- 102 -

tetracycline-resistant transformants. The resulting plasmid was designated pCGP2256 (Figure 41).

*Petunia transformation with pCGP2256*

- 5 The T-DNA contained in the binary vector plasmid pCGP2256 (Figure 41) was introduced into *Petunia hybrida* cv. Skr4 x Sw63 via *Agrobacterium*-mediated transformation.

*The binary vector pCGP2252 (CaMV 35S: kennedia F3'5'H: ocs 3')*

- 10 The binary vector pCGP2252 (Figure 42) contains a chimeric *CaMV 35S: kennedia F3'5'H: ocs 3'* gene in tandem with the *35S 5': SuRB* selectable marker cassette of the binary vector pCGP1988 (Figure 16).

*Intermediates in the preparation of the binary vector pCGP2252*

- 15 The plasmid pCGP2231 was firstly linearized upon digestion with the restriction endonuclease *XhoI*. The overhanging ends were repaired and then the kennedia *F3'5'H* cDNA clone was released upon digestion with the restriction endonuclease *PstI*. The ~1.7 kb fragment was ligated with the *ClaI* (repaired ends)/*PstI* ends of pCGP2105 (Figure 17). Correct insertion of the fragment was established by restriction endonuclease analysis of plasmid DNA isolated from ampicillin-resistant transformants. The resulting plasmid was designated pCGP2236.

- 25 A 3.6 kb fragment containing the *CaMV 35S: kennedia F3'5'H: ocs 3'* chimeric gene cassette was released from the plasmid pCGP2236 upon digestion with the restriction endonucleases *XhoI* and *NotI*. The overhanging ends were repaired and the purified fragment was ligated with *Asp718* repaired ends of the binary vector, pCGP1988 (Figure 16). Correct insertion of the fragment was established by restriction endonuclease analysis of plasmid DNA isolated from tetracycline-resistant transformants. The resulting plasmid was designated pCGP2252 (Figure 42).

- 30 *Rose transformation with pCGP2252*

The T-DNA contained in the binary vector plasmid pCGP2252 (Figure 42) was introduced into *Rosa hybrida* cv. *Kardinal* via *Agrobacterium*-mediated transformation.

#### Butterfly pea R3'5'H constructs

#### 5 Isolation of a R3'5'H cDNA clone from petals of *Clinovia ternatea* (butterfly pea)

#### Construction of butterfly pea petal cDNA library

A blue variety of *Clinovia ternatea* (butterfly pea, the seeds were kindly provided by Osaka Botanical Garden) was grown in a field in Osaka. Total RNA was isolated from fresh and pigmented petals at a pre-anthesis stage as described above. PolyA<sup>+</sup> RNA was prepared using Oligotex (Takara) according to the manufacturer's recommendation. A petal cDNA library of butterfly pea was constructed from the polyA<sup>+</sup> RNA using a directional λZAP-cDNA synthesis kit (Stratagene, USA) following the manufacturer's protocols.

#### Screening of butterfly pea cDNA library for a R3'5'H cDNA clone

15 The butterfly pea petal cDNA library was screened with DIG-labelled petunia R3'5'H petH1 cDNA clone as described previously (Tanaka *et al.*, *Plant Cell Physiol.* 37: 711-716, 1996). Two cDNA clones that showed high sequence similarity to the petunia R3'5'H petH1 were identified. The plasmid containing the longest cDNA clone was designated pBHR2 and the cDNA clone was sequenced. Alignment between the deduced amino acid sequences of the butterfly pea R3'5'H clone and the petunia R3'5'H petH1 clone (SEQ ID NO:2) revealed that the butterfly pea R3'5'H cDNA (contained in pBHR2) did not represent a full-length cDNA and lacked first 2 bases of the putative initiation codon. These two bases along with a BamHI restriction endonuclease recognition site were added to the cDNA clone using PCR and a synthetic primer, 5'-GGGATCCACAAATGTTCTTCTAAGAGAAAT-3' [SEQ ID NO:25] as described previously (Yonckura-Sakakibara *et al.*, *Plant Cell Physiol.* 41: 495-502, 2000). The resultant fragment was digested with the restriction endonucleases BamHI and PstI and the subsequent DNA fragment of about 200 bp was recovered. The DNA fragment was ligated with a 3.3 kb fragment of BamHI/EcoRI digested pBHR2 to yield pBHR2F (Figure 43). The DNA sequence was confirmed to exclude errors made during PCR (SEQ ID NO:20).

- 104 -

Comparison of the nucleotide sequence of butterfly pea *F3'5'H* clone (SEQ ID NO:20) with that of the petunia *F3'5'H* revealed around 59% identity to the petunia *F3'5'H petHf1* clone (SEQ ID NO:1) and 62% identity to the petunia *F3'5'H petHf2* clone (SEQ ID NO:3).

5

The binary vector pCGP2135 (*AmCHS* 5': butterfly pea *F3'5'H*: *petD8* 3')

The plasmid pCGP2135 (Figure 44) contains the butterfly pea *F3'5'H* cDNA clone between a snapdragon *CHS* promoter fragment (*AmCHS* 5') and a petunia *PLTP* terminator fragment (*petD8* 3') in tandem with the 35S 5': *SuRB* selectable marker gene cassette of the binary vector pCGP1988 (Figure 16).

10

The petunia *F3'5'H (petHf1)* cDNA clone in pCGP725 (described in Example 4) (Figure 7) was removed by initially digesting pCGP725 with the restriction endonucleases *Xba*I and *Bam*HI. The ends were repaired the ~4.9kb fragment containing the vector with the *AmCHS* 5' and *petD8* 3' fragments was purified and ligated with the repaired ends of the ~1.6kb *Xho*I/*Bam*HI fragment from pBHF2F (Figure 43) containing the butterfly pea *F3'5'H* cDNA clone to produce pCGP2133. Correct insertion of the butterfly pea *F3'5'H* fragment in tandem with the *AmCHS* 5' and *petD8* 3' fragments was confirmed by restriction endonuclease mapping.

15

20

The *AmCHS* 5': butterfly pea *F3'5'H*: *petD8* 3' cassette was then isolated from pCGP2133 by firstly digesting with the restriction endonuclease *No*zI. The ends of the linearised plasmid were repaired and then the chimeric *F3'5'H* gene was released upon digestion with the restriction endonuclease *Eco*RV. The ~3.6kb purified fragment was then ligated with *Asp*718 repaired ends of the binary vector pCGP1988 (described in Example 4) (Figure 16). Correct insertion of the fragment was established by restriction endonuclease analysis of plasmid DNA isolated from tetracycline-resistant transformants. The resulting plasmid was designated pCGP2135 (Figure 44).

25

**Carnation and petunia transformation with pCGP2135**

The T-DNA contained in the binary vector plasmid pCGP2135 (Figure 44) was introduced into *Dianthus caryophyllus* cultivars Kortina Chanel and Monte Lisa and *Petunia hybrida* cv. Sk4 x Sw63 via *Agrobacterium*-mediated transformation.

**The binary vector pBBBF5 (CaMV 35S; Butterfly pea F3'5'H: nos 3')**

The binary vector, pBB2113-GUS contains a GUS coding region between an enhanced *CaMV 35S* promoter and *nos* terminator in a pBI121 binary vector (Mitsunaka *et al.*, 1996, *supra*). The plasmid pBB2113-GUS was digested with the restriction endonuclease *SacI*. The overhanging ends were repaired and then ligated with a *SacI* linker to yield pBB2113-GUS. The 1.8 kb *BamHI-XhoI* fragment from pBHR2F was ligated with *BamHI-SacI* digested pBB2113-GUS to create pBBBF5 (Figure 45).

**Rose transformation with pBBBF5**

The T-DNA contained in the binary vector plasmid pBBBF5 (Figure 45) was introduced into *Rosa hybrida* cultivar Lavande via *Agrobacterium*-mediated transformation.

**The binary vector pCGP2134 (CaMV 35S; Butterfly pea F3'5'H: ocs 3')**

The binary vector pCGP2134 (Figure 46) contains a chimeric *CaMV 35S; butterfly pea F3'5'H: ocs 3'* gene cassette in a tandem orientation with the 35S 5' *SwRB* selectable marker gene cassette of the binary vector pCGP1988 (Figure 16).

**Intermediates in the preparation of the binary vector pCGP2134**

The butterfly pea *F3'5'H* cDNA clone was released upon digestion of the plasmid pBHR2F (Figure 43) with the restriction endonucleases *XhoI* and *BamHI*. The overhanging ends were repaired and the ~1.7 kb fragment was ligated with the *PstI* (repaired ends)/*EcoRV* ends of pCGP2105 (described in Example 4) (Figure 17). Correct insertion of the fragment was established by restriction endonuclease analysis of plasmid DNA isolated from ampicillin-resistant transformants. The resulting plasmid was designated pCGP2132.

- 106 -

An ~3.6 kb fragment containing the *CaMV 35S: butterfly pea F3'5'H: ocs 3'* chimeric gene cassette was released upon digestion with the restriction endonucleases *XhoI* and *XbaI*. The overhanging ends were repaired and the purified fragment was ligated with *Asp718* repaired ends of the binary vector, pCGP1988 (described in Example 4) (Figure 16).

- 5 Correct insertion of the fragment was established by restriction endonuclease analysis of plasmid DNA isolated from tetracycline-resistant transformants. The resulting plasmid was designated pCGP2134 (Figure 46).

**Rose transformation with pCGP2134**

- 10 The T-DNA contained in the binary vector plasmid pCGP2134 (Figure 46) was introduced into *Rosa hybrida* cv. Kardinal via *Agrobacterium*-mediated transformation.

**Gentia F3'5'H constructs**

**Isolation of a F3'5'H cDNA clone from petals of Gentiana triflora (gentian).**

- 15 **Construction and screening of a gentian petal cDNA library**

The isolation of a gentian cDNA encoding *F3'5'H* has been described previously (Tanaka *et al.*, 1996, *supra*) and is contained within the plasmid pG48 (Figure 47). Comparison of the nucleotide sequence of the gentia *F3'5'H* clone (*Gent#48*) (SEQ ID NO:22) contained in the plasmid pG48 (Figure 47) with that of the petunia *F3'5'H* revealed around 61% identity to the petunia *F3'5'H petHf1* clone (SEQ ID NO:1) and 64% identity to the petunia *F3'5'H petHf2* clone (SEQ ID NO:3).

**The binary vector pCGP1498 (AmCHS 5': gentia F3'5'H: petD8 3')**

- 25 The plasmid pCGP1498 (Figure 48) contains the gentia *F3'5'H* (*Gent#48*) cDNA clone between a snapdragon *CHS* promoter fragment (*AmCHS 5'*) and a petunia *PLTP* terminator fragment (*petD8 3'*) in tandem with the *35S 5': SuRB* selectable marker gene cassette of the binary vector pWTT2132 (Figure 6).

- 30 The petunia *F3'5'H (petHf1)* cDNA clone in pCGP725 (described in Example 4) (Figure 7) was removed by initially digesting pCGP725 with the restriction endonucleases *XbaI* and *BamHI*. The ends were repaired the ~4.9kb fragment containing the vector with the

*AmCHS* 5' and *petD8* 3' fragments was purified and ligated with the repaired ends of the ~1.7 kb *XhoI/BamHI* fragment from pG48 (Figure 47) containing the *genita F3'5'H* cDNA clone to produce pCGP1496. Correct insertion of the *genita F3'5'H* fragment in tandem with the *AmCHS* 5' and *petD8* 3' fragments was confirmed by restriction endonuclease mapping.

The *AmCHS* 5'; *genita F3'5'H*; *petD8* 3' cassette was then isolated from pCGP1496 by firstly digesting with the restriction endonuclease *NcoI*. The overhanging ends of the linearised plasmid were repaired and then the chimeric *F3'5'H* gene was released upon digestion with the restriction endonuclease *EcoRV*. The ~3.6kb purified fragment was then ligated with *Asp718* repaired ends of the binary vector pWTT2132 (Figure 6). Correct insertion of the fragment was established by restriction endonuclease analysis of plasmid DNA isolated from tetracycline-resistant transformants. The resulting plasmid was designated pCGP1498 (Figure 48).

#### Curatlon and *petunia* transformation with pCGP1498

The T-DNA contained in the binary vector plasmid pCGP1498 (Figure 48) was introduced into *Dianthus caryophyllus* cultivars Kortina Chanel and Monte Lisa and *Petunia hybrida* cv. Sk4 x Sw63 via *Agrobacterium*-mediated transformation.

#### The binary vector pBEGHF48 (CaMV 35S; *genita F3'5'H*; nos 3)

The *genita F3'5'H* cDNA clone was released by digestion of the plasmid pG48 with the restriction endonucleases *BamHI* and *XhoI*. The resulting ~1.7 kb DNA fragment was isolated and ligated with *BamHI/SalI* digested pBR2113-GUS (Mitsubara *et al.*, 1996, *supra*) to create pBEGHF48 (Figure 49).

#### Rose transformation with pBEGHF48

The T-DNA contained in the binary vector plasmid pBEGHF48 (Figure 49) was introduced into *Rosa hybrida* cv. Lavande via *Agrobacterium*-mediated transformation.



- 108 -

**The binary vector pCGP1982 (CaMV 35S: gentia F3'5'H: ocs 3')**

The binary vector pCGP1982 (Figure 50) contains a chimeric *CaMV 35S: gentia F3'5'H: ocs 3'* gene cassette in tandem with the *35S 5': SuRB* selectable marker gene cassette of the binary vector pWTT2132 (Figure 6).

5

**Intermediates in the preparation of the binary vector pCGP1982**

The plasmid pG48 (Figure 47) was linearised upon digestion with the restriction endonuclease *Asp718*. The overhanging ends were repaired and then the *gentia F3'5'H* cDNA clone (*Gent#48*) was released upon digestion with the restriction endonuclease *Bam*HI. The ~1.7 kb fragment was ligated with the 5.95kb *Eco*RI (repaired ends)/*Bam*HI fragment of pKIWI101 (Janssen and Gardner, 1989, *supra*). Correct insertion of the fragment was established by restriction endonuclease analysis of plasmid DNA isolated from ampicillin-resistant transformants. The resulting plasmid was designated pCGP1981.

10

15 An ~3.6 kb fragment containing the *CaMV 35S: gentia F3'5'H: ocs 3'* chimeric gene cassette was released upon digestion of the plasmid pCGP1981 with the restriction endonucleases *Xho*I and *Xba*I. The overhanging ends were repaired and the purified fragment was ligated with repaired ends of *Asp*718 digested binary vector, pWTT2132. Correct insertion of the fragment was established by restriction endonuclease analysis of plasmid DNA isolated from tetracycline-resistant transformants. The resulting plasmid was designated pCGP1982 (Figure 50).

20

**Rosa transformation with pCGP1982**

The T-DNA contained in the binary vector plasmid pCGP1982 (Figure 50) was introduced into *Rosa hybrida* cv. Kardinal via *Agrobacterium*-mediated transformation.

25

#### Lavender F3'5'H constructs

##### Isolation of a F3'5'H cDNA clone from petals of *Lavandula nil* (lavender)

##### Construction of lavender petal cDNA library

Cut flowers of a violet variety of *Lavandula nil* were purchased from a florist. Total RNA was isolated from fresh and pigmented petals as described above. PolyA<sup>+</sup> RNA was prepared using Oligotex (Takara) according to the manufacturer's recommendations. A petal cDNA library of lavender was constructed from the polyA<sup>+</sup> RNA using a directional AZAP-cDNA synthesis kit (Stratagene, USA) following the manufacturer's protocols.

##### Screening of lavender cDNA library for a F3'5'H cDNA clone

The lavender petal cDNA library was screened with DIG labelled petunia F3'5'H *petH1* cDNA clone as described previously (Tanaka *et al.*, 1996, *supra*). One cDNA clone (LBG) that showed high similarity to petunia F3'5'H *petH1* was identified and the plasmid was designated pLHF8 (Figure 51). The nucleotide sequence of the lavender F3'5'H (LBG) cDNA clone was designated as SEQ ID NO: 31.

Comparison of the nucleotide sequence of lavender F3'5'H clone with that of the petunia F3'5'H cDNA clones revealed around 59% identity to the petunia F3'5'H *petH1* clone (SEQ ID NO:1) and 60% identity to the petunia *petH2* clone (SEQ ID NO:3).

##### The binary vector pBELF8 (CaMV 35S: lavender F3'5'H: nos 3)

The plasmid of pLHF8 (Figure 51) was digested with the restriction endonucleases *Bam*HI and *Xho*I to release a DNA fragment of approximately 1.8 kb. The ~1.8kb purified fragment from pLHF8 was then ligated with the *Bam*HI-*Xho*I digested ends of the plasmid pB2113-GUS (described above) to create pBELF8 (Figure 52).

##### Rose transformation with pBELF8

The T-DNA contained in the binary vector plasmid pBELF8 (Figure 52) was introduced into *Rosa hybrida* cultivar Lavande via *Agrobacterium*-mediated transformation.

30

25

20

15

10

5

- 110 -

**EXAMPLE 8*****Analysis of transgenic carnation, petunia and rose***

The transgenic plants produced in the experiments described in Example 7 were grown to flowering. Flowers were collected and the colors of the petals were coded using the Royal Horticultural Society Colour Charts (RHSCC). The anthocyanins were extracted and the anthocyanidins analysed by spectrophotometric, TLC and/or HPLC analysis. Total RNA was also isolated from petal tissue of the appropriate stages of flower development and Northern blot analysis was used to detect transcripts of *F3'5'H* transgenes. The results of the transgenic analysis are summarised in Tables 11, 12 and 13.

**Carnation**

The *F3'5'H* genes described in Example 7 were evaluated for their ability to lead to the production of delphinidin-based pigments in carnation petals. Two carnation cultivars, Kortina Chanel (KC) and Monte Lisa (ML), were used in the transformation experiments.

The carnation cultivar Kortina Chanel produces pink colored flowers that normally accumulate cyanidin-based anthocyanins. This cultivar therefore contains a carnation *F3'H* and DFR activity that an introduced *F3'5'H* would need to compete with for substrate. The carnation cultivar Monte Lisa produces brick red colored flowers that normally accumulate pelargonidin-based anthocyanins. This cultivar is thought to lack fully functional *F3'H* activity and contain a DFR that is capable of acting on DHK and thus an introduced *F3'5'H* would only be required to compete with the endogenous DFR for substrate.

- 111 -

TABLE 11 Results of transgenic analysis of petals from carnations transformed with T-DNAs containing *F3'5'H* gene expression cassettes (*AmCHS 5'::F3'5'H::petD8 3'*).

<i>F3'5'H</i>	pCGP	cv.	#tg	TL-C+	HPLC+	Highest % del	Av. % del	Northern+
<i>Salvia#2</i>	2121	KC	22	2/16	3/4	12.5%	7%	nd
		ML	21	17/18	9/9	76%	57%	14/15
<i>Salvia#47</i>	2122	KC	23	6/12	8/8	29%	12%	nd
		ML	25	21/22	17/17	88%	56%	12/14
<i>Solysa</i>	2130	KC	30	22/27	17/17	35%	11%	nd
		ML	23	14/15	14/14	76%	49%	13/14
<i>Butterfly pea</i>	2135	KC	22	0/16	0/1	nd	nd	nd
		ML	24	19/20	13/13	23%	10%	14/14
<i>Gentian</i>	1498	KC	22	0/14	nd	nd	nd	7/8
		ML	2	2/2	1/1	nd	nd	1/2
<i>pansy BP#18</i>	1972	KC	26	18/20	12/12	14%	9%	19/19
		ML	21	15/16	8/8	80%	66%	14/16
<i>pansy BP#40</i>	1973	KC	26	11/15	7/8	18%	8%	13/17
		ML	33	19/22	20/20	72%	52%	12/15
<i>petunia</i>	1452	KC	104	41/64	nd	3.5%	1.3%	15/17
<i>petiti</i>		ML	48	39/41	26/26	75%	30%	12/13
<i>petunia</i>	1524	ML	27	18/19	17/17	81%	41%	12/14

5 *F3'5'H* = *F3'5'H* sequence contained on the T-DNA  
pCGP = plasmid pCGP number of the binary vector used in the transformation  
cv. = cultivar  
KC = carnation cultivar Kortina Chanel (cyanidin line)  
ML = carnation cultivar Monte Lisa (pelargonidin line)  
#tg = total number of transgenics produced

- 112 -

TLC+ = number of individual events in which delphinidin or delphinidin-based molecules was detected in petals (as determined by TLC) over the total number of individual events analyzed

HPLC+ = number of individual events in which delphinidin or delphinidin-based molecules was detected in petals (as determined by HPLC) over the total number of individual events analyzed

Highest % del = Highest % delphinidin or delphinidin-based molecules detected in the petals for the population of transgenic events

Av % del = average % delphinidin or delphinidin-based molecules detected in the petals for the population of transgenic events

Northern = number of individual events in which the specific intact *F3'5'H* transcripts were detected by Northern blot analysis in total RNA isolated from petals over the total number of events analyzed

nd = not done

The results suggest that all of the *F3'5'H* sequences evaluated (petunia *petHf1*, petunia *petHf2*, *Salvia Sal#2*, *Salvia Sal#47*, *Sollya Sol#5*, Butterfly pea *BpeaHF2*, pansy *BP#18*, pansy *BP#40* and *Gentian Gen#48*) were stable in carnation and resulted in the production of novel delphinidin-based pigments in carnation flowers. Intact transcripts of each *F3'5'H* were detected by Northern blot analysis in total RNA isolated from petals of the transgenic carnations.

#### Petunia

The *F3'5'H* genes described in Example 7 were evaluated for their ability to lead to the production of delphinidin-based pigments in petunia petals. The *P. hybrida* F1 hybrid Skr4 x SW63 which is homozygous recessive for *Hf1* and *Hf2*, was used in the transformation experiments. Although Skr4 x SW63 is homozygous recessive for *Hf1* and *Hf2*, these mutations do not completely block production of the endogenous *F3'5'H* (see US Patent Number 5,349,125) and low levels of malvidin are produced to give the petal limb a pale lilac color. Malvidin is the methylated derivative of the 3'5'-hydroxylated pigment, delphinidin or delphinidin-based molecules (Figures 1A and 1B). Spectrophotometric

analysis was used as a measure of total anthocyanins accumulating in petals from the transgenic petunia flowers. The increased level of anthocyanins and/or the color change detected was used as a guide to the efficacy of the *F3'5'H* gene under evaluation.

5 TABLE 12 Results of transgenic analysis of petals from *P. hybrida* cv Skra x SW63 plants transformed with T-DNAs containing *F3'5'H* gene expression cassettes (*AmCHS 5'*: *F3'5'H*: *peid8 3'*).

<i>F3'5'H</i>	pCGP	#tg	TLC+	Col	†A/c	Best	Av.	Northern+	Best color
control	na	na	na	na	na	14-	250	0	7SC
Gentiana#48	1498	22	3/5	18/20	nd			6/6	72B/78A
Butterfly pea	2135	24	18/20	22/24	23/24	4427	2397	nd	74A/78A
Kennedia	2256	24	22/24	22/24	22/24	4212	2592	nd	74A/78A
Salvia#2	2121	24	21/24	21/24	21/24	2471	1730	nd	78A
Salvia#47	2122	19	17/19	16/19	16/19	2634	1755	nd	78A/80A
Soilya#5	2130	22	14/16	13/16	13/16	3446	1565	nd	78A
penary BP#18	1972	22	nd	20/22	nd	nd	nd	9/9	74A/B
penary BP#40	1973	19	8/8	18/19	18/20	2583	1556	nd	74/78A
petunia <i>petH7</i>	484	16	nd	9/16	8/15	2683	1250	nd	74A/B
petunia <i>petH2</i>	1524	20	nd	18/20	8/8	4578	2357	8/8	74A/B

10 *F3'5'H* = *F3'5'H* sequence contained on the T-DNA  
pCGP = plasmid pCGP number of the binary vector used in the transformation  
#tg = total number of transgenes produced

- 114 -

TLC+ = number of individual events in which malvidin was detected in the flowers (at a level above the Skr4 x Sw63 background) (as determined by TLC) over the total number of individual events analyzed

Col = number of individual events that produced flowers with an altered flower color compared to the control over the total number examined

† A/c = number of individual events that had an increased level of anthocyanins in petals as measured by spectrophotometric analysis of crude extracts over the number of individual events analyzed (in  $\mu\text{moles/g}$ )

Best = highest anthocyanin amount as measured by spectrophotometric analysis of crude extracts from a flower of an individual event (in  $\mu\text{moles/g}$ )

Av = the average amount of anthocyanin detected as measured by spectrophotometric analysis of crude extracts from a flower in the population of transgenic flowers analysed (in  $\mu\text{moles/g}$ )

Northern = number of individual events in which the specific intact *F3'5'H* transcripts were detected by Northern blot analysis in total RNA isolated from petals over the total number of events analyzed

Best color = most dramatic color change recorded for the transgenic population

nd = not done

na = not applicable

Introduction of the *F3'5'H* gene expression cassettes into Skr4 x SW63 led to a dramatic flower color change from pale lilac to purple with a dramatic increase in the production of malvidin in the petals..

The results suggest that all of the *F3'5'H* sequences tested (petunia *petHf1*, petunia *petHf2*, *Salvia Sal#2*, *Salvia Sal#47*, *Sollya Sol#5*, Butterfly pea *BpeaHF2*, pansy *BP#18*, pansy *BP#40*, *Gentian Gen#48*, *Kennedia Kenn#31*) were stable in petunia petals and resulted in the complementation of the *Hf1* or *Hf2* mutation in the Skr4 x SW63 petunia line leading to dramatically increased levels of malvidin accumulation with a concomitant color change

## Rose

The *F3'5'H* genes described in Example 7 were evaluated for their ability to lead to the production of delphinidin-based pigments in rose petals. A selection of three rose cultivars, Cardinal (Kard), Soft Promise (SP) or Lavanade (Lav) were used in transformation experiments. The rose cultivar Cardinal produces red colored flowers that normally accumulate cyanidin-based anthocyanins. This cultivar therefore contains rose *F3'H* and *DFR* activities that the introduced *F3'5'H* would need to compete with for substrate. The rose cultivar Lavanade produces light pink colored flowers that normally accumulate cyanidin-based anthocyanins. This cultivar therefore contains functional rose *F3'H* and *DFR* activities that the introduced *F3'5'H* would need to compete with for substrate. The rose cultivar Soft Promise produces apricot colored flowers that normally accumulate pelargonidin. This cultivar is thought to lack a fully functional rose *F3'H* activity and contain a *DFR* that is capable of acting on DHK and thus the introduced *F3'5'H* would only be required to compete with the endogenous rose *DFR* for substrate.

TABLE 13 Results of transgenic analysis of petals from roses transformed with DNAs containing *F3'5'H* gene expression cassettes (*CaMV 35S::F3'5'H::ocs 3'*).

<i>F3'5'H</i>	plasmid	Cult	#g	TLC +	HPLC +	Highest % del	Av. % del	Northern+
Salvia2	pCGP2120	Kard	30	18/20	21/21	12%	5%	18/18
Salvia47	pCGP2119	Kard	22	11/16	9/9	7.1%	2%	12/15
Soliva	pCGP2131	Kard	27	0/23	2/2	1%	0.5%	6/6
Butterfly	pCGP2134	Kard	29	0/15	nd	na	na	0/9
pea	pBBBFS	Lav	25	nd	0/25	0%	0%	nd
Gentian	pCGP1482	Kard	27	0/23	nd	na	na	0/23
	pBBGHR48	Lav	23	nd	0/23	0%	0%	0/23
pansy	pCGP1967	Kard	56	30/33	33/34	58%	12%	21/21
		SP	36	21/24	18/18	65%	35%	16/21
pansy	pCGP1969	Kard	22	15/15	15/15	24%	9%	16/16
		SP	37	17/17	16/17	80%	54%	11/13



- 116 -

<i>F3'5'H</i>	plasmid	Cult	#tg	TLC+	HPLC+	Highest % del	Av. % del	Northern+
Petunia <i>petHf1</i>	pCGP1638	Kard	22	0/21	nd	na	na	0/16
	pCGP1392	Lav.	34	nd	0/34	0%	0%	nd
Petunia <i>petHf2</i>	pCGP2123	Kard	41	0/26	nd	na	na	0/10
Lavender	pBELF8	Lav	28	nd	4/28	4%	3.5%	nd

*F3'5'H* = the *F3'5'H* sequence contained on the T-DNA

plasmid = the plasmid number of the binary vector used in the transformation experiment

5 Cult = *Rosa hybrida* cultivar

Kard = Cardinal

SP = Soft Promise

Lav = Lavande

#tg = # of independent transgenic events produced

10 TLC+ = number of individual events that accumulated detectable delphinidin or delphinidin-based molecules (as determined by TLC) in the petals over the number of individual events analyzed

HPLC+ = number of individual events that accumulated detectable delphinidin or delphinidin-based molecules (as determined by HPLC) in the petals over the number of

15 individual events analyzed

Northern = number of individual events in which the specific intact *F3'5'H* transcripts were detected by Northern blot analysis in total RNA isolated from petals over the total number of events analyzed

20 nd = not done

- The results suggest surprisingly that not all of the *F3'5'H* sequences assessed (petunia *petH1*, petunia *petH2*, Salvia *Sal#2*, Salvia *Sal#17*, Soliya *Soli#5*, Butterfly pea *BpeaHR2*, pansy *BP#18*, pansy *BP#40*, Gentian *Gent#48*, Kennedia *Kenn#31* and Lavender *LBG*) were functional in rose. In fact transcripts of the introduced *F3'5'H* sequences isolated from *Clitoria ternatea* (butterfly pea), *Gentiana triflora*, (gentian) and *Petunia hybrida* (petunia) failed to accumulate in rose petals. Only full-length *F3'5'H* transcripts from pansy, salvia, kennedia, soliya and lavender accumulated in rose petals. However although Kennedia *F3'5'H* transcripts did accumulate in rose petals, there was either no accumulation of the enzyme or the enzyme produced was either not functional or was unable to compete with the endogenous rose *F3'H* and *DFR* enzymes to allow for the production of delphinidin or delphinidin-based molecules pigments. Of the *F3'5'H* sequences evaluated, only the *F3'5'H* sequences derived from cDNA clones from *Salvia spp.* (*Sal#2* and *Sal#47*), *Viola spp.* (*BP#18* and *BP#40*), *Soliva spp.* (*Soli#5*) and *Lavandula nil* (*LBG*) resulted in the production of delphinidin or delphinidin-based molecules based pigments in rose petals. Based on the relative percentages of delphinidin or delphinidin-based molecules produced in rose petals, the *F3'5'H* sequences from pansy (*BP#18* and *BP#40*) were revealed to be the most effective of those assessed at producing delphinidin or delphinidin-based molecules in rose petals.
- 20 Introduction of *Viola spp. F3'5'H* sequence into *Rosa hybrida* cv. Medeo and Pamela As described in the introduction, copigmentation with other flavonoids, further modification of the anthocyanidin molecule and the pH of the vacuole impact on the color produced by anthocyanins. Therefore, selection of rose cultivars with relatively high levels of flavonols and relatively high vacuolar pH would result in bluer flower colors upon production of delphinidin or delphinidin-based molecules pigments.
- 25 The rose cultivar Medeo generally produces cream-colored to pale apricot flowers (RHSCC 158C to 159A). HPLC analysis of the anthocyanidins and flavonols accumulating in Medeo rose petals revealed that the petals accumulate high levels of flavonols (2.32 mg/g kaempferol, 0.03 mg/g quercetin) and very low levels of anthocyanins (0.004 mg/g
- 30

- 118 -

cyanidin, 0.004 mg/g pelargonidin). The estimated vacuolar pH of Medeo petals is around 4.6.

5 The rose cultivar Pamela produces white to very pale pink colored flowers. It similarly accumulates low levels of anthocyanin and relatively high levels of flavonols.

The T-DNA contained in the construct pCGP1969 (Figure 30) incorporating the pansy *F3'5'H* clone, *BP#40*, was also introduced into the rose cultivars Medeo and Pamela resulting in the production of over 90% delphinidin or delphinidin-based molecules in  
10 these roses and leading to a dramatic color change and novel colored flowers. The most dramatic color change in transgenic Medeo flowers was to a purple/violet color of RHSCC 70b, 70c, 80c, 186b. The most dramatic color change in transgenic Pamela flowers was to a purple/violet color of RHSCC 71c, 60c, 71a, 80b.

15 In conclusion, two unexpected findings were revealed when gene sequences that had been proven to lead to functionality in petunia and carnation were introduced into roses.

First, the petunia *F3'5'H* *pethf1* (and *pethf2*) sequences that had resulted in novel color production in carnation and also proven to lead to synthesis of a functional enzyme in  
20 petunia did not lead to full-length (or intact) transcript accumulation (as detectable by Northern blot analysis) in rose petals. In fact, there was either no accumulation of full-length or intact transcript or the transcripts that were detected were degraded and were seen as low MW (or fast migrating) smears on RNA blots indicating the presence of low MW heterologous hybridizing RNA. Therefore in order to find a *F3'5'H* sequence that  
25 would accumulate in rose and lead to a functional enzyme, a number of *F3'5'H* sequences were isolated. Again it was not obvious which sequence would lead to an active enzyme in rose petals. All of the *F3'5'H* sequences isolated were tested for functionality in carnation and/or petunia and all led to accumulation of intact transcripts and production of a functional *F3'5'H* activity. However only *F3'5'H* sequences from pansy (*BP#18* and  
30 *BP#40*), salvia (*Sal#2* and *Sal#47*), sollya (*Soll#5*), kennedia (*Kenn#31*) and lavender (*LBG*) resulted in accumulation of intact full-length transcripts and only those from pansy

- 119 -

(BP#18 and BP#40), salvia (Sal#2 and Sal#47), solya (Soli#5) and lavender (LBG) resulted in production of a functional enzyme in rose as measured by the synthesis of delphinidin or delphinidin-based molecules.

Secondly that it was not obvious which promoters would be effective in rose. Promoter cassettes that had been tested and proven to be functional in carnation and petunia flowers did not lead to accumulation of detectable transcripts in rose petals. Of the promoters tested in rose, only *CaMY 35S*, *RoseCHS 5'*, *ChrysCHS 5'*, *mas 5'* and *nos 5'* promoters led to intact and detectable *GLS* or *nptII* or *SuRB* transcript accumulation in rose petals.

Table 14 shows a summary of the results obtained when assessing F3'5'H sequences from various species in petunia, carnation and rose.

TABLE 14 Summary of effectiveness of the F3'5'H sequences in petunia, carnation and rose

F3'5'H	Petunia			Carnation			Rose	
	Mal	RNA	Del	RNA	Del	RNA	Del	RNA
Kennedia (Kenn#31)	+	nd	nd	nd	+	-	+	+
Gentian (Gent#48)	+	+	+	+	+	-	-	-
Salvia (Sal#2)	+	nd	+	+	+	+	+	+
Salvia (Sal#47)	+	nd	+	+	+	+	+	+
Solysa (Soli#5)	+	nd	+	+	+	+	+	+
Butterfly	+	nd	+	+	+	-	-	-
pea (BpcaHF2)	+	nd	+	+	+	+	+	+
Pansy (BP#18)	+	+	+	+	+	+	+	+
Pansy (BP#40)	+	nd	+	+	+	+	+	+
Petunia (petH1)	+	+	+	+	+	+	-	-
Petunia (petH2)	+	+	+	+	+	+	-	-
Lavender (LBG)	nd	nd	nd	nd	nd	nd	+	+

- 120 -

*nd* = not done

*Mal* = malvidin detected in petals as analysed by TLC

*Del* = delphinidin or delphinidin-based molecules detected in petals as analysed by TLC or HPLC

5 + = yes

- = no

#### EXAMPLE 9

*Use of pansy F3'5'H sequences in species other than rose*

##### 10 *Gerbera*

From the examples above, it was clear that the pansy *F3'5'H* sequences, *BP#18* and *BP#40*, resulted in functional *F3'5'H* activity and lead to the production of high levels of delphinidin or delphinidin-based molecules in roses and carnations.

15 The T-DNA from binary construct pCGP1969 (described in Example 8) (Figure 30) containing the chimeric *CaMV 35S: pansy BP#40 F3'5'H: ocs 3'* gene expression cassette was introduced into the gerbera cultivar Boogie via *Agrobacterium*-mediated transformation, to test the functionality of the pansy *F3'5'H* sequence in gerbera.

20 Of six events produced to date, one (#23407) has produced flowers with a dramatic color change (RHSCC 70c) compared to the control flower color (RHSCC 38a, 38c).

The color change of the petals of the transgenic gerbera has been correlated with the presence of delphinidin or delphinidin-based molecules as detected by TLC.

25

##### *Other species*

In order to produce delphinidin or delphinidin-based molecules pigments in plants that do not normally produce delphinidin-based pigments and does not contain a flavonoid 3'5'-hydroxylase constructs containing a *F3'5'H* gene (such as but not limited to a chimaeric

30 *Viola spp.* and/or *Salvia spp.* and/or *Solhya spp.* and/or *Lavandula spp.* and/or *Kennedia spp.* *F3'5'H* gene) are introduced into a species that does not normally produce

delphinidia-based pigments. Such plants may include but are not limited to carnation, chrysanthemum, gerbera, *Euphorbia*, *Begonia* and apple.

#### EXAMPLE 10

5

#### *Characteristics of R3'5'H sequences evaluated in petunia, carnation and rose*

Geno regulation in eukaryotes is, in simple terms, facilitated by a number of factors which interact with a range of sequences proximal and distal to a nucleotide sequence coding for a given polypeptide. Engineering expression cassettes for introduction into plants for the generation of one or more traits is based on an understanding of gene regulation in eukaryotes in general and, in selected cases, plants in particular. The essential elements include a series of transcriptional regulation sequences typically, but not exclusively, located upstream or 5' to the point of transcription initiation. Such elements are typically described as enhancers and promoters, the latter being proximal to the point of transcription initiation. Immediately downstream from, or 3' to, the initiation of untranslated region (5'ut) which plays a role in transcript stability and translational efficiency. Such sequences, when engineered into expression cassettes, are frequently chimeric and may be derived from sequences naturally occurring adjacent to the coding sequence and/or adjacent to a given promoter sequence. The coding sequence (sometimes disrupted by introns) lies 3' to the 5'ut followed by a 3'ut important to transcript (mRNA) stability and translational efficiency. Sequences 3' to the end of the coding region and 3' to the 3'ut itself are denoted as terminator sequences. All these elements make up an expression cassette. In making direct comparisons between promoters or other elements it is important to maintain uniformity in the remaining elements of an expression cassette. Hence, when comparing the efficacy of various R3'5'H sequences it was possible to confine the sequences leading to instability and the subsequent autodegradation of engineered mRNA and resultant absence of tri-hydroxylated products (delphinidin or delphinidin-based molecules derivatives) to the region coding for the R3'5'H and not to other elements in the expression cassette such as 5' ut and/or 3'ut sequences for example.

- 122 -

In an attempt to identify motifs or similarities between the *F3'5'H* sequences that resulted in full-length transcripts being detected in total RNA isolated from rose flowers, and ultimately delphinidin or delphinidin-based molecules production, comparisons across a range of parameters were performed. These included sequence identities at nucleic acid and amino acid levels, sequence alignments, taxonomic classifications, % of A or T nucleotides present in the sequence, % of codons with an A or T in the third position etc.

#### *Taxonomic classification*

- 10 The taxonomy of each species from which the *F3'5'H* sequences were isolated was examined (Table 15). There appeared to be no obvious link between the subclass classification and whether the *F3'5'H* sequence resulted in an intact transcript and subsequent delphinidin or delphinidin-based molecules production in roses.
- 15 Table 15: Taxonomic classifications of the species that *F3'5'H* sequences were isolated from and whether the use of the sequences resulted in intact transcript in rose petals that were detectable by RNA blot analysis.

Flower	Species	Family	Order	Subclass	Intact transcript	Delphinidin in rose petals
gentian	<i>Gentiana iriflora</i>	Gentianaceae	Gentianales	Asteridae	NO	NO
lavender	<i>Lavandula nil</i>	Lamiaceae	Lamiales	Asteridae	YES	YES
salvia	<i>Salvia spp.</i>	Lamiaceae	Lamiales	Asteridae	YES	YES
sallya	<i>Sollya spp.</i>	Pittosporaceae	Apiales	Asteridae	YES	YES
petunia	<i>Petunia hybrida</i>	Solanaceae	Solanales	Asteridae	NO	NO
kennedia	<i>Kennedia spp.</i>	Fabaceae	Fabales	Rosidae	YES	NO
butterfly pea	<i>Clitoria ternatea</i>	Fabaceae	Fabales	Rosidae	NO	NO
pansy	<i>Viola spp.</i>	Violaceae	Malpighiales	Rosidae	YES	YES
rose	<i>Rosa hybrida</i>	Rosaceae	Rosales	Rosidae	na	na

5 The nucleotide sequence identities between each of the F3'5'H sequences evaluated were determined using the ClustalW program (Thompson *et al.*, 1994, *supra*) within the MacVector™ version 6.5.3 application program (Oxford Molecular Ltd., England) (Table 16). There were no obvious differences between the F3'5'H sequences that resulted in the detection of intact full-length transcripts in RNA isolated from rose petals and those that didn't.

transcripts being detected in RNA isolated from rose petals are underlined and in italics.

	BP18	100	82	60	61	62	51	60	62	62	59	62	BP18	Perht
	BP40		100	60	57	58	50	59	62	58	60	62	BP40	Perht
	Lx		100	68	68	48	57	57	58	59	59	60	Lx	Perht
	Sol17			100	95	48	56	57	57	59	57	58	Sol17	Perht
	Sol2				100	49	57	58	60	57	59	59	Sol2	Perht
	Sol				100	48	50	50	48	50	51	51	Sol	Perht
	Kem						100	70	56	64	60	60	Kem	Perht
	Bpes							100	59	59	62	62	Bpes	Perht
	Gant									61	64	64	Gant	Perht
	Perht1									100	84	84	Perht1	Perht
	Perht2										100	100	Perht2	Perht



- 124 -

***Comparison of F3'5'H translated nucleotide sequences***

The translated nucleotide sequence identities and similarities between each of the F3'5'H sequences evaluated were also determined using the ClustalW program (Thompson *et al.*, 1994, *supra*) within the MacVector™ version 6.5.3 application program (Oxford Molecular Ltd., England) (Table 17). There were no obvious differences between the F3'5'H sequences that resulted in the detection of intact full-length transcripts in RNA isolated from rose petals and those that didn't.

Table 17: Percentage of the amino acid sequence identity and similarity (in brackets) between F3'5'H sequences isolated from various species. F3'5'H sequences that resulted in intact transcripts being detected in RNA isolated from rose petals are underlined and in italics.

	<u><i>BF18</i></u>	<u><i>BF40</i></u>	<u><i>Lav</i></u>	<u><i>Sal47</i></u>	<u><i>Sal2</i></u>	<u><i>Sal</i></u>	<u><i>Kana</i></u>	<u><i>Bpea</i></u>	<u><i>Gent</i></u>	<u><i>PetHf1</i></u>	<u><i>PetHf2</i></u>
<u><i>BF18</i></u>	100	91 (94)	65 (77)	65 (75)	65 (76)	44 (63)	69 (83)	64 (75)	69 (80)	74 (85)	74 (85)
<u><i>BF40</i></u>		100	67 (89)	66 (77)	66 (77)	46 (64)	69 (82)	64 (75)	68 (79)	74 (85)	75 (86)
<u><i>Lav</i></u>			100	75 (86)	75 (86)	45 (63)	63 (79)	59 (74)	66 (80)	68 (82)	69 (83)
<u><i>Sal47</i></u>				100	98	45 (65)	64 (78)	60 (72)	64 (76)	68 (79)	69 (81)
<u><i>Sal2</i></u>					100	45 (65)	64 (78)	60 (72)	63 (75)	68 (79)	69 (81)
<u><i>Sal</i></u>						100	46 (66)	41 (61)	44 (62)	46 (67)	46 (66)
<u><i>Kana</i></u>							100	72 (80)	65 (75)	71 (83)	72 (83)
<u><i>Bpea</i></u>								100	69 (81)	65 (75)	65 (74)
<u><i>Gent</i></u>									100	73 (82)	73 (82)
<u><i>PetHf1</i></u>										100	93 (95)
<u><i>PetHf2</i></u>											100

There is some evidence to suggest that the choice of codons influences the rate of translation and mRNA degradation. Certain codons are used less frequently than others are and this may be related to the abundance of isoaccepting tRNAs. Transfer RNAs corresponding to rare codons are less abundant in *E. coli* and yeast than RNAs corresponding to preferred codons (van Hoof and Green, *Plant Molecular Biology*, 35: 383-387, 1997). Examples of altering codon usage and making a gene more "plant-like" are the bacterial *B.t.* toxin gene (reviewed in Diehn *et al.*, *Genet. Engin.*, 18: 83-99, 1996) and the jellyfish *gfp* gene (Haseloff *et al.*, *Proc. Natl. Acad. Sci USA*, 94: 2122-2127, 1997). However as commented in van Hoof and Green, (1997) (*supra*), the effect of eliminating the rare codons in the *B.t.* genes increased the GC content, thereby eliminating AT-rich sequences that may be responsible for improper recognition of introns and polyadenylation sites as well as removing instability determinants. Alteration of codon usage in the jellyfish *gfp* gene also resulted in removal of a cryptic intron (Haseloff *et al.*, 1996, *supra*). Studies examining the effect of codon usage and instability elements have generally been limited to differences between genes isolated from species in different kingdoms ie. bacterial versus animal versus plant. Within the plant kingdom, differences have been observed between the dicotyledons and the monocotyledons. Studies on transgenic plants have suggested that promoter fragments used to drive gene expression in dicotyledonous plants are not as effective when used in monocotyledonous plants (see Galun and Breiman, *Transgenic Plants*, Imperial College Press, London, England, 1997). Differences in the methylation and ultimate expression of a *DFR* transgene in *Pennisetum hybridum* (dicot) were detected when a maize (monocot) *DFR* cDNA was compared with a gerbera (dicot) *DFR* cDNA (Blomus *et al.*, *Molecular and General Genetics*, 248: 649-656, 1995). The conclusion was that the gerbera *DFR* cDNA had a higher AT content (lower GC content) and was more "compatible" with the genomic organization of *Pennisetum* preventing it being recognised as a foreign gene and hence silenced by methylation. (Rose along with causation and *Pennisetum* are dicotyledons and the *F3'5'H* genes tested were all isolated from dicotyledonous plants.) These points serve to illustrate that degradation and stability mechanisms are not understood in detail and differences appear between plants and other kingdoms and within the plant kingdom.

**Percentage of nucleotides A or T in the F3'5'H DNA sequences**

- 126 -

The content of A and T was examined in the *F3'5'H* cDNAs evaluated along with that of four flavonoid pathway genes (*F3'H*, *DFR*, *CHS*, *FLS*) that had been isolated from rose (Table 18). The third position of each codon (within the open reading frame) was also  
 5 examined and the percentage of codons with an A or a T in the third position was calculated (Table 18).

Table 18: Summary of the percentage amount of A or T dinucleotides in the *F3'5'H* sequences isolated and whether the *F3'5'H* resulted in full-length transcripts being  
 10 detected in rose petals by Northern blot analysis.

<i>F3'5'H</i> seq	%AT	% A or T in 3rd	RNA	Delphinidin
Viola BP#18	50	40	YES	YES
Viola BP#40	51	35	YES	YES
Salvia#2	48	33	YES	YES
Salvia#47	48	34	YES	YES
Solysa#5	54	54	YES	YES
LavenderLBG	50	37	YES*	YES
Kennedia#31	54	47	YES	NO
petunia <i>petHf1</i>	61	66	NO	NO
petunia <i>petHf2</i>	59	65	NO	NO
Gentian#48	57	57	NO	NO
Butterfly pea# <i>HF2</i>	57	53	NO	NO
rose <i>F3'H</i>	47	34	**	na
rose <i>CHS</i>	52	42	**	na

R3'5'H seq	%AT	% A or T in 3rd	RNA	Delphinidin
rose DFR	53	46	**	na
rose FLS	56	43	***	na

%AT = % of nucleotides that are A or T in the nucleic acid sequence  
 %A or T in 3<sup>rd</sup> = the percentage of codons that have an A or T in the third position

RNA = whether a full-length mRNA transcript was detected by Northern blot  
 analysis in total RNA isolated from rose petals

Del = whether any *delphinidin* or *delphinidin*-based molecules was detected by  
 TLC or HPLC in rose petals

YBS\*

—although Northern blot analysis of transgenic  
 roses transformed with the lavender R3'5'H expression cassette was not performed, it can  
 be assumed that full-length transcript was produced since *delphinidin* or *delphinidin*-based  
 molecules was detected in the rose petals.

rose R3'5'H (described in International Patent Application No. PCT/AU97/00124)

rose DFR (Tanaka et al., 1995, *supra*)

rose FLS (GenBank accession number AB038247)

rose CHS (GenBank accession number AB038246)

The AT content of the four rose sequences (above) encoding flavonoid pathway enzymes  
 had an AT content of between 47 and 56%. In general the AT content of the R3'5'H  
 sequences that resulted in intact transcripts in rose petals was between 48 and 54%.  
 However the R3'5'H sequences that did not result in intact transcripts accumulating in rose  
 petals generally had a higher AT content of between 57 and 61%. Hence the AT content of  
 the introduced R3'5'H genes into rose may be a factor in whether an intact transcript  
 accumulates in rose petals and so leads to production of R3'5'H and *delphinidin* or  
*delphinidin*-based molecules.

- 128 -

The nucleotide base at the third position of each codon of the four rose sequences encoding flavonoid pathway enzymes was generally an A or a T in 34 to 46% of the codons. In general *F3'5'H* sequences that resulted in intact transcripts in rose petals contained an A or a T in the third position of each codon in 33 to 54% of the codons. However the *F3'5'H* sequences that did not result in intact transcripts accumulating in rose petals generally contained an A or a T in the third position of each codon in 53 to 66% of the codons. So the percentage of codons with an A or a T in the third position of the introduced *F3'5'H* genes into rose may also be a factor in whether an intact transcript is accumulates in rose petals and so leads to production of *F3'5'H* and *delphinidin* or *delphinidin-based* molecules.

It may be that by altering the overall content of the nucleotides A and/or T in any *F3'5'H* DNA sequence that does not result in an intact transcript in rose such as but not limited to the *Petunia hybrida petHf1*, *Petunia hybrida petHf2*, *Clitoria ternatea* (butterfly pea) *BpeaHf2* or *Gentiana triflora* (gentian) *Gent#48*, to a level more consistent with that found in rose genes, intact transcripts will accumulate and result in the efficient translation of *F3'5'H* transcripts and so to *delphinidin* or *delphinidin-based* molecules accumulation in rose petals. One way of altering the AT content of the DNA sequence without altering the amino acid sequence is to target the degeneracy of the third position of each codon.

Those skilled in the art will appreciate that the invention described herein is susceptible to variations and modifications other than those specifically described. It is to be understood that the invention includes all such variations and modifications. The invention also includes all of the steps, features, compositions and compounds referred to or indicated in this specification, individually or collectively, and any and all combinations of any two or more of said steps or features.

## BIBLIOGRAPHY

- A look beyond transcription*, ASPP, USA, Bailey-Serres and Gallie, eds, 1998
- Altshul *et al.*, *J. Mol. Biol.* 215(3): 403-410, 1990.
- Altshul *et al.*, *Nucl. Acids Res.* 25: 3389-3402, 1997.
- Ausubel *et al.*, "Current Protocols in Molecular Biology" John Wiley & Sons Inc, 1994-1998, Chapter 15.
- Aviv and Leder, *Proc. Natl. Acad. Sci. USA* 69: 1408, 1972.
- Bevan, *Nucleic Acids Res* 12: 8711-8721, 1984.
- Bonner and Laskey, *Eur. J. Biochem.* 46: 83, 1974.
- Bodeny, Molecular and genetic regulation of Bronze-2 and other maize anthocyanin genes. Dissertation, Stanford University, USA, 1994
- Brendel *et al.*, *In A look beyond transcription*, ASPP, USA, Bailey-Serres and Gallie, eds, 1998.
- Brouillard and Dangles, *In: The Flavonoids - Advances in Research since 1986*, Harborne, J.B. (ed), Chapman and Hall, London, UK, 1-22, 1993.
- Brugliera *et al.*, *Plant J.* 5: 81-92, 1994.
- Brugliera, Characterization of floral specific genes isolated from *Petunia hybrida*. RMIT, Australia. PhD thesis, 1994.

- 130 -

Bullock *et al.*, *Biotechniques* 5: 376, 1987.

Comai *et al.*, *Plant Mol. Biol.* 15: 373-381, 1990.

Depicker *et al.*, *J Mol. and Appl. Genetics* 1: 561-573, 1982.

Diehn *et al.*, *Genet Engin*, 18: 83-99, 1996

Elomaa *et al.*, *Molecular and General Genetics*, 248: 649-656, 1995

Franck *et al.*, *Cell* 21: 285-294, 1980.

Galun and Breiman, *Transgenic Plants*, Imperial College Press, London, England, 1997

Guilley *et al.*, *Cell* 30: 763-773, 1982.

Hanahan, *J. Mol. Biol.* 166: 557, 1983.

Harpster *et al.*, *MGG* 212: 182-190, 1988.

Hasseloff *et al.*, *Proc. Natl. Acad. Sci USA*, 94: 2122-2127, 1997

Holton and Cornish, *Plant Cell* 7: 1071-1083, 1995.

Holton and Graham, *Nuc. Acids Res.* 19: 1156, 1990.

Holton *et al.*, *Nature* 366: 276-279, 1993a.

Holton *et al.*, *Plant J.* 4: 1003-1010, 1993b

- 131 -

Holton, Isolation and characterization of petal specific genes from *Petunia hybrida*. PhD thesis, University of Melbourne, Australia, 1992

Huang and Miller, *Adv. Appl. Math.* 12: 373-381, 1991.

Inoue *et al.*, *Gene* 96: 23-28, 1990.

Janssen and Gardner, *Plant Molecular Biology*, 14: 61-72, 1989

Jefferson, *et al.*, *EMBO J.* 6: 3901-3907, 1987.

Johnson *et al.*, In *A look beyond transcription*, ASPP, USA, Bailey-Serres and Gallie, eds, 1998.

Karn *et al.*, *Gene* 32: 217-224, 1984.

Lazo *et al.*, *Bio/Technology* 9: 963-967, 1991.

Lee *et al.*, *EMBO J.* 7: 1241-1248, 1988.

Lu *et al.*, *Bio/Technology* 9: 864-868, 1991.

Marchant *et al.*, *Molecular Breeding* 4: 187-194, 1998.

Marmur and Doty, *J. Mol. Biol.* 5: 109, 1962.

McBride and Summerville, *Plant Molecular Biology* 14: 269-276, 1990.

McIntire, *J. Am. Chem. Soc.* 85: 2149, 1964.

Mitsubara *et al.*, *Plant Cell Physiol.* 37: 49-59, 1996.



Mol *et al.*, *Trends Plant Sci.* 3: 212-217, 1998.

Pearson and Lipman, *Proc. Natl. Acad. Sci. USA* 85(8): 2444-2448, 1988.

Plant Molecular Biology Labfax, Croy (ed), Bios scientific Publishers, Oxford, UK, 1993.

Plant Molecular Biology Manual (2<sup>nd</sup> edition), Gelvin and Schilperoot (eds), Kluwer Academic Publisher, The Netherlands, 1994.

Robinson and Firoozabady, *Scientia Horticulturae* 55: 83-99, 1993.

Rout *et al.*, *Scientia Horticulturae* 81: 201-238, 1999.

Sambrook *et al.*, Molecular Cloning: A Laboratory Manual, Cold Spring Harbor Laboratories, Cold Spring Harbor, NY, USA, 1989.

Sambrook and Russell, Molecular Cloning: A Laboratory Manual 3<sup>rd</sup> edition, Cold Spring Harbor Laboratories, Cold Spring Harbor, NY, USA, 2001

Seitz and Hinderer, Anthocyanins. In: *Cell Culture and Somatic Cell Genetics of Plants*. Constabel, F. and Vasil, I.K. (eds.), Academic Press, New York, USA, 5: 49-76, 1988.

Short *et al.*, *Nucl. Acids Res.* 16: 7583-7600, 1988.

Sommer and Saedler, *Mol Gen. Gent.*, 202: 429-434, 1986.

Strack and Wray, In: *The Flavonoids -Advances in Research since 1986*. Harborne, J.B. (ed), Chapman and Hall, London, UK, 1-22, 1993.

Tanaka *et al.*, *Plant Cell Physiol.* 36: 1023-1031, 1995.

- Tanaka *et al*, *Plant Cell Physiol.* 37: 711-716, 1996.
- Thompson *et al*, *Nucleic Acids Research* 22: 4673-4680, 1994.
- Turpen and Griffith, *BioTechniques* 4: 11-15, 1986.
- van Hoof and Green, *Plant Molecular Biology*, 35: 383-387, 1997
- Winkel-Shirley, *Plant Physiol.* 126: 485-493, 2001a.
- Winkel-Shirley, *Plant Physiol.* 127: 1399-1404, 2001b.
- Yonekura-Sakakibara *et al*, *Plant Cell Physiol.* 41: 495-502, 2000.

## THE CLAIMS DEFINING THE INVENTION ARE AS FOLLOWS:

1. An isolated nucleic acid molecule comprising a sequence of nucleotides encoding or complementary to a sequence encoding a flavonoid 3', 5' hydroxylase (F3'5'H) or a polypeptide having F3'5'H activity wherein expression of said nucleic acid molecule in a rose petal tissue results in detectable levels of *delphinidin* or *delphinidin-based molecules* as measured by a chromatographic technique.
2. An isolated nucleic acid molecule comprising a sequence of nucleotides encoding or complementary to a sequence encoding a F3'5'H or a polypeptide having F3'5'H activity wherein expression of said nucleic acid molecule in a rose petal tissue results in a sufficient level and length of transcript which is translated to said F3'5'H as determined by detectable levels of *delphinidin* or *delphinidin-based molecules* as measured by a chromatographic technique.
3. The isolated nucleic acid molecule of claim 1 or 2 wherein expression of said nucleic acid molecule in said rose petal results in a visually detectable colour change.
4. The isolated nucleic acid molecule of any one of claims 1 to 3, wherein the nucleic acid molecule is derived from a plant selected from the list comprising a *Viola* spp., *Salvia* spp., *Solysa* spp., *Lavandula* spp. and *Kennedia* spp.
5. The isolated nucleic acid molecule of claim 4, wherein the nucleic acid molecule is derived from a *Viola* spp. plant.
6. The isolated nucleic acid molecule of claim 5, wherein the nucleic acid molecule is derived from the *Viola* spp., cultivar Black Pansy.

7. The isolated nucleic acid molecule of any one of claims 5 or 6, wherein the nucleotide sequence encodes a R3'5'H comprising an amino acid sequence selected from SEQ ID NO:10, SEQ ID NO:12, an amino acid sequence having at least about 40% similarity to SEQ ID NO:10 and an amino acid sequence having at least about 40% similarity to SEQ ID NO:12.

8. The isolated nucleic acid molecule of claim 7, comprising a nucleotide sequence selected from SEQ ID NO:9, SEQ ID NO:11, a nucleotide sequence having at least about 40% identity to SEQ ID NO:9, a nucleotide sequence capable of hybridizing to SEQ ID NO:9 least about 40% identity to SEQ ID NO:11, a nucleotide sequence capable of hybridizing to SEQ ID NO:11 or its complement under low stringency conditions and a nucleotide sequence capable of hybridizing to SEQ ID NO:11 or its complement under low stringency conditions.

9. The isolated nucleic acid molecule of claim 8, comprising the nucleotide sequence set forth in SEQ ID NO:9.

10. The isolated nucleic acid molecule of claim 8, comprising the nucleotide sequence set forth in SEQ ID NO:11.

11. The isolated nucleic acid molecule of claim 4, wherein the nucleic acid molecule is derived from *Salvia spp.*

12. The isolated nucleic acid molecule of claim 11, wherein the nucleotide sequence encodes a R3'5'H comprising an amino acid sequence selected from SEQ ID NO:14, SEQ ID NO:16, an amino acid sequence having at least about 40% similarity to SEQ ID NO:14 and an amino acid sequence having at least about 40% similarity to SEQ ID NO:16.

- 136 -

13. The isolated nucleic acid molecule of claim 12, comprising a nucleotide sequence selected from SEQ ID NO:13, SEQ ID NO:15, a nucleotide sequence having at least about 40% identity to SEQ ID NO:13, a nucleotide sequence having at least about 40% identity to SEQ ID NO:15, a nucleotide sequence capable of hybridizing to SEQ ID NO:13 or its complement under low stringency conditions and a nucleotide sequence capable of hybridizing to SEQ ID NO:15 or its complement under low stringency conditions.

14. The isolated nucleic acid molecule of claim 13, comprising the nucleotide sequence set forth in SEQ ID NO:13.

15. The isolated nucleic acid molecule of claim 13, comprising the nucleotide sequence set forth in SEQ ID NO:15.

16. The isolated nucleic acid molecule of claim 4, wherein the nucleic acid molecule is derived from *Solha* spp.

17. The isolated nucleic acid molecule of claim 16, wherein the nucleotide sequence encodes a F3'5'H comprising an amino acid sequence selected from SEQ ID NO:18 and an amino acid sequence having at least about 40% similarity to SEQ ID NO:18.

18. The isolated nucleic acid molecule of claim 17, comprising a nucleotide sequence selected from SEQ ID NO:17, a nucleotide sequence having at least about 40% identity to SEQ ID NO:17 and a nucleotide sequence capable of hybridizing to SEQ ID NO:17 or its complement under low stringency conditions.

19. The isolated nucleic acid molecule of claim 18, comprising the nucleotide sequence set forth in SEQ ID NO:17.

20. The isolated nucleic acid molecule of claim 4, wherein the nucleic acid molecule is derived from *Kennedia* spp.

21. The isolated nucleic acid molecule of claim 20, wherein the nucleotide sequence encodes a R<sup>3</sup>'5'H comprising an amino acid sequence selected from SEQ ID NO:27 and an amino acid sequence having at least about 40% similarity to SEQ ID NO:27.

22. The isolated nucleic acid molecule of claim 21, comprising a nucleotide sequence selected from SEQ ID NO:26, a nucleotide sequence having at least about 40% identity to SEQ ID NO:26 and a nucleotide sequence capable of hybridizing to SEQ ID NO:26 or its complement under low stringency conditions.

23. The isolated nucleic acid molecule of claim 22, comprising the nucleotide sequence set forth in SEQ ID NO:26.

24. The isolated nucleic acid molecule of claim 4, wherein the nucleic acid molecule is derived from *Lavanthula* spp.

25. The isolated nucleic acid molecule of claim 24, wherein the nucleotide sequence encodes a R<sup>3</sup>'5'H comprising an amino acid sequence selected from SEQ ID NO:32 and an amino acid sequence having at least about 40% similarity to SEQ ID NO:32.

26. The isolated nucleic acid molecule of claim 25, comprising a nucleotide sequence selected from SEQ ID NO:31, a nucleotide sequence having at least about 40% identity to SEQ ID NO:31 and a nucleotide sequence capable of hybridizing to SEQ ID NO:31 or its complement under low stringency conditions.

27. The isolated nucleic acid molecule of claim 26, comprising the nucleotide sequence set forth in SEQ ID NO:31.

- 138 -

28. The isolated nucleic acid molecule of any one of claims 1- to 4, wherein the nucleotide sequence comprises an overall percentage of less than or equal to 54% of the nucleotides

- (i) A, or
- (ii) T, or
- (iii) A and T

in the third nucleotide position of each codon.

29. A construct comprising a sequence of nucleotides comprising:

- (i) a promoter which is operable in rose petal tissue and wherein said promoter is operably linked to,
- (ii) a nucleic acid molecule encoding F3'5'H, or complementary to a sequence encoding a F3'5'H or a polypeptide having F3'5'H activity wherein expression of said nucleic acid molecule in a rose petal tissue results in detectable levels of *delphinidin* or *delphinidin-based molecules* as measured by a chromatographic technique and wherein said nucleic acid molecule is derived from a plant selected from the group consisting of a *Viola spp.*, *Salvia spp.*, *Solhya spp.*, *Lavandula spp.* and *Kennedia spp.*

30. A construct comprising a sequence of nucleotides comprising:

- (i) a promoter which is operable in rose petal tissue and wherein said promoter is operably linked to,
- (ii) a nucleic acid molecule encoding F3'5'H, or complementary to a sequence encoding a F3'5'H or a polypeptide having F3'5'H activity wherein expression of said nucleic acid molecule in a rose petal tissue results in a sufficient level and length of transcript which is translated to said F3'5'H as determined by detectable levels of *delphinidin* or *delphinidin-based molecules* as measured by a chromatographic technique.

31. The construct of claim 29 or 30, wherein expression of said construct in said rose petal results in a visually detectable colour change.

32. The construct of any one of claims 29 to 31, wherein said promoter is selected from the group consisting of rose CHS, chrysanthemum CHS and CaMV 35S.

33. A construct of any one of claims 29 to 31 wherein said promoter comprises SEQ ID NO:5, or a functional equivalent thereof.

34. A construct of any one of claims 29 to 31 wherein said promoter comprises SEQ ID NO:30, or a functional equivalent thereof.

35. The construct of any one of claims 29 to 34, wherein the nucleic acid molecule is derived from a *Viola* spp.

36. The isolated nucleic acid molecule of claim 35, wherein the nucleotide sequence encodes R<sup>3</sup>5'H comprising an amino acid sequence selected from SEQ ID NO:10, SEQ ID NO:12, an amino acid sequence having at least about 40% similarity to SEQ ID NO:10, SEQ ID NO:12 and an amino acid sequence having at least about 40% similarity to SEQ ID NO:12.

37. The isolated nucleic acid molecule of claim 36, comprising a nucleotide sequence selected from SEQ ID NO:9, SEQ ID NO:11, a nucleotide sequence having at least about 40% identity to SEQ ID NO:9, a nucleotide sequence having at least about 40% identity to SEQ ID NO:11, a nucleotide sequence capable of hybridizing to SEQ ID NO:9 or its complement under low stringent conditions and a nucleotide sequence capable of hybridizing to SEQ ID NO:11 or its complement under low stringent conditions.

38. The isolated nucleic acid molecule of claim 37, wherein the gene comprises the nucleotide sequence set forth in SEQ ID NO:9.



- 140 -

39. The isolated nucleic acid molecule of claim 37, wherein the gene comprises the nucleotide sequence set forth in SEQ ID NO:11.

40. The isolated nucleic acid molecule of any one of claims 29 to 34, wherein the nucleic acid molecule is derived from *Salvia spp.*

41. The isolated nucleic acid molecule of claim 40, wherein the gene comprises a nucleotide sequence encoding F3'5'H comprising an amino acid sequence selected from SEQ ID NO:14, SEQ ID NO:16, an amino acid sequence having at least about 40% similarity to SEQ ID NO:14 and an amino acid sequence having at least about 40% similarity to SEQ ID NO:16.

42. The isolated nucleic acid molecule of claim 41, wherein the gene comprises a nucleotide sequence selected from SEQ ID NO:13, SEQ ID NO:15, a nucleotide sequence having at least about 40% identity to SEQ ID NO:13, a nucleic sequence having at least about 40% identity to SEQ ID NO:15, a nucleotide sequence capable of hybridizing to SEQ ID NO:13 or its complements under low stringent conditions and a nucleotide sequence capable of hybridizing to SEQ ID NO:15 or is complement under low stringent conditions.

43. The isolated nucleic acid molecule of claim 42, wherein the gene comprises the nucleotide sequence set forth in SEQ ID NO:13.

44. The isolated nucleic acid molecule of claim 42, wherein the gene comprises the nucleotide sequence set forth in SEQ ID NO:15.

45. The isolated nucleic acid molecule of any one of claims 29 to 34, wherein the gene is derived from *Salvia spp.*

- WO 2004/020637 PCT/AU2003/001111
46. The isolated nucleic acid molecule of claim 45, wherein the gene encodes a P3'5'H comprising an amino acid sequence selected from SEQ ID NO:18 and an amino acid sequence having at least about 40% similarity to SEQ ID NO:18.
47. The isolated nucleic acid molecule of claim 46, wherein the gene comprises a nucleotide sequence selected from SEQ ID NO:17, a nucleotide sequence having at least about 40% identity to SEQ ID NO:17 and a nucleotide sequence capable of hybridizing to SEQ ID NO:17 or its complements under low stringent.
48. The isolated nucleic acid molecule of claim 47, wherein the gene comprises the nucleotide sequence set forth in SEQ ID NO:17.
49. The isolated nucleic acid molecule of any one of claims 29 to 34, wherein the gene is derived from *Kennedia spp.*
50. The isolated nucleic acid molecule of claim 49, wherein the gene encodes P3'5'H comprising an amino acid sequence selected from SEQ ID NO:27 and an amino acid sequence having at least about 40% similarity to SEQ ID NO:27.
51. The isolated nucleic acid molecule of claim 50, wherein the gene comprises a nucleotide sequence selected from SEQ ID NO:26, a nucleotide sequence having at least about 40% identity to SEQ ID NO:26 and a nucleotide sequence capable of hybridizing to SEQ ID NO:26 or its complements under low stringent.
52. The isolated nucleic acid molecule of claim 51, wherein the gene comprises the nucleotide sequence set forth in SEQ ID NO:26.
53. The isolated nucleic acid molecule of any one of claims 29 to 34, wherein the gene is derived from *Lavandula spp.*

- 142 -

54. The isolated nucleic acid molecule of claim 53, wherein the gene encodes  $F3'5'H$  comprising an amino acid sequence selected from SEQ ID NO:32 and an amino acid sequence having at least about 40% similarity to SEQ ID NO:32.

55. The isolated nucleic acid molecule of claim 54, wherein the gene comprises a nucleotide sequence selected from SEQ ID NO:31 and a nucleotide sequence having at least about 40% identity to SEQ ID NO:31, a nucleotide sequence capable of hybridizing to SEQ ID NO:31 or its complements under low stringent conditions.

56. The isolated nucleic acid molecule of claim 55, wherein the gene comprises the nucleotide sequence set forth in SEQ ID NO:31.

57. A method for producing a transgenic flowering plant capable of synthesizing a  $F3'5'H$ , said method comprising stably transforming a cell of a suitable plant with a nucleic acid sequence as defined in any one of claims 1 to 28, under conditions permitting the eventual expression of said nucleic acid sequence, regenerating a transgenic plant from the cell and growing said transgenic plant for a time and under conditions sufficient to permit the expression of the nucleic acid sequence.

58. A method for producing a transgenic plant with reduced indigenous or existing  $F3'5'H$  activity, said method comprising stably transforming a cell of a suitable plant with a nucleic acid molecule as defined in any one of claims 1 to 28, regenerating a transgenic plant from the cell and where necessary growing said transgenic plant under conditions sufficient to permit the expression of the nucleic acid.

59. A method for producing a genetically modified plant with reduced indigenous or existing  $F3'5'H$  activity, said method comprising altering the  $F3'5'H$  gene through modification of the indigenous sequences via homologous recombination from an appropriately altered  $F3'5'H$  gene as defined in any one of claims 1 to 28, or a derivative or part thereof introduced into the plant cell, and regenerating the genetically modified plant from the cell.

64. A genetically modified plant or part thereof or cells therefrom comprising an isolated nucleic acid molecule of any one of claims 1 to 28 or comprising a reduced level of expression of a nucleic acid molecule of any one of claims 1 to 28.

63. A genetically modified plant or part thereof or cells therefrom comprising an isolated nucleic acid molecule of any one of claims 1 to 28.

62. A method for producing a transgenic plant capable of expressing a recombinant gene encoding F3'5'H as defined in any one of claims 1 to 28, or part thereof or which carries a nucleic acid sequence which is substantially complementary to all or a part of an mRNA molecule encoding said F3'5'H, said method comprising stably transforming a cell of a suitable plant with the isolated nucleic acid molecule comprising a sequence of nucleotides encoding, or complementary to a sequence encoding F3'5'H, where necessary under conditions permitting the eventual expression of said isolated nucleic acid molecule, and regenerating a transgenic plant from the cell.

61. A method for producing a flowering plant exhibiting altered inflorescence properties, said method comprising alteration of a F3'5'H gene as defined in any one of claims 1 to 28, through modification of the indigenous sequences *via* homologous recombination from an appropriately altered F3'5'H gene or derivative or part thereof introduced into the plant cell, and regenerating the genetically modified plant from the cell.

60. A method for producing a transgenic flowering plant exhibiting altered inflorescence properties, said method comprising stably transforming a cell of a suitable plant with a nucleic acid sequence as defined in any one of claims 1 to 28, regenerating a transgenic plant from the cell and growing said transgenic plant for a time and under conditions sufficient to permit the expression of the nucleic acid sequence.

- 144 -

65. A genetically modified plant or part thereof or cells therefrom comprising an isolated nucleic acid molecule of any one of claims 1 to 28 or comprising an increased level of expression of a nucleic acid molecule of any one of claims 1 to 28.

66. The genetically modified plant or part thereof or cells therefrom any one of claims 63 to 65, wherein the plant part is selected from sepal, bract, petiole, peduncle, ovaries, anthers, flowers, fruits, nuts, roots, stems, leaves, seeds.

67. The genetically modified plant or part thereof or cells therefrom of any one of claims 63 to 66, wherein the plant is a horticultural species, agricultural species or ornamental species.

68. Use of an isolated nucleic acid molecule as defined in any one of claims 1 to 28, in the manufacture of a genetic construct capable of expressing F3'5'H or down-regulating an indigenous F3'5'H enzyme in a plant.

69. A gene silencing construct comprising an isolated nucleic acid molecule as defined in any one of claims 1 to 28 or a complex thereof.

70. The genetically modified plant or part thereof or cells therefrom of 63 to 66, wherein the plant is selected from a rose, carnation, lisianthus, petunia, lily, pansy, gerbera, chrysanthemum, geranium, *Torenia*, *Begonia*, *Cyclamen*, *Nierembergia*, *Catharanthus*, *Pelargonium*, orchid, grape, apple, *Euphorbia* or *Fuchsia*.

71. An extract from a genetically modified plant or part thereof or cells therefrom from any one of claims 63 to 67 and 70.

72. The extract of claim 71, wherein the extract is a flavouring or food additive or health product or beverage or juice or colouring.

73. The method of any one of claims 57 to 62 wherein the genetically modified plant or part thereof or cells therefrom exhibit altered fluorescence.
74. An isolated recombinant R3'5'H or peptide having R3'5'H activity encoded by a nucleic acid molecule as defined in any one of claims 1 to 28.
75. The isolated recombinant R3'5'H or peptide having R3'5'H activity of claim 74, wherein the recombinant R3'5'H or peptide having R3'5'H activity is a fusion molecule comprising two or more heterologous amino acid sequences.
76. An isolated recombinant R3'5'H or peptide having R3'5'H activity nucleic acid molecule of any one of claims 1 to 28 comprising a fusion of two or more heterologous nucleotide sequences.
77. A prokaryotic organism carrying a genetic sequence encoding a R3'5'H molecule according to any one of claims 1 to 28 extrachromasomally in plasmid form.
78. A eukaryotic organism carrying a genetic sequence encoding a R3'5'H molecule according to any one of claims 1 to 28 extrachromasomally in plasmid form.
79. The use of a nucleic acid molecule of any one of claims 1 to 28 in the manufacture of a genetically modified plant or part thereof or cells therefrom.
80. The genetically modified plant or part thereof or cells therefrom of claim 79, wherein the genetically modified plant or part thereof or cells therefrom exhibit altered flowers or inflorescence.
81. The use of a nucleic acid sequence as defined in any one of claims 1 to 28 in the manufacture of a genetic construct capable of expressing R3'5'H or down-regulating an indigenous R3'5'H enzyme in a plant.

- 146 -

82. An isolated nucleic acid molecule comprising a sequence of nucleotides encoding or complementary to a sequence encoding a F3'5'H or a polypeptide having F3'5'H activity, wherein said nucleic acid molecule is derived from butterfly pea.

83. The isolated nucleic acid molecule of claim 81, wherein the nucleotide sequence encodes a F3'5'H comprising an amino acid sequence selecting for SEQ ID NO:21 and an amino acid sequence having at least about 40% similarity to SEQ ID NO:21.

84. The isolated nucleic acid molecule of claim 83, comprising a nucleotide sequence selected from SEQ ID NO:20, a nucleotide sequence having at least about 40% identity to SEQ ID NO:20 and a nucleotide sequence capable of hybridizing to SEQ ID NO:20 or its complement under low stringency conditions.

85. The isolated nucleic acid molecule of claim 84, comprising the nucleotide sequence set forth in SEQ ID NO:20.

86. The isolated nucleic acid molecule of any one of claims 1 to 4, wherein the nucleotide sequence comprises an overall percentage of less than or equal to 55% of the nucleotides

- (i) A, or
- (ii) T, or
- (iii) A and T.

87. An isolated nucleic acid molecule comprising SEQ ID NO:5 or a functional equivalent thereof.

88. An isolated nucleic acid molecule comprising SEQ ID NO:30 or a functional equivalent thereof.

- 147 -

89. An isolated nucleic acid molecule which has been modified so as to comprise a sequence of nucleotides encoding or complementary to a sequence encoding R3'S'H or a polypeptide having R3'S'H activity wherein expression of said nucleic acid molecule in a rose petal tissue results in detectable levels of *delphinidin* or *delphinidin-based molecules* as measured by a chromatographic technique wherein the nucleic acid molecule is derived from a plant selected from the list comprising *Petunia spp.*, *Crenata spp.* and *Clitoria spp.*

90. An isolated nucleic acid molecule which has been modified so as to comprise a sequence of nucleotides encoding or complementary to a sequence encoding R3'S'H or a polypeptide having R3'S'H activity wherein expression of said nucleic acid molecule in a rose petal tissue results in a sufficient level and length of transcript which is translated to said R3'S'H as determined by detectable levels of *delphinidin* or *delphinidin-based molecules* as measured by a chromatographic technique wherein the nucleic acid molecule is derived from a plant selected from the list comprising *Petunia*, *geranium* and butterfly pea.

91. The use of a nucleic acid sequence as defined in claim 89 or 90 in the manufacture of a genetic construct capable of expressing R3'S'H or down-regulating an indigenous R3'S'H enzyme in a plant.

92. The use of a nucleic acid sequence as defined in claim 89 or 90 in the manufacture of a genetically modified plant or part thereof or cells therefrom.






Figure 1a







Figure 1b



Replicon: pBluescript SK (+) vector 2.95kb


Insert: ~1.8kb petunia F3'5'H *petHf1* cDNA  
homologs from *P. hybrida* cv. OGB

Figure 2



Replicon: pBluescript SK (+) vector 2.95kb  
 Insert: ~1.8kb petunia F3'5'H petH2 cDNA  
 from *P. hybrida* cv. OGB


Figure 3



Replicon: ~2.7kb *EcoRI* (blunted) pUC19 vector

Insert: ~1.6kb *BspHI* (blunted)/*FspI* fragment containing petunia F3'5'H *petHf1* cDNA from pCGP601

Figure 4



Replicon: ~18.7kb SmaI pWTT2132 vector  
 Insert: ~3.5kb PstI (blunted) fragment  
 containing *AmCHS* 5': *petH1*: *petD8* 3' gene  
 from pCGP485

Figure 5





Figure 6

①

②






Replicon: 2.95kb (BamHI/XbaI) blunted vector fragment of pBluescript II KS (+).


Insert: ~3.5kb PstI (blunted) fragment containing *AmCHS* 5': *petHf1*: *petD8* 3' gene from pCGP483

Figure 7



Replicon: ~18.7kb *SmaI* pWTT2132 vector  
 Insert: ~3.9kb *PstI* (blunted) fragment  
 containing *Mac*: *petH1*: *mas 3'* gene from  
 pCGP628


Figure 8



Replicon: ~18.7kb SmaI/PstI pWTT2132 vector

Insert: ~5.3kb XbaI (blunted)/PstI fragment containing *petD8* 5': *petHf1*: *petD8* 3' gene from pCGP1107


Figure 9



Replicon: ~18.7kb PstI (blunted)/KpnI  
pWTT2132 vector

Insert: ~4.35kb SacI (blunted)/KpnI fragment  
containing *shortFLS 5': petHf1: petFLS 3'* gene  
from pCGP497


Figure 10



Replicon: ~18.7kb PstI/BamHI pWTT2132 vector


Insert: ~3kb PstI/BamHI fragment containing *petRT5'*: *pethHf1*: *nos 3'* gene from pCGP846

Figure 11



Replicon: ~18.7kb SalI pWTT2132 vector  
 Insert: ~4.9kb XhoI fragment containing  
 mas/35S:petHf1:ocs 3' gene from pCGP1619


Figure 12



Replicon: ~18.7kb SmaI pWTT2132 vector

Insert: ~2.6kb (PstI/EcoRI) blunted fragment containing *CaMV* 35S: *petHf1*: *ocs* 3' gene from pCGP1636

Figure 13




Replicon: ~18.7kb BamHI pVTT2132  
vector

Insert: ~4.9kb BglII fragment from  
containing *RoseCHS 5'*:*petH1*: *nos 3'* gene  
from pCGP200

Figure 14






Replicon: ~18.4kb Asp718 (blunted)  
pCGP1988 vector

Insert: ~3.7kb (Asp718/XbaI) blunted  
fragment containing *CaMV 35S*: *petHf2*:  
*ocs 3'* gene from pCGP2109


Figure 15



Replicon: ~18.4kb Sall (blunted)/PstI vector  
 fragment from pWTT2132

Insert: ~66bp EcoRI (blunted)/PstI fragment  
 containing multi-cloning site from pNEB193


Figure 16



Replicon: ~3.3 kb HincII/XhoI vector fragment from pCGP2000 (containing *CaMV* 35S promoter fragment in pBluescript SK)


Insert: ~1.6kb EcoRI (blunted)/ XhoI ocs 3' fragment from pKIWI101

Figure 17.



Replicon: ~15kb HindIII/PstI pCGN1548 vector  
 Insert: ~5.3kb HindIII/PstI fragment containing  
*petD8* 5': *GUS*: *petD8* 3' gene from pCGP1106


Figure 18



Replicon: ~11.8kb BamHI (GA-filled)/SacI  
pBIN19 vector


Insert: ~6.7kb XhoI (TC-filled)/SacI  
fragment containing *longpetFLS* 5': *GUS*:  
*petFLS* 3' gene from pCGP496

Figure 19



Replicon: ~18.7kb PstI/BamHI pVST132  
vector  
Insert: ~4.6kb PstI/BglII fragment  
containing *ChrysoCHS 5'*: *GUS*: *nos 3'*  
gene from pCGP1622


Figure 20



Replicon: ~18.7kb PstI pWTT2132 vector

Insert: ~5.4kb PstI fragment containing  
*petRT 5'*; *GUS*; *petRT 3'* gene from  
pCGP1628


Figure 21



Replicon: ~18.7kb BamHI pWTT2132  
vector  
Insert: ~5kb BglII fragment containing  
RoseCHS 5': GUS: nos 3' gene from  
pCGP197

Figure 22





Replicon: ~18.7kb Asp718 (blunted)  
pWTT2132 vector

Insert: ~3.8kb (EagI/PstI) blunted  
fragment containing *AmCHS* 5': *GUS*:  
*petD8* 3' gene from pCGP1952

Figure 23






Figure 24



Replicon: pBluescript SK II (+) vector 2.95kb

Insert: ~1.6kb *pansy* F3'5'H BP#18 cDNA from *Viola* spp. cv. Black Pansy


Figure 25



Replicon: pBluescript SK II (+) vector  
2.95kb

Insert: ~1.6kb pansy F3'5'H BP#40 cDNA  
from *Viola* spp. cv. Black Pansy


Figure 26



Replicon: ~18.7kb Asp718 (blunted)  
pWTT2132 vector

Insert: ~3.8kb NotI (blunted)/ EcoRV  
fragment containing *AmCHS* 5': *BP#18*:  
*petD8* 3' gene from pCGP1970


Figure 27



Replicon: ~18.7kb *Asp718* (blunted)  
 pWTT2132 vector

Insert: ~3.8kb *NotI* (blunted) / *EcoRV*  
 fragment containing *AmCHS 5'*: BP#40:  
*petD8 3'* gene from pCGP1971


Figure 28



Replicon: ~18.7kb Asp718 (blunted)  
pWTT2132 vector

Insert: ~3.6kb (XhoI /XbaI) blunted  
fragment containing CaMV 35S: BP#18:  
ocs 3' gene from pCGP1965

Figure 29




Replicon: ~18.7kb Asp718 (blunted)  
pWTT2132 vector

Insert: ~3.6kb (XhoI/XbaI) blunted  
fragment containing CaMV 35S: BP#40:  
ocs 3' gene from pCGP1966

Figure 30






Replicon: pBluescript SK II (+) vector 2.95kb

Insert: ~1.6kb F3'5'H Sal#2 cDNA from *Salvia spp.*


Figure 31



Replicon: pBluescript SK II (+) vector  
2.95kb

Insert: ~1.6kb F3'5'H SalI#47 cDNA from  
*Salvia* spp.


Figure 32



Replicon: ~18.4kb *Asp718* (blunted)  
pCGP1988 vector

Insert: ~3.6kb *NotI* (blunted)/ *EcoRV*  
fragment containing *AmCHS* 5': *Sal*#2:  
*petD8* 3' gene from pCGP2116


Figure 33



Replicon: ~18.4kb Asp718 (blunted)  
pCGP1988 vector

Insert: ~3.6kb NotI (blunted)/EcoRV  
fragment containing AmCHS 5'  
SalI#47: petD8 3' gene from  
pCGP2117


Figure 34



Replicon: ~18.4kb Asp718 (blunted)  
pCGP1988 vector

Insert: ~3.6kb (XhoI/XbaI) blunted  
fragment containing CaMV 35S:  
Sal#2: ocs 3' gene from pCGP2112


Figure 35



Replicon: ~18.4kb Asp718 (blunted)  
pCGP1988 vector

Insert: ~3.6kb (XhoI/XbaI) blunted  
fragment containing CaMV 35S: SaII#47:  
ocs 3' gene from pCGP2111


Figure 36



Replicon: pBluescript SK II (+) vector  
2.95kb

Insert: ~1.7kb *F3'5'H SolI#5* cDNA  
from *SolIya spp.*

Figure 37




Replicon: ~18.4kb Asp718 (blunted)  
 PCGP1988 vector

Insert: ~3.5kb NotI (blunted)/EcoRV  
 fragment containing *AmCHS* 5': *Sol/#5*:  
*petD8* 3' gene from PCGP2128

Figure 38






Replicon: ~18.4kb Asp718 (blunted)  
pCGP1988 vector


Insert: ~3.6kb (Asp718/XbaI) blunted  
fragment containing *CaMV* 35S:  
*Soll*#5: *ocs* 3' gene from pCGP2129

Figure 39



Replicon: pBluescript SK II (+) vector 2.95kb  
 Insert: ~1.8kb F3'5'H Kenn#31 cDNA from  
*Kennedia* spp.


Figure 40



Replicon: ~18.4kb *Asp718* (blunted)  
pCGP1988 vector.

Insert: ~3.7kb (*NotI*/ *EcoRI*) blunted  
fragment containing *AmCHS* 5':  
Kenn#31: *petD8* 3' gene from  
pCGP2242


Figure 41



Replicon: ~18.4kb Asp718 (blunted)  
pCGP1988 vector

Insert: ~3.6kb (XhoI/NotI) blunted fragment  
containing CaMV 35S: Kenn#31: ocs 3' gene  
from pCGP2236


Figure 42



Replicon: pBHF2 BamHI/PstI 4.5kb vector + partial *BpeaHF2* insert (backbone = pBluescript SK II (+) vector)

Insert: ~200bp BamHI/PstI fragment from PCR using pBHF2 as template (5' fragment of butterfly pea *F3'5'H* cDNA (*BpeaHF2*) from *Clitoria ternatea* including putative initiating codon (ATG))


Figure 43



Replicon: ~18.4kb Asp718 (blunted)  
pCGP1988 vector

Insert: ~3.6kb NotI (blunted)/EcoRV  
fragment containing AmCHS 5': BpeaHF2:  
petD8 3' gene from pCGP2133


Figure 44



Replicon: ~12.8kb pBE2113-GUSs BamHI/Sall  
(pBI121 backbone)

Insert: ~1.7kb BamHI/XhoI fragment containing  
*Clitoria F3'5'H BpeaHF2* cDNA clone from  
pBHF2F

Figure 45




Replicon: ~18.4kb Asp718 (blunted)  
pcGP1988 vector

Insert: ~3.6kb (XhoI/XbaI) blunted  
fragment containing CaMV 35S:

BpeaHF2: ocs 3' gene from pcGP2132

Figure 46






Replicon: pBluescript SK II (+) vector 2.95kb

Insert: ~1.7kb F3'5'H Gen#48 cDNA from *Gentiana triflora*


Figure 47



Replicon: ~18.7kb Asp718 (blunted)  
pWTT2132 vector

Insert: ~3.6kb NotI (blunted)/ EcoRV  
fragment containing AmCHS 5': Gen#48:  
petD8 3' gene from pCGP1496


Figure 48



Replicon: ~12.8kb pBE2113-GUSs BamHI/Sall  
(pBI121 backbone)

Insert: ~1.8kb BamHI/XhoI fragment containing  
*gentian F3'5'H* (*Genn#48*) cDNA clone from  
pG48


Figure 49



Replicon: ~18.7kb Asp718 (blunted)  
pWTT2132 vector

Insert: ~3.6kb (XhoI/XbaI) blunted fragment  
containing CaMV 35S: Gen#48: ocs 3' gene  
from pCGP1981

Figure 50



Replicon: pBluescript SK II (+) vector 2.95kb

Insert: ~1.8kb *lavender F3'5'H LBG cDNA*  
from *Lavendula nil*

Figure 51



- 1 -

## SEQUENCE LISTING

<110> International Flower Developments Pty. Ltd.  
 Brugliera, Filippa (US only)  
 Tanaka, Yoshikazu (US only)  
 Mason, John (US only)

<120> Genetic Sequences and Uses Therefor

<130> 12322720/EJH

<150> AU 2002951088

<151> 2002-08-30

<150> AU 2002952835

<151> 2002-09-16

<160> 32

<170> PatentIn version 3.1

<210> 1

<211> 1812

<212> DNA

<213> petunia

<400> 1

ctttctacta gctacttcgt tatatatatg taaaattgtg actttgaaaa tcattttaa	60
tatcataagg ttcatTTTTat ctTgatcaaa atatttactt cggccatata cgttttcc	120
tagtcatgat gctacttact gagcttggtg cagcaacttc aatctttcta atagcacaca	180
taatcatttc aactcttatt tcaaaacta cgggcggca tctaccgccg gggcagag	240
ggtggccggt gatcggagca ctccacttt taggagccat gccacatgtt tccctagcta	300
aastggcaaa aaatatgga gcaatcatgt atctcaaagt tggaacatgt ggcattgag	360
ttgctttctac cctgatgct gctaaagcat tcttgaaaac aattgatatc aacttctoca	420
atgctccacc taatgcagggt gccactcaat tagcttataa tgctcagac atggtttttg	480
cacattatgg accacgatgg aagttgctaa ggaanttaag caacttgcac atgctagggg	540

gaaagagccct agagagcttg gcaaatgtcc gtagcaatga gctagagagcac atgcataaat  
600 caatgcctga tctagagctcga gaggagccaga aggtgtgtgtgt ggcggagagatg ttagaatgtg  
720 caatggccga tatgagctcaga caagtgtatga taagcaaaag agtatgtga gataaagtg  
780 ttgaggtcaa tgaattcaa gacatgggtg tagagttaat gacataagca ggttatcca  
840 acatgtgtga ttttatccct tgtttagctt gtaggtatct acaggggtat gaaagagaa  
900 tgaacgtctt acataagag ttgtatgctt tatgtgcaaa gattgttgat gaacacaaag  
960 caactaccta tgaagcttag gggaaacag atcttcttga tgttgttatg gaaaatgggg  
1020 acatctcga agaggaaga ctcagtcaa caacatcaa agaatcttg ctgaattctg  
1080 taagagctgg tagggcaact tatcttagg caatagatg ggcacttgca gaatgtgtg  
1140 agaacctgc catcttgaaa aagagcaag cagaaatga tcaagtcat ggaagaaata  
1200 ggcgttact gaatcagat atcccaatc tcccttacct ccgagcaatt tgaagaaagaa  
1260 catctcgaaa acacccctc acacaccaa atcttctcag gactcagaa gaaocatga  
1320 tagtcatggt tcttatcaat caaaaaaa ctatgtcttag gtttaacaa tgggcaattg  
1380 gaagagatcc caagtgttg gaaatccaa tagagttaa tccogaaag tctgtgatg  
1440 gaagaaatcc caagtctgat cctcagaggga agatcttga atgtatcaca tctgtgtcgtg  
1500 gacgaagaa ttgtgcagga acaagacgg gaattgtaat ggttgaaat atattagga  
1560 cttgtgttca tctcatgtgat tggaaatca caagtgaat tcttgagttg aataggaag  
1620 aagcctcttg cttagctctg cagaaagctg tccctcttga agctatgtt actcaaggt  
1680 taacaattga tgttatga caatagctat agatgtgat tggctatga tggcagatg  
1740 tgtgtgtgt agcatgagat atcaaaaga gtacatcgaag cgcattgcat ggttcaat  
1800 tgaagccct taatatctt gcatcttca aatcaatga tctgtgttg tgggtcaaa  
1812 aaaaaaa aa

<210> 2  
<211> 506  
<212> ERI  
<213> Pctm1a  
<400> 2

Met Met Leu The Gly Ala The Ser Ile Phe Ile  
5 10 15



- 3 -

Ala His Ile Ile Ile Ser Thr Leu Ile Ser Lys Thr Thr Gly Arg His  
 20 25 30

Leu Pro Pro Gly Pro Arg Gly Trp Pro Val Ile Gly Ala Leu Pro Leu  
 35 40 45

Leu Gly Ala Met Pro His Val Ser Leu Ala Lys Met Ala Lys Lys Tyr  
 50 55 60

Gly Ala Ile Met Tyr Leu Lys Val Gly Thr Cys Gly Met Ala Val Ala  
 65 70 75 80

Ser Thr Pro Asp Ala Ala Lys Ala Phe Leu Lys Thr Leu Asp Ile Asn  
 85 90 95

Phe Ser Asn Arg Pro Pro Asn Ala Gly Ala Thr His Leu Ala Tyr Asn  
 100 105 110

Ala Gln Asp Met Val Phe Ala His Tyr Gly Pro Arg Trp Lys Leu Leu  
 115 120 125

Arg Lys Leu Ser Asn Leu His Met Leu Gly Gly Lys Ala Leu Glu Asn  
 130 135 140

Trp Ala Asn Val Arg Ala Asn Glu Leu Gly His Met Leu Lys Ser Met  
 145 150 155 160

Ser Asp Met Ser Arg Glu Gly Gln Arg Val Val Val Ala Glu Met Leu  
 165 170 175

Thr Phe Ala Met Ala Asn Met Ile Gly Gln Val Met Leu Ser Lys Arg  
 180 185 190

Val Phe Val Asp Lys Gly Val Glu Val Asn Glu Phe Lys Asp Met Val  
 195 200 205

Val Glu Leu Met Thr Ile Ala Gly Tyr Phe Asn Ile Gly Asp Phe Ile  
 210 215 220

Pro Cys Leu Ala Trp Met Asp Leu Gln Gly Ile Glu Lys Arg Met Lys  
 225 230 235 240

Arg Leu His Lys Lys Phe Asp Ala Leu Leu Thr Lys Met Phe Asp Glu 255  
 His Lys Ala Thr Thr Tyr Glu Arg Lys Gly Lys Pro Asp Phe Leu Asp 260  
 Val Val Met Glu Asn Gly Asp Asn Ser Glu Gly Arg Leu Ser Thr 275  
 Thr Asn Ile Lys Ala Leu Leu Asn Leu Phe Thr Ala Gly Thr Asp 290  
 Thr Ser Ser Ser Ala Ile Glu Trp Ala Leu Ala Glu Met Met Lys Asn 305  
 Pro Ala Ile Leu Lys Lys Ala Glu Ala Glu Met Asp Glu Val Ile Gly 325  
 Arg Asn Arg Arg Leu Leu Glu Ser Asp Ile Pro Asn Leu Pro Tyr Leu 340  
 Arg Ala Ile Cys Lys Glu Thr Phe Arg Lys His Pro Ser Thr Pro Leu 355  
 Asn Leu Pro Arg Ile Ser Asn Glu Pro Cys Ile Val Asp Gly Tyr Tyr 370  
 Ile Pro Lys Asn Thr Arg Leu Ser Val Asn Ile Trp Ala Ile Gly Arg 385  
 Asp pro Glu Val Trp Glu Asn Pro Leu Glu Phe Asn Pro Glu Arg Phe 405  
 Leu Ser Gly Arg Asn Ser Lys Ile Asp Pro Arg Gly Asn Asp Phe Glu 420  
 Leu Ile Pro Phe Gly Ala Gly Arg Arg Ile Cys Ala Gly Thr Arg Met 435  
 Gly Ile Val Met Val Glu Tyr Ile Leu Gly Thr Leu Val His Ser Phe 450

- 5 -

Asp Trp Lys Leu Pro Ser Glu Val Ile Glu Leu Asn Met Glu Glu Ala  
465 470 475 480

Phe Gly Leu Ala Leu Gln Lys Ala Val Pro Leu Glu Ala Met Val Thr  
485 490 495

Pro Arg Leu Gln Leu Asp Val Tyr Val Pro  
500 505

<210> 3  
<211> 1756  
<212> DNA  
<213> petunia

<400> 3  
ttgaatccag ctctatctgg cttagacaa tgggtctact tagtgagctt gctgcagcaa 60  
ccttaatctt tctaacaaca catatcttca ttcaactct tctttctata actaacggcc 120  
ggcgtctccc gccagggcca agagggtggc oggtgatcgg agcacttcca cttttaggag 180  
ccatgccaca tgtttcetta gctaaaatgg caaaaaata tggagcaatc atgtatctca 240  
aagttggaac gtgtggcatg gtagttgctt ctacccctga tgcgtctaaa gcgttcttga 300  
aaacacttga tctcaacttc tccaatcgtc cacctaattg aggtgccacc cacttagcct 360  
atggtgctca agacatggtt ttgacacatt atggaccaag atggaagttg ctaaggaaat 420  
taagcaactt acatctgcta ggggggaaag ccttagaana ttgggcaaat gtctgtgcca 480  
atgagctagg acacatgcta aaatcgatgt ttgatattgag cagagaaggg gagagagttg 540  
tgggtggcga gatgttgaca ttgccatgg cgaatatgat cggacaggtg atacttagca 600  
aaagagtatt tgbaaataaa ggtgttgagg taaatgaatt taaggaaatg gtggttagagt 660  
taatgacaac agcagggtat ttaacattg gtgattttat tcttctttta gotttggatgg 720  
atttacaagg gatagaaaaa ggaatgaaac gtttacataa gaagtttgat gctttattga 780  
caagatgtt tgatgaacac aaagcaacta gctatgaacg taaggggaaa ccagattttc 840  
ttgatttgtt tatggaaaat agggacaatt ctgaaggaga aaggctcagt acaaccaaca 900  
tcaaagcaat cttgctgaat ttgttcaacg ctggtacaga cacttcttct agtgcaatag 960  
aatgggcaat tgcagagatg atgaagaacc ctgccatttt aaagaaagca caaggagaaa 1020  
tggatcaagt cattggaaac aataggogtc tgcctgaatc ggatatccca aatctccctt 1080  
acctccgagc aatttgcaaa gaaacatttc gaaagcaccg ttctacacca ttaaatctcc 1140  
ctaggatctc gaacgaacca tgcattgtcg atggttatta cataccaaaa aacactaggc 1200

ttatgttttt catatgggca atgggaagat atccggagat ttgggaagaa caactagagt 1260  
ttatctctga aggttcttg agtgggaagaa actggagat ttgatctcga gggaacgagat 1320  
ttgaattgat acaatttggt gctgggaagaa gaatttggtc aggggaagaa atgggaatcg 1380  
taattgttga acatatttt gaacttttg tccattcatt ttgatggaaa ttacaaagt 1440  
aattattga gctaaatag gaagaact ttgattagc ttggcagaaa gctgtccctc 1500  
ttgaagctat gttactcca aggttcctta ttgatltta tgcacctta gcttgaaaaa 1560  
tgcttttag ttgtttcag ttgttggtag taattgttg ttgttttg ctatagaat 1620  
atcaaat gatagatc tgaaggogcg tgaaggogcg ggggttgct ttatagatga 1680  
gtaatatgtt agcttctt ttatctctg ttgttgtag aatcttgata ttcttcttg 1740  
gaaaaaaa aaaaaa 1756

<210> 4  
<211> 608  
<212> PRT  
<213> peptide  
<400> 4

Met Val Leu Leu Ser Glu Leu Ala Ala Thr Leu Ile Phe Leu Thr  
1 5 10 15

Thr His Ile Phe Ile Ser Thr Leu Ser Ile Thr Asn Gly Arg  
20 25 30

Leu Pro Pro Gly Pro Arg Gly Trp Pro Val Ile Gly Ala Leu Pro Leu  
35 40 45

Leu Gly Ala Met Pro His Val Ser Leu Ala Lys Met Ala Lys Lys Tyr  
50 55 60

Gly Ala Ile Met Tyr Leu Lys Val Gly Thr Cys Gly Met Val Val Ala  
65 70 75 80

Ser Thr Pro Asp Ala Lys Ala Phe Leu Lys Thr Leu Asp Leu Asn  
85 90 95

Phe Ser Asn Arg Pro Asn Ala Gly Ala Thr His Leu Ala Tyr Gly  
100 105 110

- 7 -

Ala Gln Asp Met Val Phe Ala His Tyr Gly Pro Arg Trp Lys Leu Leu  
 115 120 125

Arg Lys Leu Ser Asn Leu His Met Leu Gly Gly Lys Ala Leu Glu Asn  
 130 135 140

Trp Ala Asn Val Arg Ala Asn Glu Leu Gly His Met Leu Lys Ser Met  
 145 150 155 160

Phe Asp Met Ser Arg Glu Gly Glu Arg Val Val Val Ala Glu Met Leu  
 165 170 175

Thr Phe Ala Met Ala Asn Met Ile Gly Gln Val Ile Leu Ser Lys Arg  
 180 185 190

Val Phe Val Asn Lys Gly Val Glu Val Asn Glu Phe Lys Asp Met Val  
 195 200 205

Val Glu Leu Met Thr Thr Ala Gly Tyr Phe Asn Ile Gly Asp Phe Ile  
 210 215 220

Pro Cys Leu Ala Trp Met Asp Leu Gln Gly Ile Glu Lys Gly Met Lys  
 225 230 235 240

Arg Leu His Lys Lys Phe Asp Ala Leu Leu Thr Lys Met Phe Asp Glu  
 245 250 255

His Lys Ala Thr Ser Tyr Glu Arg Lys Gly Lys Pro Asp Phe Leu Asp  
 260 265 270

Cys Val Met Glu Asn Arg Asp Asn Ser Glu Gly Glu Arg Leu Ser Thr  
 275 280 285

Thr Asn Ile Lys Ala Leu Leu Leu Asn Leu Phe Thr Ala Gly Thr Asp  
 290 295 300

Thr Ser Ser Ser Ala Ile Glu Trp Ala Leu Ala Glu Met Met Lys Asn  
 305 310 315 320

Pro Ala Ile Leu Lys Lys Ala Gln Gly Glu Met Asp Gln Val Ile Gly  
 325 330 335

- 8 -

Asn Asn Arg Arg Leu Leu Glu Ser Asp Ile Pro Asn Leu Pro Tyr Leu 340 345 350

Arg Ala Ile Cys Lys Glu Thr Phe Arg Lys His Pro Ser Thr Pro Leu 355 360 365

Asn Leu Pro Arg Ile Ser Asn Glu Pro Cys Ile Val Asp Gly Tyr Tyr 370 375 380

Ile Pro Lys Asn Thr Arg Leu Ser Val Asn Ile Trp Ala Ile Gly Arg 385 390 395 400

Asp Pro Glu Val Trp Glu Asn Pro Leu Glu Phe Tyr Pro Glu Arg Phe 405 410 415

Leu Ser Gly Arg Asn Ser Lys Ile Asp Pro Arg Gly Asn Asp Phe Glu 420 425 430

Leu Ile Pro Phe Gly Ala Gly Arg Arg Ile Cys Ala Gly Thr Arg Met 435 440 445

Gly Ile Val Met Val Glu Tyr Ile Leu Gly Thr Leu Val His Ser Phe 450 455 460

Asp Trp Lys Leu Pro Ser Glu Val Ile Glu Leu Asn Met Glu Glu Ala 465 470 475 480

Phe Gly Leu Ala Leu Glu Lys Ala Val Pro Leu Glu Ala Met Val Thr 485 490 495

Pro Arg Leu Pro Ile Asp Val Tyr Ala Pro Leu Ala 500 505

<210> 5  
<211> 2934  
<212> DNA  
<213> rose

<400> 5  
aagcttcagc aagagttgaa gaataaggga cagagccatc catgtgctt gatgaatcga 60  
atggatatac aaatgtgana gattcaactg ctgatttacc cagatttct tcaataagtg 120  
aggagaatgt tgaagatct aatgatagay actctgttaa actagaagga attcattgtgc 180  
agcaagagtg tcaatgaggc agtgaagaa agaaacctga tggtaagagc ggtgagatg 240

- 9 -

cagttgatct ggctaatacat ggcattggctc gaactgattt ttgtcagata acagaagaga	300
ttgagaatgg agtagtcato actgagatga gcaacattgc caaccotgat aaaactgata	360
ttccaaaacgg ggtgcctcaa aatgagactg atgatggatt taataaacct caggatgatg	420
ctaatacaaa ggaagtgaca gaagagaatt ctgacagacg tgcgaaggaa gtgacagaag	480
agaattctga caaagatggt ttgaagaata tcttgaatt ctacagtgtt tottctgttg	540
tggattttga aattccagtg ttggatgtga aatttaettc tottgaagt tgcagtgcga	600
cttgttctct tgcagccctt ttgtctgaat cgcgggaatc aatgactgaa gcaccttgtg	660
tgaggcaaat tgatgatgtg ccccggttg gtgaggagtc tagettgatt ttggtggaag	720
atcgggagcc gtttggtoct actcctgatg gtaatttttc tgtggatatg gattactata	780
gtgtagcaga acctttgagc acatgggatg cgaatctgca gtgtgaaaca tcaaatagcc	840
atgagacttt tgcctgaagt ctatattgat agcttctgtg ttaataaact tgttagtctg	900
tacataaatt tgtctagaca agaattggtc gtgtactatc gtgtgttttt gcctgtgctt	960
agtactcatg aaccaattca gaaaaactg gctgcataat ttgaggagtc totgaattct	1020
tcaatgctca actggatgac atgtagggtg catatcaact cagggtattct totattcttt	1080
aaotttacgc atcttgacat tttgtatata acaaaatcag gtctattggg tgaaagtaat	1140
tggctagaat ggaagctct acggttttac cgcaggtaaa ttttcatagc tccacaagtg	1200
aattgaaaat gctcatagga tttatgtttg tctccact ctggcgaaga tgtttgttg	1260
ggagttaact caaacctacc accaaaactg aacctatct ccataattta taatacaaat	1320
ttgcgatcat ttgttcatec aattattgtg aactcggct accacccaa atatcggta	1380
cagacccaaa cgtattgtca caacaaatcg tgtctctcgc attaaacaca gctagaaga	1440
agagttgaac ccaaatctg agacccact acctatgtac gaagtcatga gttcagatca	1500
ccataggggt agaagtgaat tcatttgatc atctttaaag aaataaaaag aagagttgaa	1560
cccacaattg gctcttgtcc caaaaagaac taatagttca gtgcaccgac gtgtatttgc	1620
acogacataa atggattgtt agattatatt aaatacaact ttaggttatt aataaaaata	1680
ttaattataa atatcaaaag ttgagatcat cttataaatg ttgggtcagt tacacggtcg	1740
gtgcatagaa taatttcaa acctatataat agccttcatt ttctgattta gctcatggga	1800
catgattgct ataaataatt gtactcgtag aggcataact gtgtcttttt atacagttgt	1860
actgaagctc agaaaagttt atgaagggtg gaactgagaa gggcaaggca tttggtagtt	1920

- 10 -

gaggtatarg agagcatgaa cccatggat tgcagctaac accctctt ttccctct  
 1980  
 cccatgaaaa taaaacaaac tctctccac taagtctaac atcttctt atggcagctc  
 2040  
 ttgccttaac agtccatcta tatctataa ttctatcta atattgtcta ctctgtctac  
 2100  
 ctacagccc aaaaataaac tgaataagc cxaattgtag atatttctg tttctgtct  
 2160  
 tgttttctc ttttcttat gatttttta aaatttaaat gacttcaatt ttgtttctg  
 2220  
 tttttttt tatttttt tatagaaaa tgcgaaaac ttcatctac gttattgtag  
 2280  
 acaatttagc catataaac tatataatat tctcttcaa ttcgagtaaa tttaaaacgg  
 2340  
 tgaataata aatatgtac gttatctaa atgaataaaa ctcaatctaat atagttataa  
 2400  
 ctgaattcac atctaacgac atagattctc ttcatctagt cttaaccagt ttgaattatat  
 2460  
 agatttgat tatgtttatg tcttgtcca catttgtgtc tgaataaaaa tgtgcacagg  
 2520  
 aggtatgta ctgttgtctc atcaattcaa agtttgaaat aaagaaaaaa aaaaaagagc  
 2580  
 atatttctg cgttctgtct gtttaggtaaa acgaggaaac aaagcctac caaatcatgc  
 2640  
 gatttctgat cggcaaaaaa caacaaaaaa kaacgttaaac gatgcacagt ccaatttgcg  
 2700  
 ggggttcta acaagttant tgggcagggc cgttgatcccc cagctaacca cctctcgtt  
 2760  
 cccctctcaa actaatcttc catgtatat taacaaacct ttctcagac catctatctc  
 2820  
 taacatttc gcttgttat tgaacgcaa caaaaaatgc tcatctcaco ctgtctctc  
 2880  
 cccatttga tcttctctcg accctctcc gagatggga ccgaactcga attc  
 2934

<210> 6  
 <211> 24  
 <212> DNA  
 <213> Petunia  
 <400> 6  
 gtctcgaag aagatataa caat  
 24

<210> 7  
 <211> 20  
 <212> DNA  
 <213> Petunia  
 <400> 7  
 caagatcga ggaactgcatg  
 20

<210> 8  
 <211> 20  
 <212> DNA  
 <213> chrysanthemum



- 11 -

<400> 8  
gttaaggaag ccatgggtgt

20

<210> 9  
<211> 1648  
<212> DNA  
<213> viola

<400> 9  
agccaatatg gcaattccag teactgaact tgctgtcgcg gttatccttt tottgatcac 60  
tcgcttctta gttcgttctc ttttcaagaa accaaccgga ccgctcccgc cgggtccttc 120  
aggctggccc ttggtgggcg cgtccctctt cctaggcgcc atgctccacg tcccaactagc 180  
caacctcgct aaaaaatacg gtccgatcat gtacctaaaa atgggoacgt gcgacatggt 240  
ggtcgcgtcc aotcccgact cggctcgagc cttcctcaaa accctagacc tcaacttctc 300  
cgaccgcccg cccaacgcg gcgccaccca tttggcgtae ggccgcgagc acttggtctt 360  
cgcgaagtac ggtccaaagt ggaagacct aaganaattg agcaacctcc acatgctagg 420  
cgggaaggcg ctggaagatt gggctcacgt gagggttaac gagctaggcc acatgcttaa 480  
cgccatgtgc gaggcgagcc ggtgcgaga gcccggtgtg ctggccgaga tgcacacgta 540  
cgccatggcc aacatgatcg gtcagtgat actgagtcgg cgcgtgttcg tcaccaaagg 600  
gacagagtcg aacgagttca aagatatggt ggtcgagttg atgacttccg cggggtattt 660  
caacattggt gacttcatac cgtcgattgc ttggatgat ttgcaagga togagcgagg 720  
gatgaagaaa ttgcacacga aattcgatgt tttgttgacg aagatgatga aggagcacag 780  
agcgacgagt catgagcgcg aagggaatc ggatttctc gaogtctct tggagaatg 840  
cgagaataca aatggcgaga agcttaatgt taccacgctc aagctgtcc tcttgaactt 900  
attcacggcg ggtacggaca catottcaag cataatcgaa tgggcgttaa ccgaaatgat 960  
gaagaatccg acgatcttaa azaagaccca agaagagatg gatcgagtca tcggtcgcga 1020  
tcggagattg ctggaatccg acgtttcgaa actcccgat ttacaagccn tagcgaaaga 1080  
aacatatcgt aaacacccat cgacacctct aacactgcg aggattgcga tccaaagcat 1140  
tgaagttagt ggctactaca tccccaaaga cagaggctt agcgtcaaca tttggcgat 1200  
cggctgggac ccaagtgtt gggagaatc atcggagttc tgcctgaaa gattcttctc 1260  
tgaggagaat ggggaagatc gtccaggcg gaatgatttt gagctgattc cgtttggagc 1320  
agggaggaga atttgtgctg ggacaaggat gggaatggtc cttgtaagtt atattttggg 1380

1440 cacttgltc nactcttg attgaaat accaaatgg gtaagtga taaatga  
1500 tgaagtctt ggtcttgat tgaatggc ggtctctc tggctagc tgaatcag  
1560 atgggocca aggtgagc ttaatgag tgaatggc ggtctgac caaatat  
1620 ggttgltc tcttgat ttaatat ataatg taaatgta ttaatgta  
1648 ataatgta aaaaaa

<210> 10  
<211> 506  
<212> PRI  
<213> vtoia  
<400> 10

1 Met Ala Ile Pro Val Thr Asp Leu Ala Val Ile Leu Phe Leu

20 Ile Thr Arg Phe Leu Val Arg Ser Leu Phe Lys Lys Pro Thr Gly Pro

35 Leu Pro Pro Gly Pro Ser Gly Tyr Pro Leu Val Gly Ala Leu Pro Leu

50 Leu Gly Ala Met Pro His Val Thr Leu Ala Asn Leu Ala Lys Tyr

65 Gly Pro Ile Met Tyr Leu Lys Met Gly Thr Cys Asp Met Val Val Ala

85 Ser Thr Pro Asp Ser Ala Arg Ala Phe Leu Lys Thr Leu Asp Asn

100 Phe Ser Asp Arg Pro Pro Asn Ala Gly Ala Thr His Leu Ala Tyr Gly

115 Ala Gln Asp Leu Val Phe Ala Lys Tyr Gly Pro Arg Tyr Lys Thr Leu

130 Arg Lys Leu Ser Asn Leu His Met Leu Gly Gly Lys Ala Leu Asp Asp

145 Trp Ala His Val Arg Ala Asn Gln Leu Gly His Met Leu Asn Ala Met

150

155

160

- 13 -

Cys Glu Ala Ser Arg Cys Gly Glu Pro Val Val Leu Ala Glu Met Leu  
 165 170 175

Thr Tyr Ala Met Ala Asn Met Ile Gly Gln Val Ile Leu Ser Arg Arg  
 180 185 190

Val Phe Val Thr Lys Gly Thr Glu Ser Asn Glu Phe Lys Asp Met Val  
 195 200 205

Val Glu Leu Met Thr Ser Ala Gly Tyr Phe Asn Ile Gly Asp Phe Ile  
 210 215 220

Pro Ser Ile Ala Trp Met Asp Leu Gln Gly Ile Glu Arg Gly Met Lys  
 225 230 235 240

Lys Leu His Thr Lys Phe Asp Val Leu Leu Thr Lys Met Met Lys Glu  
 245 250 255

His Arg Ala Thr Ser His Glu Arg Glu Gly Lys Ser Asp Phe Leu Asp  
 260 265 270

Val Leu Leu Glu Glu Cys Glu Asn Thr Asn Gly Glu Lys Leu Asn Val  
 275 280 285

Thr Asn Val Lys Ala Val Leu Leu Asn Leu Phe Thr Ala Gly Thr Asp  
 290 295 300

Thr Ser Ser Ser Ile Ile Glu Trp Ala Leu Thr Glu Met Met Lys Asn  
 305 310 315 320

Pro Thr Ile Leu Lys Lys Thr Gln Glu Glu Met Asp Arg Val Ile Gly  
 325 330 335

Arg Asp Arg Arg Leu Leu Glu Ser Asp Val Ser Lys Leu Pro Tyr Leu  
 340 345 350

Gln Ala Ile Ala Lys Glu Thr Tyr Arg Lys His Pro Ser Thr Pro Leu  
 355 360 365

Asn Leu Pro Arg Ile Ala Ile Gln Ala Cys Glu Val Asp Gly Tyr Tyr  
 370 375 380

- 14 -

Ile Pro Lys Asp Thr Arg Leu Ser Val Asn Ile Trp Ala Ile Gly Arg 385  
390 395

Asp Pro Ser Val Trp Gln Asn Pro Ser Gln Phe Ser Pro Gln Arg Phe 405  
410 415

Leu Ser Gln Gln Asn Gly Lys Ile Ser Pro Gly Gly Asn Asp Phe Gln 420  
430 435

Leu Ile Pro Phe Gly Ala Gly Arg Arg Ile Cys Ala Gly Thr Arg Met 440  
445

Gly Met Val Leu Val Ser Tyr Ile Leu Gly Thr Leu Val His Ser Phe 450  
455

Asp Trp Lys Leu Pro Asn Gly Val Ser Gln Ile Asn Met Asp Gln Ser 465  
470 475

Phe Gly Leu Ala Leu Gln Lys Ala Val Pro Leu Ser Ala Thr Val Ser 485  
490 495

Pro Arg Leu Ala Pro Ser Ala Tyr Val Ile 500  
505

<210> 11

<211> 1782

<212> DNA

<213> v101a

<220>

<221> misc feature

<222> (307)..(307)

<223> n = any nucleotide

<400> 11

bgacaaatag gcaattctag tcaacagact cgttgtccgg gctatattt tcttgatcaa 60  
tcgtttatc gttcgttata ttctcaagaa accaaacaga ccgtccccc ccgttctct 120  
ccgttggccc ttgttggggg cctccctct cctcagggcc atgcttcag tcgacatagc 180  
caactcgtc aagaaatag gtccatcat gcaactaaa atgggcaagt ggcacatagt 240  
gttcgcgtcc accccagat cggctogaga ctctctcaaa agcttagac tcaactctc 300  
caacgncac ccaacggcg ggcattcca cctagcgctac ggcgcggcag acttagctct 360  
cgccaagtac ggtccgaggt ggaagactt agaaatlg agcaacctc acatgcttag 420

- 15 -

cgggaaggcg ttggatgatt gggcaaatgt gagggtoacc gagctaggac acatgottaa 480  
 agccatgtgc gaggcgagcc ggtgcgggga gcccggtgtg ctggccgaga tgctcacgta 540  
 cgcoatggcg aacatgatcg gtcaagtgat actcaqccgg cgcgtgttcg tgacczaagg 600  
 gaccgagtcct aacgagttca aagacatggt ggtcgagttg atgacgtccg ccgggtactt 660  
 caacatcggg gaottcatac cctcgatcgc ttggatggat ttgcaaggga tcgagcgagg 720  
 gatgaagaag ctgcacaogm agtttgatgt gttattqacg aagatggtga aggagcatag 780  
 agcgacgagt catgagcgca aagggaaggc agatttcctc gacgtttctc tggagaagtg 840  
 cgacaataca aatggggaga agcttagtat taccaatata aaagctgtcc ttttgaatct 900  
 attcaaggcg ggcacggaca catottcgag catxatogaa tgggcgttaa cggagatgat 960  
 caagaatccg acgatottaa aaaaggcgca agaggagatg gatcgagtca tcggctcgtga 1020  
 tcggaggctg ctggaatcgg acatatcgag cctcccgtae ctacaagcca ttgctaaga 1080  
 aacgtatcgc aaacacccgt cgacgcctct caacttgcgg aggattgcga tccaagcatg 1140  
 tgaagttgat ggctactaca tccctaagga cgcgaggctt agcgtgaaca tttgggcgat 1200  
 cggtcgggac ccgaatgttt gggagaatcc gttggagttc ttgcggaaa gattcttgtc 1260  
 tgaagagaat gggaagatca atcccggttg gaatgatttt aagctgattc cgtttggagc 1320  
 cgggaggaga atttgtcggg ggacaaggat gggaatggtc cttgtaagtt atattttggg 1380  
 caotttggtc cattottttg attggaatt accaatggtt gtcgctgagc ttaatatgga 1440  
 tgaagttttt gggcttgcac tgcaaaaggo cgtgcgcctc tcggccttgg tcagcccacg 1500  
 gttggcctca aaccgtaag caaccbgagc taatgggctg ggccctagttt tgtgggccct 1560  
 aatttagaga cttttgtgtt ttaagggtgt tactttatta attgggtgtt taaatgtgtg 1620  
 ttttaatttg tatttatggt taattatgao tttattgtat aattatttat tttcccttc 1680  
 tgggtatttt atccatttaa ttttcttca gaattatgat catagttatc agaataaaat 1740  
 tgaataaat gaatcgaaa aaaaaaaaaa aaaaaaaaaa aa 1782

<210> 12  
 <211> 506  
 <212> FRT  
 <213> viola

<400> 12

Met Ala Ile Leu Val Thr Asp Phe Val Val Ala Ala Ile Ile Phe Leu  
 1 5 10 15

Ile Thr Arg Phe Leu Val Arg Ser Leu Phe Lys Lys Phe Thr Arg Pro 20  
 25  
 30  
 Leu Pro Pro Gly Pro Leu Gly Trp Pro Leu Val Gly Ala Leu Pro Leu 35  
 40  
 45  
 Leu Gly Ala Met Pro His Val Ala Leu Ala Lys Leu Ala Lys Tyr 50  
 55  
 60  
 Gly Pro Ile Met His Leu Lys Met Gly Thr Cys Asp Met Val Val Ala 65  
 70  
 75  
 Ser Thr Pro Glu Ser Ala Arg Ala Phe Leu Lys Thr Leu Asp Leu Asn 85  
 90  
 95  
 Phe Ser Asn Arg Pro Pro Asn Ala Gly Ala Ser His Leu Ala Tyr Gly 100  
 105  
 110  
 Ala Glu Asp Leu Val Phe Ala Lys Tyr Gly Pro Arg Trp Lys Thr Leu 115  
 120  
 125  
 Arg Lys Leu Ser Asn Leu His Met Leu Gly Gly Lys Ala Leu Asp Asp 130  
 135  
 140  
 Trp Ala Asn Val Arg Val Thr Glu Leu Gly His Met Leu Lys Ala Met 145  
 150  
 155  
 Cys Glu Ala Ser Arg Cys Gly Glu Pro Val Val Leu Ala Glu Met Leu 165  
 170  
 175  
 Thr Tyr Ala Met Ala Asn Met Ile Gly Glu Val Ile Leu Ser Arg Arg 180  
 185  
 190  
 Val Phe Val Thr Lys Gly Thr Glu Ser Asn Glu Phe Lys Asp Met Val 195  
 200  
 205  
 Val Glu Leu Met Thr Ser Ala Gly Tyr Phe Asn Ile Gly Asp Phe Ile 210  
 215  
 220  
 Pro Ser Ile Ala Trp Met Asp Leu Glu Gly Ile Glu Arg Gly Met Lys 230  
 235  
 240

- 17 -

Lys Leu His Thr Lys Phe Asp Val Leu Leu Thr Lys Met Val Lys Glu  
 245 250 255

His Arg Ala Thr Ser His Glu Arg Lys Gly Lys Ala Asp Phe Leu Asp  
 260 265 270

Val Leu Leu Glu Glu Cys Asp Asn Thr Asn Gly Glu Lys Leu Ser Ile  
 275 280 285

Thr Asn Ile Lys Ala Val Leu Leu Asn Leu Phe Thr Ala Gly Thr Asp  
 290 295 300

Thr Ser Ser Ser Ile Ile Glu Trp Ala Leu Thr Glu Met Ile Lys Asn  
 305 310 315 320

Pro Thr Ile Leu Lys Lys Ala Gln Glu Glu Met Asp Arg Val Ile Gly  
 325 330 335

Arg Asp Arg Arg Leu Leu Glu Ser Asp Ile Ser Ser Leu Pro Tyr Leu  
 340 345 350

Gln Ala Ile Ala Lys Glu Thr Tyr Arg Lys His Pro Ser Thr Pro Leu  
 355 360 365

Asn Leu Pro Arg Ile Ala Ile Gln Ala Cys Glu Val Asp Gly Tyr Tyr  
 370 375 380

Ile Pro Lys Asp Ala Arg Leu Ser Val Asn Ile Trp Ala Ile Gly Arg  
 385 390 395 400

Asp Pro Asn Val Trp Glu Asn Pro Leu Glu Phe Leu Pro Glu Arg Phe  
 405 410 415

Leu Ser Glu Glu Asn Gly Lys Ile Asn Pro Gly Gly Asn Asp Phe Lys  
 420 425 430

Leu Ile Pro Phe Gly Ala Gly Arg Arg Ile Cys Ala Gly Thr Arg Met  
 435 440 445

Gly Met Val Leu Val Ser Tyr Ile Leu Gly Thr Leu Val His Ser Phe  
 450 455 460

- 18 -

ASP TRP LYS LEU PRO ASN GLY VAL ALA GLN LEU ASN MET ASP GLN SER 468  
470 475 480

PRO GLY LEU ALA LEU GLN LYS ALA VAL PRO LEU SER ALA LEU VAL SER 485  
490 495

PRO ARG LEU ALA SER ASN PRO TYR ALA THR 500  
505

<210> 13  
<211> 1659  
<212> DNA  
<213> SALYER

<400> 13  
CATGGAGGCC AAGAAATA TGTGTGAT TGCTAGGCG CTTGTGTAG GATCCTTACT 60  
CTACATTTTG ATCGGTATGT TTATCTCASA ATGAGGCAAC ACGGCAAC CTCGTCCCCC 120  
GGGGCGGAGG AGCTTCTTAG TGGTGGGCTC CTTCTCCCTG CTGGGCGACA TGCCCATGTT 180  
CGCCCTAGCA AAAATGGCCA AAACTTACGG CCGATCATAG TACTTGAABA TGGGCAAGAT 240  
CGGCATGTC GTGGGCTCCA CGCAAGACGC GGGCGGGGCG TTCTTAAGA CCGCAAGAGC 300  
TAATTTCTCG AACGGGCGGG TCAACGGGGG TGCCACATCC CTGGCATACA ATGCCCAAGG 360  
CATGTCTGTTT GCGCGGTAGG GCGCCAAAGT GAGATCTGCT AGGAAGGTGA GCAGTCTCCA 420  
CATGTCTGGCG GCGATCGCAAG AGGCGCAAGCG CTCGGTGGGAG GTCTGTGGGTT TGGCGGAGAT 540  
GCTGTCTTAC GCGACGGCGG AACATGATCGG GCAAGGTGATA TTGAGCGGGA GAGTTTCTCT 600  
GAGGAAAGGG AAGGAGATGA ATGATATCCA GGAATCTGTG GTGGTGGCTC TGCACCAAGC 660  
TGGCTATTC AACATTTGTTG ATTTCAATCC ATGGCTCTGCT TGGATGGGAT TCGAAGGGGAT 720  
TGGAGAGAGG ATGAAAGAA TGGCAAGGA GTGGGAGCGC TTGATCTGGT AAGATCTGGG 780  
TGATCTGATG AATCAACCT ACAACGCA CCAAGGCA GATCTCTCTG ATCTCTCTCT 840  
GGCAATCAT GTGTGTGGA GTAAAGGGA TGTAGGAGAT TGCAGGAT TGCAGGCTCA CCAACCAACA 900  
CATTAAGAC CTTTACCTGA ATTTACTTA TGGAGGGGCA GACCATCTGT CGAGCATTAAT 960  
AGATGGGCA TTAGGCGGGA TCAATCAAGA TCAATCAAGA TCAATCAAGA TCAATCAAGA 1020



- 19 -

```

gttgagcgctc aacatctggg ccataggccg agatcccgac gtttgggaga atcccottga 1260
cttcaaccct gacaggttta tgtcagggtt gcaggggatt gagccggag ggaatcactt 1320
cgagctcatt ccccttgggg cggggcgag gatctgcgc gccagcaga tggggattgt 1380
aatgtggag tatttgcctg ccacactcgt gcactcttcc gaatgggatt tgcgggcgg 1440
ctcagcggag atggacatgg aggaggtgtt cgggctggcc ttgcagaaag ctgtaccact 1500
tgctgctagg ctcaactcta ggttgccttc acattgctat gcacctcctt ctatttaatt 1560
tgcataatta catatgttgt gttacattga gcccttgcct atgttctatc caaootatct 1620
tataacttgt gcataaatt gaaaaaaaaa aaaaaaaaaa 1659

```

```

<210> 14
<211> 520
<212> PRT
<213> salvia

```

```

<400> 14

```

```

Gly Thr Ser Met Glu Ala Gln Glu Asn Met Leu Leu Ile Ala Arg Ala
1 5 10 15

```

```

Leu Val Val Ala Ser Leu Leu Tyr Ile Leu Ile Arg Met Phe Ile Ser
20 25 30

```

```

Lys Leu Ser Thr Thr Gly His Pro Leu Pro Pro Gly Pro Arg Gly Phe
35 40 45

```

```

Leu Val Val Gly Ser Leu Pro Leu Leu Gly Asp Met Pro His Val Ala
50 55 60

```

```

Leu Ala Lys Met Ala Lys Thr Tyr Gly Pro Ile Met Tyr Leu Lys Met
65 70 75 80

```

```

Gly Thr Val Gly Met Val Val Ala Ser Thr Pro Asp Ala Ala Arg Ala
85 90 95

```

```

Phe Leu Lys Thr His Asp Ala Asn Phe Ser Asn Arg Pro Val Asn Ala
100 105 110

```

```

Gly Ala Thr Ile Leu Ala Tyr Asn Ala Gln Asp Met Val Phe Ala Pro
115 120 125

```



- 21 -

Arg Leu Leu Glu Ser Asp Ile Pro Asn Leu Pro Tyr Leu Lys Ala Ile  
355 360 365

Cys Lys Glu Ala Tyr Arg Lys His Pro Ser Thr Pro Leu Asn Leu Pro  
370 375 380

Arg Ile Ser Thr Asp Ala Cys Val Val Asp Gly Tyr His Ile Pro Lys  
385 390 395 400

Asn Thr Arg Leu Ser Val Asn Ile Trp Ala Ile Gly Arg Asp Pro Asp  
405 410 415

Val Trp Glu Asn Pro Leu Asp Phe Asn Pro Asp Arg Phe Met Ser Gly  
420 425 430

Leu Gln Gly Ile Glu Pro Gly Gly Asn His Phe Glu Leu Ile Pro Phe  
435 440 445

Gly Ala Gly Arg Arg Ile Cys Ala Gly Ser Arg Met Gly Ile Val Ile  
450 455 460

Val Glu Tyr Leu Leu Ala Thr Leu Val His Ser Phe Glu Trp Asp Leu  
465 470 475 480

Pro Ala Gly Ser Ala Glu Met Asp Met Glu Glu Val Phe Gly Leu Ala  
485 490 495

Leu Gln Lys Ala Val Pro Leu Ala Ala Arg Leu Thr Pro Arg Leu Pro  
500 505 510

Ser His Cys Tyr Ala Pro Pro Ser  
515 520

<210> 15  
<211> 1617  
<212> DNA  
<213> *salvia*

<400> 15  
agatagtaag catggaagcc caagaaaata tgttgttgat tgotaggcca ctgtttgtag 60  
catccttaet ctacattttg atccgtatgt ttatctcaaa attgagcaac ccgggccacc 120  
ctctgccccg ggggcgagg ggctttccag tggtyggctc ccttccattg ctgggcgaca 180  
tgccacatgt tgccctagca aaaatggcca aaacttatgg ccgatcatg tacttgaaaa 240



- 23 -

Ala Ser Leu Leu Tyr Ile Leu Ile Arg Met Phe Ile Ser Lys Leu Ser  
 20 25 30

Thr Pro Gly His Pro Leu Pro Pro Gly Pro Arg Gly Phe Pro Val Val  
 35 40 45

Gly Ser Leu Pro Leu Leu Gly Asp Met Pro His Val Ala Leu Ala Lys  
 50 55 60

Met Ala Lys Thr Tyr Gly Pro Ile Met Tyr Leu Lys Met Gly Thr Val  
 65 70 75 80

Gly Met Val Val Ala Ser Thr Pro Asp Ala Ala Arg Ala Phe Leu Lys  
 85 90 95

Thr Gln Asp Ala Asn Phe Ser Asn Arg Pro Val Asn Ala Gly Ala Thr  
 100 105 110

Ile Leu Ala Tyr Asn Ala Gln Asp Met Val Phe Ala Pro Tyr Gly Pro  
 115 120 125

Lys Trp Arg Leu Leu Arg Lys Leu Ser Ser Leu His Met Leu Gly Ser  
 130 135 140

Lys Ala Leu Glu Glu Trp Ala Asp Val Arg Thr Ser Glu Val Gly His  
 145 150 155 160

Met Leu Ala Ala Met His Glu Ala Ser Arg Leu Asp Glu Ala Val Gly  
 165 170 175

Leu Pro Glu Met Leu Val Tyr Ala Thr Ala Asn Met Ile Gly Lys Val  
 180 185 190

Ile Leu Ser Arg Arg Val Phe Val Thr Lys Gly Lys Glu Met Asn Glu  
 195 200 205

Phe Lys Glu Met Val Val Glu Leu Met Thr Thr Ala Gly Tyr Phe Asn  
 210 215 220

Ile Gly Asp Phe Ile Pro Trp Leu Ala Trp Met Asp Leu Gln Gly Ile  
 225 230 235 240

Arg Arg ile cys ala gly ser Arg Met gly ile Val ile Val gin Tyr  
450 455  
ile gin pro gly gly Asn His phe gin leu ile pro phe gly ala gly  
435 440 445  
Asn pro leu Asp phe Asn pro Asp Arg phe Met ser gly leu gin gly  
420 425 430  
leu ser val Asn ile trip ala ile gly Arg Asp pro Asp val trip gin  
405 410 415  
the Asp ala cys val val Asp gly Tyr His ile pro lys Asn the Arg  
385 390 395 400  
ala Tyr Arg lys His pro ser the pro leu Asn leu pro Arg ile ser  
370 375 380  
gin ser Asp ile pro Asn leu pro Tyr leu lys ala ile cys lys gin  
355 360 365  
ala His gin gin Met Asp Arg val ile gly Arg gin Arg Arg leu leu  
340 345 350  
gin trip ala leu ala gin Met ile lys Asn pro ser ile gin lys Arg  
325 330 335  
leu leu Asn leu phe the ala gly the Asp the ser ser ser ile ile  
305 310 315 320  
asp Asp Asp gin Asp cys lys leu the the the the the lys ala leu  
290 295 300  
pro Asp leu leu Asp ser leu leu ala Asn His Asp Asp gin ser lys  
275 280 285  
lys Met leu Asp Asp Arg leu lys ser the Tyr lys Arg Asn Asp lys  
260 265 270  
gin Arg gly Met lys lys leu His lys lys trip Asp Arg leu ile gly  
245 250 255

- 25 -

Leu Leu Ala Thr Leu Val His Ser Phe Glu Trp Asp Leu Pro Ala Gly  
465 470 475 480

Ser Ala Glu Met Asp Met Glu Glu Val Phe Gly Leu Ala Leu Gln Lys  
485 490 495

Ala Val Pro Leu Ala Ala Arg Leu Thr Pro Arg Leu Pro Ser His Cys  
500 505 510

Tyr Ala Pro Pro Ser Ile  
515

<210> 17  
<211> 1730  
<212> DNA  
<213> sollya

<220>  
<221> misc\_feature  
<222> (1372)..(1372)  
<223> n = any nucleotide

<400> 17  
gatggctaact accttagaat tcattctatg cttoaccatt actgcacttc cttttctata 60  
ttgcataact aacatggcga tcttccttaa ccgtcaaccg aggtcaactc caccaggtcc 120  
aagaccatgg cctattgtag gaaacctccc acaccttggc aocaaagccac accactccat 180  
agctgccatg gctgggaat accgtccct cctgcacctc cgcattggga tegtgcacgt 240  
ggtgggttgc gcctctgtg atgtggcggc acagttcttg aagaatgatg ccaacttctc 300  
tagccgggca ccgaattctg gtgctaagca tatggcttat aactatocag acatgggttt 360  
tgcacctac ggtccaggt ggccgatgtt gaggaaaatt tgtgcccttc atatatcttc 420  
cgtaagggt ctcgatgatt ttcctcggt gcgtgaggag gaggttgcca tactcgcag 480  
gacctagca cagcagggcc aaaagccggt gaatttggg cagttgttct ctacgtgtaa 540  
tgctaattgg ctatcagtc tgatgctagg caggaggttg ttcagcacag aagttgatto 600  
aaaagcatat gatttcaaac aaatggtggt ggagctgatg actctagccg gtgagtttaa 660  
cgtcagtgat ttcattccac cctcgagtg gctagacttg caaggcgtgg cagcgaat 720  
gaagaacgtg cacaatcgat tcgatgcgtt totgaatga attttggagg agcataagct 780  
gaaactaat aatagtggac atggggaaca aaacatatg gacttggtga gtacgttgat 840  
tttgcctaag gatgatgctg atagtgggg aggaaaactc actgatactg aaatcaagc 900

gctgcttctg aattgcttc ctgcctggag ggaactca tccagccaa tagaatggt 960  
tatacttag ctatagca atctaaat cttagccaa gcccagaag agtggactt 1020  
ggtggttctg ccaatagac ttgtaacga ttggaactc aaacattaa cctaacaa 1080  
agcaatctc aaagaaact tccgtaca tccgtcac ccaactcac tccagagat 1140  
cgcaaccga agcgtgaaa tcaaggtt tcaattca agggctcaa cactctctg 1200  
taaatatgg gcaatagcc gtaacaaa caactggct gaaacttgg tattccagaa 1260  
tgaacttc tcatagatg ttgaagttc taacttga gtcnaagaa gtaatttga 1320  
attgatacaa ttggggcggt ggagaaat ttgtgctgg atgaacttg gntacagcat 1380  
gtccagta gtaactgaa ogtaattca tgaattcaa tggagttcg cagaaaggga 1440  
attgocagaa aatagaaia ttgaggaag ctatggat agctgcaac ggaactggcc 1500  
atgattct caccnaagc cagacaga ccaagatt taccagttc atggagtgt 1560  
aaactagta cactagaa ctgaacaga agtgcaga ttctgctta tatgctta 1620  
tgaatggcga coctgggtg ttttatgt atgttctgt acaagtctg agocctggcg 1680  
ggcaatggac aattatata gtaattta aaaaaaa aaaaaaa 1730

<210> 18  
<211> 521  
<212> PRT  
<213> sollya

1 Met Ala Thr Thr Leu Glu Phe Ile Leu Cys Phe Thr Ile Thr Ala Leu

20 Pro Phe Leu Tyr Cys Ile Leu Asn Met Arg Ile Leu Asn Arg His

35 Pro Arg Ser Leu Pro Gly Pro Arg Pro Arg Pro Ile Val Gly Asn

50 Leu Pro His Leu Gly Thr Lys Pro His His Ser Ile Ala Ala Met Ala

65 Arg Lys Tyr Gly Pro Leu Leu His Leu Arg Met Gly Ile Val His Val

80



- 27 -

Val Val Ala Ala Ser Ala Asp Val Ala Ala Gln Phe Leu Lys Asn Asp  
85 90 95

Ala Asn Phe Ser Ser Arg Pro Pro Asn Ser Gly Ala Lys His Met Ala  
100 105 110

Tyr Asn Tyr His Asp Met Val Phe Ala Pro Tyr Gly Pro Arg Trp Arg  
115 120 125

Met Leu Arg Lys Ile Cys Ala Leu His Ile Phe Ser Ala Lys Ala Leu  
130 135 140

Asp Asp Phe His Arg Val Arg Glu Glu Glu Val Ala Ile Leu Ala Arg  
145 150 155 160

Thr Leu Ala His Ala Gly Gln Lys Pro Val Asn Leu Gly Gln Leu Phe  
165 170 175

Ser Thr Cys Asn Ala Asn Ala Leu Ser Val Leu Met Leu Gly Arg Arg  
180 185 190

Leu Phe Ser Thr Glu Val Asp Ser Lys Ala Tyr Asp Phe Lys Gln Met  
195 200 205

Val Val Glu Leu Met Thr Leu Ala Gly Glu Phe Asn Val Ser Asp Phe  
210 215 220

Ile Pro Pro Leu Glu Trp Leu Asp Leu Gln Gly Val Ala Ala Lys Met  
225 230 235 240

Lys Asn Val His Asn Arg Phe Asp Ala Phe Leu Asn Val Ile Leu Glu  
245 250 255

Glu His Lys Leu Lys Leu Asn Asn Ser Gly His Gly Glu Gln Lys His  
260 265 270

Met Asp Leu Leu Ser Thr Leu Ile Leu Leu Lys Asp Asp Ala Asp Ser  
275 280 285

Glu Gly Gly Lys Leu Thr Asp Thr Glu Ile Lys Ala Leu Leu Leu Asn  
290 295 300

Leu Phe Ser Ala Gly Thr Asp Thr Ser Ser Ser Thr Ile Glu Trp Val 305  
 310  
 315  
 320  
 Ile Ala Glu Leu Ile Arg Asn Pro Lys Ile Leu Ala Glu Ala Arg 325  
 330  
 335  
 Glu Leu Asp Leu Val Val Gly Pro Asn Arg Leu Val Thr Asp Leu Asp 340  
 345  
 350  
 Leu Lys Glu Leu Thr Tyr Leu Glu Ala Ile Val Lys Glu Thr Phe Arg 355  
 360  
 365  
 Leu His Pro Ala Thr Pro Leu Ser Leu Pro Arg Ile Ala Thr Glu Ser 370  
 375  
 380  
 Cys Glu Ile Asn Gly Phe Tyr Ile Pro Lys Gly Ser Thr Leu Leu Val 385  
 390  
 395  
 Asn Ile Trp Ala Ile Gly Arg Asp Pro Asn Thr Trp Ala Glu Pro Leu 405  
 410  
 415  
 Val Phe Arg Pro Glu Arg Phe Leu Ser Asp Gly Glu Ser Pro Asn Val 420  
 425  
 430  
 Asp Val Lys Gly Arg Asn Phe Glu Leu Ile Pro Phe Gly Ala Gly Arg 435  
 440  
 445  
 Arg Ile Cys Ala Gly Met Asn Phe Gly Leu Arg Met Val Glu Leu Val 450  
 455  
 Thr Ala Thr Leu Ile His Ala Phe Asn Trp Glu Leu Pro Glu Gly Glu 465  
 470  
 475  
 480  
 Leu Pro Glu Asn Met Asn Met Glu Glu Asp Tyr Gly Ile Ser Leu Glu 485  
 490  
 495  
 Arg Thr Val Pro Leu Val Val His Pro Lys Pro Arg Leu Asp His Glu 500  
 505  
 510  
 Val Tyr Glu Ser His Gly Val Asn 515  
 520

- 29 -

<211> 37  
 <212> DNA  
 <213> petunia

<400> 19  
 aaaatcgata ccatgggtctt tttttctttg totatac

37

<210> 20  
 <211> 1748  
 <212> DNA  
 <213> clitoria

<400> 20  
 ggatccaaca atgttctctt taagagaaat tgggtatca attttgatct toatgatcac 60  
 ccatottgtg attogtttag ttctgaaaga gaaggaaaca cggaaacttc caccagggcc 120  
 aaaaggttgg ccaattgtgg gtgcactgoc totaatggga agcatgcccc atgtcacact 180  
 otcagaaatg gctaaaaaat atggacctgt tatgtacott aaaatgggca caaacaacat 240  
 ggctgtagea totactccct ctgcagctcg tgcattcttc aaaaaccttg accttaactt 300  
 ctccaatcgc ccccaaatg ctggggcaac tcacttagct tatgatgccc aggcactggt 360  
 gtttgotgat tacggatcta ggtggaagtt gottagaaaa ctaagcaact tacacatgot 420  
 tggaggaaag gctcttgaag aatggtcaca agttaqagag attgagatgg ggcacatgct 480  
 togtgcaatg tacgattgta gtggtggcgg tgacggcac aacgcaatg atggcaaca 540  
 gaaaaaggtt actcgtcatg agcctattgt ggtggtgaa atgttaacat acgcgatggc 600  
 caacatgata ggtcaagtga tottgagcog togtgtatto gagacaaagg gttcggaato 660  
 gaacgagttt aaggacatgg tggttcagot catgacggtt gctggctact ttazoattgg 720  
 tgattttatt ccttttttgg ctgccttoga cctccaaggc atcgagcgtg gcatgaaaac 780  
 tttgcataac aagttcgatg ttttgttgac gacaatgatt catgagcatg tggcttctgc 840  
 toatasacga aagggtasac ctgatttctt ggtgttctc atggctcacc ataccaacga 900  
 gtctcatgaa ctgtcgctca ccaacatcaa agcctctctc ttacatctat ttactgcagg 960  
 cacagataca tcataagta tcatagagtg ggcactagca gagatgttga taaaccctaa 1020  
 aatcatgaag aagtgcatg aggaaatgga caaagtgata ggcaaggata gaaggctaaa 1080  
 agaatccgac atagaaaato tcccttactt gcaggcaatt tgcnaagaga catatagaaa 1140  
 gcacccatac acgccactca acttgccatg aatctcctcc caagcatgcc aagtgaatgg 1200  
 ctactacatc ccaaagaaca ctaggcttag tgtcaacatc tgggocattg gaagagaccc 1260  
 taatgtgtgg gagaacctt tggagttcaa tccagagagg tttatgggtg ccaataagac 1320

1380      tarracaa ctyggagatg attgagot aatccatt ggtcggga gaagattg  
1440      tgcgggaca agatggga ctygtgtgt tcaatcat ttgggacat tggatata  
1500      cctgattg agttacaa atggtgtgt ggaattgaa atggagaa ctttgat  
1560      tgccttgag aaaaagatc cacttcgc ttgattacc ctaggtgc cccaactg  
1620      ttacatgt aatccat aattgact tagttatg gtaattata accaatag  
1680      taatcatgt ttgtatat gttctgat tccagatga ttctctgt gtaggaaa  
1740      tctgaccc ccaactggg taatgttt aatattt catattt aaaaaaaa  
1748      aaaaaa

<210> 21  
<211> 525  
<212> 527  
<213> 527  
<400> 21

Met Phe Leu Leu Arg Gln Ile Gly Val Ser Ile Leu Ile Phe Met Ile  
1  
5  
10  
15

Thr His Leu Val Ile Arg Leu Val Leu Lys Gln Lys Gln Arg Lys  
20  
25  
30

Leu Pro Pro Gly Pro Lys Gly Trp Pro Ile Val Gly Ala Leu Pro Leu  
35  
40  
45

Met Gly Ser Met Pro His Val Thr Leu Ser Gln Met Ala Lys Lys Tyr  
50  
55  
60

Gly Pro Val Met Tyr Leu Lys Met Gly Thr Asn Asn Met Ala Val Ala  
65  
70  
75  
80

Ser Thr Pro Ser Ala Ala Arg Ala Phe Leu Lys Thr Leu Asp Leu Asn  
85  
90  
95

Phe Ser Asn Arg Pro Pro Asn Ala Gly Ala Thr His Leu Ala Tyr Asp  
100  
105  
110

Ala Gln Asp Met Val Phe Ala Asp Tyr Gly Ser Arg Trp Lys Leu Leu  
115  
120  
125

-31-

Arg Lys Leu Ser Asn Leu His Met Leu Gly Gly Lys Ala Leu Glu Glu  
130 135 140

Trp Ser Gln Val Arg Glu Ile Glu Met Gly His Met Leu Arg Ala Met  
145 150 155 160

Tyr Asp Cys Ser Gly Gly Gly Asp Gly Asn Asn Asp Asn Asp Gly Asn  
165 170 175

Lys Lys Lys Gly Thr Arg His Glu Pro Ile Val Val Ala Glu Met Leu  
180 185 190

Thr Tyr Ala Met Ala Asn Met Ile Gly Pro Ser Asp Leu Glu Pro Ser  
195 200 205

Cys Ile Pro Arg Gln Arg Val Arg Asn Arg Thr Ser Leu Arg Thr Trp  
210 215 220

Trp Phe Lys Leu Met Thr Val Ala Gly Tyr Phe Asn Ile Gly Asp Phe  
225 230 235 240

Phe Pro Phe Leu Ala Arg Arg Arg Arg Gln Gly Ile Glu Arg Gly Met  
245 250 255

Lys Thr Leu His Asn Lys Lys Asp Asp Leu Leu Thr Thr Met Ile His  
260 265 270

Glu His Val Ala Ser Ala His Lys Arg Lys Gly Lys Pro Pro Phe Leu  
275 280 285

Asp Val Leu Met Ala His His Thr Asn Glu Ser His Glu Leu Ser Leu  
290 295 300

Thr Asn Ile Lys Ala Leu Leu Leu Asn Leu Phe Thr Ala Gly Thr Asp  
305 310 315 320

Thr Ser Ser Ser Ile Ile Glu Trp Ala Leu Ala Glu Met Leu Ile Asn  
325 330 335

Pro Lys Ile Met Lys Lys Val His Glu Glu Met Asp Lys Val Ile Gly  
340 345 350

370 375 380

ASN LEU PRO ARG ILE SER SER GLN ALA CYS GLN VAL ASN GLY TYR TYR 385  
390  
395  
400

105  
410  
415

420  
425  
430

Met Gly Ala Asn Lys Thr Ile Asp Pro Arg Gly Asn Asp Phe Glu Leu 435  
440  
445

150 The pro the G1y ALA G1y Arg Arg. The G1y ALA G1y The Arg Met G1y  
455  
460

465 The Val Ten Val Gln Tyr 470 The Ten Gln Tyr 475 The Ten Val His Ser Phe Asp 480

TRP LYE TEN PRO ASN GUY VAL VAL GUY TEN LEU ASN MET GUY GUY THE  
485 490 495

500 505 510

Arg Ieu Pro Pro Thx Ala Tyr Asn Val Ile Asn Ser Ser 515  
520  
525

22	<210>
1584	<211>
DNA	<212>
gentiana	<213>

[illegible]

- 33 -

```

aagtcggtag coattggctta gcaatagcgt cgacacccga cgttgctaaa gcgttcotca 300
aaaccctcga tttaaatttc tcgaacccgc caccaaatgc cggagctacc catttagcct 360
ataacgctca agatatggtt tttgcacatt atggtcctaa atggzaattg ttacgtaaac 420
tcagttaactt acucattgcta ggtggcaag ccttggaaaa ttgggctgat gttagaaaaa 480
cagagcttgg ttatatgctt aaagccatgt ttgaatcgag tcaaaaacaat gaggccggtga 540
tgatttcgga gatgctaacg taaggccatgg cgaacatggt aagccaagtt atacttagcc 600
gtcgcgtatt caataaaaaa ggcgcgaat caaacgagtt taaagatatg gtggtcgaat 660
taatgacgag tgccgggtat ttcaatatag gtgattttat accatcaatt ggttggatgg 720
atttgcaagg gattgaaggt ggaatgaana gattgcacaa aaagttozao gttttgttga 780
ctcgattatt ggatgacat aaagaaaga gtcaggagcg taaacaaaaa ccgatttttc 840
ttgattttgt gattgcaat ggcgataatt ctgatggtga aaggtcnaac accgacaaac 900
tcaaggctct tttattgaac ttgtttactg ctggtacgga tacatcatca agcatcattg 960
agtgggcact agcagaactg ctaagaatc ggaoactcct cacccgagcc caggacgaaa 1020
tggtatgggt atcggggcga gaccgogtc ttottgaato agacatcccc aacttaccat 1080
atcttcaagc aatctgcaaa gaaacattcc gtaaacaccc ttcaacacca ttaaccttc 1140
caagggaattg catcagaggc catgttggatg taaatgggtc ctacattccg aaagggaactc 1200
ggctcaacgt caacatatgg gcgattggaa gagaccatc ggtttggggg gataacccga 1260
acgagttcga cccggagagg tttttgtatg ggaggaaatgc taagattgat ccacgaggaa 1320
accattttga attgatccca ttgggtgctg gacgaagaat ttgtgcagga zcaagaatgg 1380
ggatattgct tgttgagtat attttgggga cattggatga tagttttgat tggaaactgg 1440
gattttctga ggatgagott aatatggatg agacatttgg gcttgatctg cagaagctg 1500
tgoctttago ggccatggtt attccacgco ttctcttca tgrttatgct ccttaattca 1560
gagatttaat ttcattgctt gttttattaa tcattttott aatatgaatt gatggaggtt 1620
atctagttat gaaaaataat aatggaggat ttgtttatca tcatgcaaa aaaaaaaaa 1680
aaaa 1684

```

```

<210> 23
<211> 516
<212> FRT
<213> gentiana

```

<400> 23  
 1 Met Ser Pro Ile Tyr Thr Thr Leu Thr Leu His Leu Ala Thr Ala Leu  
 5 10 15  
 20 Phe Leu Phe Phe His Val Gln Tyr Leu Val His Tyr Leu His Gly Lys  
 25 30  
 35 Ala Thr Gly His Arg Cys Arg Arg Leu Pro Pro Gly Pro Thr Gly Trp  
 40 45  
 50 Pro Ile Leu Gly Ala Leu Pro Leu Leu Gly Asp Met Pro His Val Thr  
 55 60  
 65 Phe Ala Asn Met Ala Lys Lys Tyr Gly Ser Val Met Tyr Leu Lys Val  
 70 75  
 80 Gly Ser His Gly Leu Ala Ile Ala Ser Thr Pro Asp Ala Ala Lys Ala  
 85 90 95  
 100 Phe Leu Lys Thr Leu Asp Leu Asn Phe Ser Asn Arg Pro Pro Asn Ala  
 105 110  
 115 Gly Ala Thr His Leu Ala Tyr Asn Ala Gln Asp Met Val Phe Ala His  
 120 125  
 130 Tyr Gly Pro Lys Trp Lys Leu Leu Arg Lys Leu Ser Asn Leu His Met  
 135 140  
 145 Leu Gly Gly Lys Ala Leu Gln Asn Trp Ala Asp Val Arg Lys Thr Gln  
 150 155 160  
 165 Leu Gly Tyr Met Leu Lys Ala Met Phe Gln Ser Ser Gln Asn Gln  
 170 175  
 180 Pro Val Met Ile Ser Gln Met Leu Thr Tyr Ala Met Ala Asn Met Leu  
 185 190  
 195 Ser Gln Val Ile Leu Ser Arg Arg Val Phe Asn Lys Lys Gly Ala Lys  
 200 205  
 210 Ser Asn Gln Phe Lys Asp Met Val Gln Leu Met Thr Ser Ala Gly  
 215 220



- 35 -

Tyr Phe Asn Ile Gly Asp Phe Ile Pro Ser Ile Gly Trp Met Asp Leu  
 225 230 235 240  
 Gln Gly Ile Glu Gly Gly Met Lys Arg Leu His Lys Lys Phe Asp Val  
 245 250 255  
 Leu Leu Thr Arg Leu Leu Asp Asp His Lys Arg Thr Ser Gln Glu Arg  
 260 265 270  
 Lys Gln Lys Pro Asp Phe Leu Asp Phe Val Ile Ala Asn Gly Asp Asn  
 275 280 285  
 Ser Asp Gly Glu Arg Leu Asn Thr Asp Asn Ile Lys Ala Leu Leu Leu  
 290 295 300  
 Asn Leu Phe Thr Ala Gly Thr Asp Thr Ser Ser Ser Ile Ile Glu Trp  
 305 310 315 320  
 Ala Leu Ala Glu Leu Leu Lys Asn Arg Thr Leu Leu Thr Arg Ala Gln  
 325 330 335  
 Asp Glu Met Asp Arg Val Ile Gly Arg Asp Arg Arg Leu Leu Glu Ser  
 340 345 350  
 Asp Ile Pro Asn Leu Pro Tyr Leu Gln Ala Ile Cys Lys Glu Thr Phe  
 355 360 365  
 Arg Lys His Pro Ser Thr Pro Leu Asn Leu Pro Arg Asn Cys Ile Arg  
 370 375 380  
 Gly His Val Asp Val Asn Gly Tyr Tyr Ile Pro Lys Gly Thr Arg Leu  
 385 390 395 400  
 Asn Val Asn Ile Trp Ala Ile Gly Arg Asp Pro Ser Val Trp Gly Asp  
 405 410 415  
 Asn Pro Asn Glu Phe Asp Pro Glu Arg Phe Leu Tyr Gly Arg Asn Ala  
 420 425 430  
 Lys Ile Asp Pro Arg Gly Asn His Phe Glu Leu Ile Pro Phe Gly Ala  
 435 440 445

Gly Arg Arg Ile Cys Ala Gly Thr Arg Met Gly Ile Leu Leu Val Glu  
450  
455  
460

Lys Ile Leu Gly Thr Leu Val His Ser Phe Asp Trp Lys Leu Gly Phe  
470  
475  
480

Ser Glu Asp Glu Leu Asn Met Asp Glu Thr Phe Gly Leu Ala Leu Glu  
485  
490  
495

Lys Ala Val Pro Leu Ala Ala Met Val Ile Pro Arg Leu Pro Leu His  
500  
505  
510

Val Tyr Ala Pro  
515

<210> 24  
<211> 3731  
<212> DNA  
<213> Protein

tcagatag catttggc gagaaatca caaatltga caaagatgt aatttttt  
60  
120 ttcctttt cccctttaa cattaaatgt gataagtag taacaacaac taaccagt  
180 tatttaca gtggagctg gggaggtga atgtacga caacttaca caaccaggt  
240 ggaagggcag ttccggtag agcctcggt gaagaaata ttccgagac aggttga  
300 aatggagcag aagaaaca ctataaata ataaactaa agcatatac tatataac  
360 atagtagcag tagtataaa ggcactgact agcagagaa taactatata ataggaga  
420 agaacatc ccatatcct ccttctact ttactcatg acatcagac ttccctact  
480 aggtcatgt cctcggtgat ctgacttg gcatgtct atctaatcac ctcggtcag  
540 tctctctag gtcctcctc acctcctgt agaatcaca ctcggaact ctcacacct  
600 ctacacgagc atctggactt cctccttta cgtatctga ccatctcat cttgtctc  
660 gcatctttc tgcacatga gatcctca ccttctca agtgcaca tcatatct  
720 tgcacctct agtatcaca aactcctc tgaagatct tatctctgc aacacaaa  
780 tcttaagt gtggttctt gactggaaaa cactcagcc catcaaat agctgtata  
840 accaacatgt atatatat gaagctaat aatactta atcaactt gtaactag  
900 caggaagag aatctcttg acaatcttg gataatga ttgggtcca cactctgtg  
960 aggaatat tttctcag acaaaaaat ttcacagc atgagatg

- 37 -

atagatcaag gtgtcctttt tcatcttatt aggaataaa aatttgaaga tgcaaatcca 1020  
 aggacgatca tgcattotaa attaastagtt aatgattota attttcattt atttaaattt 1080  
 tgattttttgc ggcttcaaat tcatatttga ttttcaacca cataaatatt taattatgac 1140  
 ttgtgttagg caacaaatac caaaagtctt actactttct tttggagatt gacctttota 1200  
 tatecccttc ccaatttgat ogatcgagct ctogtaatot attgctaactg tgtctttttt 1260  
 gttctggcta gcaagacag aatattctac gtaaatctota tgtcaagtca aacogtgcca 1320  
 cataaataa aatgtaggga atatagatca attagtgtct aagtgtacaa ctatatatat 1380  
 gttatctaa aatagattag attacaaaaa tgtatttacc attgattgat cactactttt 1440  
 cagogaatta gtcaagaggt cagggttatt tataaacat gcataatgta tacatatcat 1500  
 gcogtggcca ttgtacnatt atgttataaa aggtatacat taatataaac ttgtatttat 1560  
 tttttataca tgtcagcaat gtttgaaagt gtgaagttcc ctggttotta taattcaatc 1620  
 ttctgataat gtttttaacc ggccagaaca cagcattaac tccattaatc ataccaatta 1680  
 tgaccatgga atcagattct agtaacaaca taccoggtca tattctacaa gtggggcctg 1740  
 aggagggtaa ggtatacgt aaccatacac ttaaggtaga gaaattattt ctgaaagatc 1800  
 ctgactcac acgtttcgaa aaataggttt aaaaaatact tacataaaaa aaaaaataa 1860  
 aaataaagat acagataaag ataaacagag taaacaaaac aacaaatgga atatacagac 1920  
 aaaaagcaag ataaaatag tatatggaag aaaaatacaa atgcttatat gacagaagtc 1980  
 gtcgactac cttcaaacgt tctactotaa tcttgacct ccatacactc ctatctaagg 2040  
 ttatgtctc gattatcga aagattgaat cttgaatcaa atttatata tacaagaatt 2100  
 atttaactcc tgtattacaa gttcttatat ttcactgaac aaccactttt ggttcacaa 2160  
 gaatagtga atatagataa aattatctct aataactttg gacgagggat tatcatttaa 2220  
 atgtaataag aaaaatgtcg atggtattgg aagtcaaac aaagtccac atcggtagtt 2280  
 gaaaagttt gaatccaacg tataaggtgt atgtatctct taatggtgta agacatttta 2340  
 tgaaaactgt gttggcttag ccaaaagcga acaatatcac tccatttcaa gaatatcttg 2400  
 gttgttttag tccagcaact ggtatcagag caaatggttt agcaaatga gtatgttgtg 2460  
 tagtgattgc gtggggcatg acttagcctt tacccttgac ttggagaggg ccgggttaat 2520  
 gtctttgctc atctacagcc agtttattac ctttccatgt agctttaaag acgcacacag 2580  
 aggtattoyg ttgatgtggg tgacacaaa tazatctoca aataaaccca atagtggtag 2640

2700 ttgtctatgt gaactatgt tgggggggga gatttttgag agtctgagaa aaggtctca  
 2760 oattatagag aagattttag gaggctatct ataaagttata tggatctctc aataatgtgg  
 2820 gggtttatag gaatttatgt ttctctata tgccttata atgattagb tacyggagaa  
 2880 tcaagcttcc taanaattag gcaagtcca tgcctcaatg tccatctga ttgttaggct  
 2940 ataaagtag aatacgttta agaaagttta taattacgt ataatccaa aagtgaatg  
 3000 tgtttatna ttgttatac tcaatttgt gttatgtga tcaatttc gtcaggatca  
 3060 gttccaaag attagccaa aataatag tgttataa atgttaact tgggaactaa  
 3120 cttatagt agtcgatct attagatag gtaaaacttc acgttgttat ttgcaagcaag  
 3180 acaatnaatg cagatnatat gattatcat aaattttat catttgatca tcatgtttaa  
 3240 tacttcaac gttccaaat agatgtgtag ttctacatc ttattatcaa aataatgtt  
 3300 aattagat ataatgaaa ttaattttt ttctaaacca tattgtcttc gctattgaaa  
 3360 agtgaata tgaatctta ttctttaca tatacaatg aataatna gaataatag  
 3420 gtaaatata tgtatgtag ataatgttc ttctaatgg gcaataaat gttgaatca  
 3480 ataatnaog gataaggggt atattctca gctcaacta atcaatttg gttgaatna  
 3540 cgaatgcag tattaaoc agttttgga ataatgtacc atcaagaaac gcaatcgatg  
 3600 taactcaaaa atagtcaaaa caaattccct cactgtcca ttgcaattt atctagttc  
 3660 aattttacat atataaaagc caatagatc tatatagct ttgtactcaa caatccagca  
 3720 maactatag caatatcttc caacttcc tccgaatct ctctcaattg tatatcttt  
 3731 cctcttaac a

<210> 25  
 <211> 31  
 <212> DNA  
 <213> clloria  
 <400> 25  
 ggtatccaa atgttctctc ctagaagaa t

<210> 26  
 <211> 1831  
 <212> DNA  
 <213> kenna  
 <400> 26  
 cccctgata tcccaatba agggaaaaa agctgagct ccaacgaggt ggcggcagct  
 60  
 ctagaactag tggatccccc gggctgcaag aattttta atctnaaga ttctgataaa  
 120

- 39 -

attttaaato	ttagtacggc	atggccaact	tagatcaatt	gttcootttc	aaagaaattg	180
ctatgtocat	tttgatcttc	ttgatcactc	aactcaccat	tcattcactc	ttcacaaacc	240
gtcacaataa	gottocacca	gggootagag	gttgccaat	cgtaggtgac	ctccctgtct	300
tgggaagcat	gcctaatgtc	aocctotcta	gaatggcaa	aaagtatgga	gcctgcatgt	360
acctcaagat	gggcacccaa	aacttggttg	tgacctctac	tcocgotgca	gctcgtgcat	420
ccctcaaac	ccttgatcaa	aacttctcca	accgcoctcc	aaatgctggt	gcaactcaact	480
tagcttatga	ttcacaggac	atggtgtttg	ccactatagg	ctctaggtgg	aggttgctta	540
ggaaactgag	caacttgca	atgctgggtg	gaaaggctct	tgatgattgg	gcacatgttc	600
gggagaaaga	gatgaggtac	atgcttggtt	caatgtatga	ttgtagcaaa	aggggtgagg	660
ctgtggtggt	ggctgagatg	ttgacatag	ctatggccaa	tatgattggt	caagtgarat	720
tgagccgtcg	tgtgttcgag	tcaagggtt	cggaatcaaa	cgagttcaag	gacatggttg	780
ttgagctcat	gaccgttgcc	gggtacttca	acattggaga	ttttgtccct	tttcttgctt	840
ggtttgactt	gcaaggcata	gagcgtgaga	tgaaggcctt	gcataagaag	tttgatgcgt	900
tgttgacaag	gatgattgag	gagcatgtgg	ctctagatg	tcacaaaggt	aaaggaaact	960
atgatttctt	agaagtgtg	atggatcatt	ctagcgaag	cagtgatgga	gagagaactca	1020
cactcaccaa	tgtcaaggca	ctgctottga	atcttttcac	agcaggcact	gatacatctt	1080
cgagtgtgat	agagtgggca	ctagcggaga	tgttgaaaaa	tccccacata	acaaagagag	1140
ctcatgagga	aatggacca	gtcataggca	aggatcgacg	cctcaaggaa	tctgacctaa	1200
ggaaccttcc	ttacttgcaa	gotatttgca	aagaggcatt	gagaaagcac	ccttcaaccc	1260
cattgaactt	gcctagagtc	tcatacaaac	cgtagccaagt	gaatggatat	tacatcccca	1320
agaacactag	gttgagtgtg	aacatatggg	ccattggag	agcccccgag	gtgtgggaga	1380
agccttgtga	gttcaatcct	gagaggttta	tgagtggaaa	aggtagccaa	gttgatccac	1440
atgggaatga	ttttgagctg	attccgtttg	gtgctgggag	aagggtgtgt	gctgggacaa	1500
ggatggggat	tgtgatggtt	cagtacatat	tgggcacttt	ggtgcactca	tttgaatggg	1560
agctaccaaa	tgggttggtg	gagttgaaca	tggaaagagc	ctttgggctt	gccttgacga	1620
aaaagggtgc	actctcggtt	ttggttagcc	ctaggttgca	accaagttct	tatatccagt	1680
agagttgggt	ttggtttggt	tcaccaactc	tgttcaaaac	ttatgtctag	ctatttaaaa	1740
attacaatat	atgctttaag	gttatgtgac	tatatattgc	gcaaacggcg	caataataa	1800

1831

stgtgotttg gctcaaaaaa a

<210> 27  
<211> 513  
<212> PRT  
<213> kennedia  
<400> 27

Met Ala Asn Leu Asp His Leu Phe Leu Leu Lys Gln Ile Ala Met Ser  
1  
5  
10  
15  
20  
25  
30  
35  
40  
45  
50  
55  
60  
65  
70  
75  
80  
85  
90  
95  
100  
105  
110  
115  
120  
125  
130  
135  
140  
145  
150  
155  
160  
165  
170  
175  
180  
185  
190  
195  
200  
205  
210  
215  
220  
225  
230  
235  
240  
245  
250  
255  
260  
265  
270  
275  
280  
285  
290  
295  
300  
305  
310  
315  
320  
325  
330  
335  
340  
345  
350  
355  
360  
365  
370  
375  
380  
385  
390  
395  
400  
405  
410  
415  
420  
425  
430  
435  
440  
445  
450  
455  
460  
465  
470  
475  
480  
485  
490  
495  
500  
505  
510  
515  
520  
525  
530  
535  
540  
545  
550  
555  
560  
565  
570  
575  
580  
585  
590  
595  
600  
605  
610  
615  
620  
625  
630  
635  
640  
645  
650  
655  
660  
665  
670  
675  
680  
685  
690  
695  
700  
705  
710  
715  
720  
725  
730  
735  
740  
745  
750  
755  
760  
765  
770  
775  
780  
785  
790  
795  
800  
805  
810  
815  
820  
825  
830  
835  
840  
845  
850  
855  
860  
865  
870  
875  
880  
885  
890  
895  
900  
905  
910  
915  
920  
925  
930  
935  
940  
945  
950  
955  
960  
965  
970  
975  
980  
985  
990  
995  
1000  
1005  
1010  
1015  
1020  
1025  
1030  
1035  
1040  
1045  
1050  
1055  
1060  
1065  
1070  
1075  
1080  
1085  
1090  
1095  
1100  
1105  
1110  
1115  
1120  
1125  
1130  
1135  
1140  
1145  
1150  
1155  
1160  
1165  
1170  
1175  
1180  
1185  
1190  
1195  
1200  
1205  
1210  
1215  
1220  
1225  
1230  
1235  
1240  
1245  
1250  
1255  
1260  
1265  
1270  
1275  
1280  
1285  
1290  
1295  
1300  
1305  
1310  
1315  
1320  
1325  
1330  
1335  
1340  
1345  
1350  
1355  
1360  
1365  
1370  
1375  
1380  
1385  
1390  
1395  
1400  
1405  
1410  
1415  
1420  
1425  
1430  
1435  
1440  
1445  
1450  
1455  
1460  
1465  
1470  
1475  
1480  
1485  
1490  
1495  
1500  
1505  
1510  
1515  
1520  
1525  
1530  
1535  
1540  
1545  
1550  
1555  
1560  
1565  
1570  
1575  
1580  
1585  
1590  
1595  
1600  
1605  
1610  
1615  
1620  
1625  
1630  
1635  
1640  
1645  
1650  
1655  
1660  
1665  
1670  
1675  
1680  
1685  
1690  
1695  
1700  
1705  
1710  
1715  
1720  
1725  
1730  
1735  
1740  
1745  
1750  
1755  
1760  
1765  
1770  
1775  
1780  
1785  
1790  
1795  
1800  
1805  
1810  
1815  
1820  
1825  
1830  
1835  
1840  
1845  
1850  
1855  
1860  
1865  
1870  
1875  
1880  
1885  
1890  
1895  
1900  
1905  
1910  
1915  
1920  
1925  
1930  
1935  
1940  
1945  
1950  
1955  
1960  
1965  
1970  
1975  
1980  
1985  
1990  
1995  
2000  
2005  
2010  
2015  
2020  
2025  
2030  
2035  
2040  
2045  
2050  
2055  
2060  
2065  
2070  
2075  
2080  
2085  
2090  
2095  
2100  
2105  
2110  
2115  
2120  
2125  
2130  
2135  
2140  
2145  
2150  
2155  
2160  
2165  
2170  
2175  
2180  
2185  
2190  
2195  
2200  
2205  
2210  
2215  
2220  
2225  
2230  
2235  
2240  
2245  
2250  
2255  
2260  
2265  
2270  
2275  
2280  
2285  
2290  
2295  
2300  
2305  
2310  
2315  
2320  
2325  
2330  
2335  
2340  
2345  
2350  
2355  
2360  
2365  
2370  
2375  
2380  
2385  
2390  
2395  
2400  
2405  
2410  
2415  
2420  
2425  
2430  
2435  
2440  
2445  
2450  
2455  
2460  
2465  
2470  
2475  
2480  
2485  
2490  
2495  
2500  
2505  
2510  
2515  
2520  
2525  
2530  
2535  
2540  
2545  
2550  
2555  
2560  
2565  
2570  
2575  
2580  
2585  
2590  
2595  
2600  
2605  
2610  
2615  
2620  
2625  
2630  
2635  
2640  
2645  
2650  
2655  
2660  
2665  
2670  
2675  
2680  
2685  
2690  
2695  
2700  
2705  
2710  
2715  
2720  
2725  
2730  
2735  
2740  
2745  
2750  
2755  
2760  
2765  
2770  
2775  
2780  
2785  
2790  
2795  
2800  
2805  
2810  
2815  
2820  
2825  
2830  
2835  
2840  
2845  
2850  
2855  
2860  
2865  
2870  
2875  
2880  
2885  
2890  
2895  
2900  
2905  
2910  
2915  
2920  
2925  
2930  
2935  
2940  
2945  
2950  
2955  
2960  
2965  
2970  
2975  
2980  
2985  
2990  
2995  
3000  
3005  
3010  
3015  
3020  
3025  
3030  
3035  
3040  
3045  
3050  
3055  
3060  
3065  
3070  
3075  
3080  
3085  
3090  
3095  
3100  
3105  
3110  
3115  
3120  
3125  
3130  
3135  
3140  
3145  
3150  
3155  
3160  
3165  
3170  
3175  
3180  
3185  
3190  
3195  
3200  
3205  
3210  
3215  
3220  
3225  
3230  
3235  
3240  
3245  
3250  
3255  
3260  
3265  
3270  
3275  
3280  
3285  
3290  
3295  
3300  
3305  
3310  
3315  
3320  
3325  
3330  
3335  
3340  
3345  
3350  
3355  
3360  
3365  
3370  
3375  
3380  
3385  
3390  
3395  
3400  
3405  
3410  
3415  
3420  
3425  
3430  
3435  
3440  
3445  
3450  
3455  
3460  
3465  
3470  
3475  
3480  
3485  
3490  
3495  
3500  
3505  
3510  
3515  
3520  
3525  
3530  
3535  
3540  
3545  
3550  
3555  
3560  
3565  
3570  
3575  
3580  
3585  
3590  
3595  
3600  
3605  
3610  
3615  
3620  
3625  
3630  
3635  
3640  
3645  
3650  
3655  
3660  
3665  
3670  
3675  
3680  
3685  
3690  
3695  
3700  
3705  
3710  
3715  
3720  
3725  
3730  
3735  
3740  
3745  
3750  
3755  
3760  
3765  
3770  
3775  
3780  
3785  
3790  
3795  
3800  
3805  
3810  
3815  
3820  
3825  
3830  
3835  
3840  
3845  
3850  
3855  
3860  
3865  
3870  
3875  
3880  
3885  
3890  
3895  
3900  
3905  
3910  
3915  
3920  
3925  
3930  
3935  
3940  
3945  
3950  
3955  
3960  
3965  
3970  
3975  
3980  
3985  
3990  
3995  
4000  
4005  
4010  
4015  
4020  
4025  
4030  
4035  
4040  
4045  
4050  
4055  
4060  
4065  
4070  
4075  
4080  
4085  
4090  
4095  
4100  
4105  
4110  
4115  
4120  
4125  
4130  
4135  
4140  
4145  
4150  
4155  
4160  
4165  
4170  
4175  
4180  
4185  
4190  
4195  
4200  
4205  
4210  
4215  
4220  
4225  
4230  
4235  
4240  
4245  
4250  
4255  
4260  
4265  
4270  
4275  
4280  
4285  
4290  
4295  
4300  
4305  
4310  
4315  
4320  
4325  
4330  
4335  
4340  
4345  
4350  
4355  
4360  
4365  
4370  
4375  
4380  
4385  
4390  
4395  
4400  
4405  
4410  
4415  
4420  
4425  
4430  
4435  
4440  
4445  
4450  
4455  
4460  
4465  
4470  
4475  
4480  
4485  
4490  
4495  
4500  
4505  
4510  
4515  
4520  
4525  
4530  
4535  
4540  
4545  
4550  
4555  
4560  
4565  
4570  
4575  
4580  
4585  
4590  
4595  
4600  
4605  
4610  
4615  
4620  
4625  
4630  
4635  
4640  
4645  
4650  
4655  
4660  
4665  
4670  
4675  
4680  
4685  
4690  
4695  
4700  
4705  
4710  
4715  
4720  
4725  
4730  
4735  
4740  
4745  
4750  
4755  
4760  
4765  
4770  
4775  
4780  
4785  
4790  
4795  
4800  
4805  
4810  
4815  
4820  
4825  
4830  
4835  
4840  
4845  
4850  
4855  
4860  
4865  
4870  
4875  
4880  
4885  
4890  
4895  
4900  
4905  
4910  
4915  
4920  
4925  
4930  
4935  
4940  
4945  
4950  
4955  
4960  
4965  
4970  
4975  
4980  
4985  
4990  
4995  
5000  
5005  
5010  
5015  
5020  
5025  
5030  
5035  
5040  
5045  
5050  
5055  
5060  
5065  
5070  
5075  
5080  
5085  
5090  
5095  
5100  
5105  
5110  
5115  
5120  
5125  
5130  
5135  
5140  
5145  
5150  
5155  
5160  
5165  
5170  
5175  
5180  
5185  
5190  
5195  
5200  
5205  
5210  
5215  
5220  
5225  
5230  
5235  
5240  
5245  
5250  
5255  
5260  
5265  
5270  
5275  
5280  
5285  
5290  
5295  
5300  
5305  
5310  
5315  
5320  
5325  
5330  
5335  
5340  
5345  
5350  
5355  
5360  
5365  
5370  
5375  
5380  
5385  
5390  
5395  
5400  
5405  
5410  
5415  
5420  
5425  
5430  
5435  
5440  
5445  
5450  
5455  
5460  
5465  
5470  
5475  
5480  
5485  
5490  
5495  
5500  
5505  
5510  
5515  
5520  
5525  
5530  
5535  
5540  
5545  
5550  
5555  
5560  
5565  
5570  
5575  
5580  
5585  
5590  
5595  
5600  
5605  
5610  
5615  
5620  
5625  
5630  
5635  
5640  
5645  
5650  
5655  
5660  
5665  
5670  
5675  
5680  
5685  
5690  
5695  
5700  
5705  
5710  
5715  
5720  
5725  
5730  
5735  
5740  
5745  
5750  
5755  
5760  
5765  
5770  
5775  
5780  
5785  
5790  
5795  
5800  
5805  
5810  
5815  
5820  
5825  
5830  
5835  
5840  
5845  
5850  
5855  
5860  
5865  
5870  
5875  
5880  
5885  
5890  
5895  
5900  
5905  
5910  
5915  
5920  
5925  
5930  
5935  
5940  
5945  
5950  
5955  
5960  
5965  
5970  
5975  
5980  
5985  
5990  
5995  
6000  
6005  
6010  
6015  
6020  
6025  
6030  
6035  
6040  
6045  
6050  
6055  
6060  
6065  
6070  
6075  
6080  
6085  
6090  
6095  
6100  
6105  
6110  
6115  
6120  
6125  
6130  
6135  
6140  
6145  
6150  
6155  
6160  
6165  
6170  
6175  
6180  
6185  
6190  
6195  
6200  
6205  
6210  
6215  
6220  
6225  
6230  
6235  
6240  
6245  
6250  
6255  
6260  
6265  
6270  
6275  
6280  
6285  
6290  
6295  
6300  
6305  
6310  
6315  
6320  
6325  
6330  
6335  
6340  
6345  
6350  
6355  
6360  
6365  
6370  
6375  
6380  
6385  
6390  
6395  
6400  
6405  
6410  
6415  
6420  
6425  
6430  
6435  
6440  
6445  
6450  
6455  
6460  
6465  
6470  
6475  
6480  
6485  
6490  
6495  
6500  
6505  
6510  
6515  
6520  
6525  
6530  
6535  
6540  
6545  
6550  
6555  
6560  
6565  
6570  
6575  
6580  
6585  
6590  
6595  
6600  
6605  
6610  
6615  
6620  
6625  
6630  
6635  
6640  
6645  
6650  
6655  
6660  
6665  
6670  
6675  
6680  
6685  
6690  
6695  
6700  
6705  
6710  
6715  
6720  
6725  
6730  
6735  
6740  
6745  
6750  
6755  
6760  
6765  
6770  
6775  
6780  
6785  
6790  
6795  
6800  
6805  
6810  
6815  
6820  
6825  
6830  
6835  
6840  
6845  
6850  
6855  
6860  
6865  
6870  
6875  
6880  
6885  
6890  
6895  
6900  
6905  
6910  
6915  
6920  
6925  
6930  
6935  
6940  
6945  
6950  
6955  
6960  
6965  
6970  
6975  
6980  
6985  
6990  
6995  
7000  
7005  
7010  
7015  
7020  
7025  
7030  
7035  
7040  
7045  
7050  
7055  
7060  
7065  
7070  
7075  
7080  
7085  
7090  
7095  
7100  
7105  
7110  
7115  
7120  
7125  
7130  
7135  
7140  
7145  
7150  
7155  
7160  
7165  
7170  
7175  
7180  
7185  
7190  
7195  
7200  
7205  
7210  
7215  
7220  
7225  
7230  
7235  
7240  
7245  
7250  
7255  
7260  
7265  
7270  
7275  
7280  
7285  
7290  
7295  
7300  
7305  
7310  
7315  
7320  
7325  
7330  
7335  
7340  
7345  
7350  
7355  
7360  
7365  
7370  
7375  
7380  
7385  
7390  
7395  
7400  
7405  
7410  
7415  
7420  
7425  
7430  
7435  
7440  
7445  
7450  
7455  
7460  
7465  
7470  
7475  
7480  
7485  
7490  
7495  
7500  
7505  
7510  
7515  
7520  
7525  
7530  
7535  
7540  
7545  
7550  
7555  
7560  
7565  
7570  
7575  
7580  
7585  
7590  
7595  
7600  
7605  
7610  
7615  
7620  
7625  
7630  
7635  
7640  
7645  
7650  
7655  
7660  
7665  
7670  
7675  
7680  
7685  
7690  
7695  
7700  
7705  
7710  
7715  
7720  
7725  
7730  
7735  
7740  
7745  
7750  
7755  
7760  
7765  
7770  
7775  
7780  
7785  
7790  
7795  
7800  
7805  
7810  
7815  
7820  
7825  
7830  
7835  
7840  
7845  
7850  
7855  
7860  
7865  
7870  
7875  
7880  
7885  
7890  
7895  
7900  
7905  
7910  
7915  
7920  
7925  
7930  
7935  
7940  
7945  
7950  
7955  
7960  
7965  
7970  
7975  
7980  
7985  
7990  
7995  
8000  
8005  
8010  
8015  
8020  
8025  
8030  
8035  
8040  
8045  
8050  
8055  
8060  
8065  
8070  
8075  
8080  
8085  
8090  
8095  
8100  
8105  
8110  
8115  
8120  
8125  
8130  
8135  
8140  
8145  
8150  
8155  
8160  
8165  
8170  
8175  
8180  
8185  
8190  
8195  
8200  
8205  
8210  
8215  
8220  
8225  
8230  
8235  
8240  
8245  
8250  
8255  
8260  
8265  
8270  
8275  
8280  
8285  
8290  
8295  
8300  
8305  
8310  
8315  
8320  
8325  
8330  
8335  
8340  
8345  
8350  
8355  
8360  
8365  
8370  
8375  
8380  
8385  
8390  
8395  
8400  
8405  
8410  
8415  
8420  
8425  
8430  
8435  
8440  
8445  
8450  
8455  
8460  
8465  
8470  
8475  
8480  
8485  
8490  
8495  
8500  
8505  
8510  
8515  
8520  
8525  
8530  
8535  
8540  
8545  
8550  
8555  
8560  
8565  
8570  
8575  
8580  
8585  
8590  
8595  
8600  
8605  
8610  
8615  
8620  
8625  
8630  
8635  
8640  
8645  
8650  
8655  
8660  
8665  
8670  
8675  
8680  
8685  
8690  
8695  
8700  
8705  
8710  
8715  
8720  
8725  
8730  
8735  
8740  
8745  
8750  
8755  
8760  
8765  
8770  
8775  
8780  
8785  
8790  
8795  
8800  
8805  
8810  
8815  
8820  
8825  
8830  
8835  
8840  
8845  
8850  
8855  
8860  
8865  
8870  
8875  
8880  
8885  
8890  
8895  
8900  
8905  
8910  
8915  
8920  
8925  
8930  
8935  
8940  
8945  
8950  
8955  
8960  
8965  
8970  
8975  
8980  
8985  
8990  
8995  
9000  
9005  
9010  
9015  
9020  
9025  
9030  
9035  
9040  
9045  
9050  
9055  
9060  
9065  
9070  
9075  
9080  
9085  
9090  
9095  
9100  
9105  
9110  
9115  
9120  
9125  
9130  
9135  
9140  
9145  
9150  
9155  
9160  
9165  
9170  
9175  
9180  
9185  
9190  
9195  
9200  
9205  
9210  
9215  
9220  
9225  
9230  
9235  
9240  
9245  
9250  
9255  
9260  
9265  
9270  
9275  
9280  
9285  
9290  
9295  
9300  
9305  
9310  
9315  
9320  
9325  
9330  
9335  
9340  
9345  
9350  
9355  
9360  
9365  
9370  
9375  
9380  
9385  
9390  
9395  
9400  
9405  
9410  
9415  
9420  
9425  
9430  
9435  
9440  
9445  
9450  
9455  
9460  
9465  
9470  
9475  
9480  
9485  
9490  
9495  
9500  
9505  
9510  
9515  
9520  
9525  
9530  
9535  
9540  
9545  
9550  
9555  
9560  
9565  
9570  
9575  
9580  
9585  
9590  
9595  
9600  
9605  
9610  
9615  
9620  
9625  
9630  
9635  
9640  
9645  
9650  
9655  
9660  
9665  
9670  
9675  
9680  
9685  
9690  
9695  
9700  
9705  
9710  
9715  
9720  
9725  
9730  
9735  
9740  
9745  
9750  
9755  
9760  
9765  
9770  
9775  
9780  
9785  
9790  
9795  
9800  
9805  
9810  
9815  
9820  
9825  
9830  
9835  
9840  
9845  
9850  
9855  
9860  
9865  
9870  
9875  
9880  
9885  
9890  
9895  
9900  
9905  
9910  
9915  
9920  
9925  
9930  
9935  
9940  
9945  
9950  
9955  
9960  
9965  
9970  
9975  
9980  
9985  
9990  
9995  
10000  
10005  
10010  
10015  
10020  
10025  
10030  
10035  
10040  
10045  
10050  
10055  
10060  
10065  
10070  
10075  
10080  
10085  
10090  
10095  
10100  
10105  
10110  
10115  
10120  
10125  
10130  
10135  
10140  
10145  
10150  
10155  
10160  
10165  
10170

- 41 -

Ile Leu Ser Arg Arg Val Phe Glu Ser Lys Gly Ser Glu Ser Asn Glu  
195 200 205

Phe Lys Asp Met Val Val Glu Leu Met Thr Val Ala Gly Tyr Phe Asn  
210 215 220

Ile Gly Asp Phe Val Pro Phe Leu Ala Trp Phe Asp Leu Gln Gly Ile  
225 230 235 240

Glu Arg Glu Met Lys Ala Leu His Lys Lys Phe Asp Ala Leu Leu Thr  
245 250 255

Arg Met Ile Glu Glu His Val Ala Ser Arg Cys His Lys Gly Lys Gly  
260 265 270

Asn Tyr Asp Phe Leu Asp Val Val Met Asp His Ser Ser Glu Ser Ser  
275 280 285

Asp Gly Glu Arg Leu Thr Leu Thr Asn Val Lys Ala Leu Leu Leu Asn  
290 295 300

Leu Phe Thr Ala Gly Thr Asp Thr Ser Ser Ser Val Ile Glu Trp Ala  
305 310 315 320

Leu Ala Glu Met Leu Lys Asn Pro His Ile Thr Lys Arg Ala His Glu  
325 330 335

Glu Met Asp Gln Val Ile Gly Lys Asp Arg Arg Leu Lys Glu Ser Asp  
340 345 350

Leu Arg Asn Leu Pro Tyr Leu Gln Ala Ile Cys Lys Glu Ala Leu Arg  
355 360 365

Lys His Pro Ser Thr Pro Leu Asn Leu Pro Arg Val Ser Ser Gln Pro  
370 375 380

Cys Gln Val Asn Gly Tyr Tyr Ile Pro Lys Asn Thr Arg Leu Ser Val  
385 390 395 400

Asn Ile Trp Ala Ile Gly Arg Asp Pro Glu Val Trp Glu Asn Pro Cys  
405 410 415

-42-

GLU PHE ASN PRO GLU ARG PHE MET SER GLY LYS GLY ALA LYS VAL ASP  
420 425 430

PRO HIS GLY ASN ASP PHE GLU LEU ILE PRO PHE GLY ALA GLY ARG ARG  
440 445

VAL CYS ALA GLY THR ARG MET GLY ILE VAL MET VAL GLN TYR ILE LEU  
450 455 460

GLY THR LEU VAL HIS SER PHE GLU TRP LYS LEU PRO ASN GLY VAL VAL  
470 475 480

GLN LEU ASN MET GLN GLN THR PHE GLY LEU ALA LEU GLN LYS LYS VAL  
485 490 495

PRO LEU SER ALA LEU VAL SER PRO ARG LEU HIS PRO SER TYR ILE  
500 505 510

GLN

<210> 28

<211> 1374

<212> DNA

<213> chrysanthemum

<400> 28

GAATTCCGTC GTGTCGCTA CTTCAAGA TCAATACAT TTCGTTCC CAGATTA  
60  
CAAGGATGCG TCCTTAAC GACATGCGG CACTAGAGA GGTCCAAGG GTCCAAGT  
120  
CAGCTACCAT TCAAGCAT GGCACAGCA CTCGGCTAA TTGTGATAT CAGGCTAT  
180  
ATCCCGATTA CTAATTCGG ATCACTAAN GTGAACCAT GGTGGATCT AAGAGAA  
240  
TCAAGCGCAT GTGCGAAG TATATGATA GAAACGATA CAGGCACTC ACGGAGG  
300  
ATCTTAAGA GAACCAAC CTTGTGAGT AACTGGCTCC GTCCTCGAT GTCGCGC  
360  
ATGTGTGTGT GTTGTGATC CCAAGCTG GAAAGAGAG CCGAAGAAA GCTATTAA  
420  
AATGGAGCA ACCAATAT AATACACC AACTAATCT CTGCAACA TCTGTGTAG  
480  
AATAGCCCGG GTGTGATTA CAACTCAAC AATCCTCGG CTCGCGCAT TCGTCCAA  
540  
GTTTATGAT GTACCAACA GGTGTGCTTG AAGTGTGGA CGTCTTCGT CTAGCAAA  
600  
ACCTCGAAG AAAAACAAG GATGCAAGTG TCCAGTTGT TTGTCAGAG ATCACTG  
660  
TCAATCCG TGTCTTAC GACCTCATC TTGATCCT CTTGTGTCAA GCTCTGTTG  
720



- 43 -

```

gggatggagc tggggtgtc attgttggtt cagaccotga cttgacaaaa gagcgtccat 780
tggttcgagat gatattctgt gctcaxaacta tcttaaccaga ctccggaggga gcaatcgatg 840
ggcacttgag ggaagtcggg ctacacatttc atctctcaca agacgtacct ggggttgatct 900
ccaagaacat agagaaggca ttgacacaag cctttttctcc attaggtata agtgactgga 960
aotcgatctt ttggatcgct oactctggtg gtccagctat tctggacca gttgagotta 1020
agctcgggtct caaggaggag aagatgagag coantagaca cgttcttagt gagtatggaa 1080
acatgtcaag tgcttggtt ttgttcatta tggatgaaat gaggaagaaa tcggctgagg 1140
aaggtgagc cacaaccggt gaagggttag attggggtgt tttattcggg ttccgtcctg 1200
gittgacggt cgaaccggtg gtctccaca gcctcccaac cactgtatcg gttgcaatt 1260
aatttagttg catggttatg gatataagcg tcttttgttg gaacaattaa atttttactg 1320
tttttgtttt ctactaaata atgtgtgtt tgcaaaaaa aaaaaaaaa aaaa 1374

```

<210> 29  
 <211> 398  
 <212> PRT  
 <213> chrysanthemum

<400> 29

Met Ala Ser Leu Thr Asp Ile Ala Ala Ile Arg Glu Ala Gln Arg Ala  
 1 5 10 15

Gln Gly Pro Ala Thr Ile Leu Ala Ile Gly Thr Ala Thr Pro Ala Asn  
 20 25 30

Cys Val Tyr Gln Ala Asp Tyr Pro Asp Tyr Tyr Phe Arg Ile Thr Lys  
 35 40 45

Ser Glu His Met Val Asp Leu Lys Glu Lys Phe Lys Arg Met Cys Asp  
 50 55 60

Lys Ser Met Ile Arg Lys Arg Tyr Met His Leu Thr Glu Glu Tyr Leu  
 65 70 75 80

Lys Glu Asn Pro Asn Leu Cys Glu Tyr Met Ala Pro Ser Leu Asp Ala  
 85 90 95

Arg Gln Asp Val Val Val Val Glu Val Pro Lys Leu Gly Lys Glu Ala  
 100 105 110

Ala Thr Lys Ala Ile Lys Glu Trp Gly Glu Pro Lys Ser Lys Ile Thr  
 115 120 125  
 His Leu Ile Phe Cys Thr Thr Ser Gly Val Asp Met Pro Gly Ala Asp  
 130 135 140  
 Tyr Glu Leu Thr Lys Leu Gly Leu Arg Pro Ser Val Lys Arg Phe  
 145 150 155  
 Met Met Tyr Glu Glu Gly Cys Phe Ala Gly Gly Thr Val Leu Arg Leu  
 165 170 175  
 Ala Lys Asp Leu Ala Glu Asn Asn Lys Asp Ala Arg Val Leu Val Val  
 180 185 190  
 Cys Ser Glu Ile Thr Ala Val Thr Phe Arg Gly Pro Asn Asp Thr His  
 195 200 205  
 Leu Asp Ser Leu Val Gly Glu Ala Leu Phe Gly Asp Gly Ala Ala Ala  
 210 215 220  
 Val Ile Val Gly Ser Asp Pro Asp Leu Thr Lys Glu Arg Pro Leu Phe  
 225 230 235 240  
 Glu Met Ile Ser Ala Ala Glu Thr Ile Leu Pro Asp Ser Glu Gly Ala  
 245 250 255  
 Ile Asp Gly His Leu Arg Glu Val Gly Leu Thr Phe His Leu Lys  
 260 265 270  
 Asp Val Pro Gly Leu Ile Ser Lys Asn Ile Glu Lys Ala Leu Thr Glu  
 275 280 285  
 Ala Phe Ser Pro Leu Gly Ile Ser Asp Trp Asn Ser Ile Phe Trp Ile  
 290 295 300  
 Ala His Pro Gly Gly Pro Ala Ile Leu Asp Glu Val Glu Lys Leu  
 305 310 315 320  
 Gly Leu Lys Glu Lys Met Arg Ala Thr Arg His Val Leu Ser Glu  
 325 330 335

- 45 -

Tyr Gly Asn Met Ser Ser Ala Cys Val Leu Phe Ile Met Asp Glu Met  
340 345 350

Arg Lys Lys Ser Ala Glu Glu Gly Ala Ala Thr Thr Gly Glu Gly Leu  
355 360 365

Asp Trp Gly Val Leu Phe Gly Phe Gly Pro Gly Leu Thr Val Glu Thr  
370 375 380

Val Val Leu His Ser Leu Pro Thr Thr Val Ser Val Ala Asn  
385 390 395

<210> 30  
<211> 2979  
<212> DNA  
<213> chrysanthemum

<220>  
<221> misc feature  
<222> (2051)..(2051)  
<223> n = any nucleotide

<400> 30  
ttttaacgcg cattagcctc actaaagggg caaaagctgg agctcccccg ggtggcggcc 60  
getctgaact agtggatccc ccgggctgca ggaattcgat tggatgactc gaacagctat 120  
ggccatgatt acttagtacc acatgtact gagacttgca atggacaagt acttattatc 180  
ctacaacctc acttccttgt tgtgtttaag tccaaaaagt tatgcgtgcg gccagaatcc 240  
atcaaaatgt gtagcatttg ttccaacaca tgacctttaa ccgtatagt gttatgagtt 300  
ggtactooag acaatacatt aagagatata ttgggtatgc attgttgtgt atcatctccc 360  
acacatgctc agttatgtca gtatcaaat ctctctctc caaacacaaat totaattctc 420  
cttctctctc atctctaata totaaagtaa acatttgacc ttcaactta tgaccatgca 480  
tatacttctt atcagatata aaacatagat tcttagccct cttttcagca tacttttttt 540  
taacttaatta ctttctaggt gtaggcctgg tgtttgcaa ctgtttggtt accaactgag 600  
ttgtaggcaa agctaaaata ttgaaatat tttttagtag gataagtcac attcctgttt 660  
acattccaat tattattgta cttaggagta ggcaacaatg agacactttt ttctttaaca 720  
actgetagcc tagcttcttc catcttagcc caacaataga catcatttaa actagttggt 780  
ttaaacatgc taaccagcat aactatttgg ttttcaatc caccatata caaactaata 840



- 47 -

aaggcccttct gaanaattta atcaacagtt acctaatggt agattgatat gaacacaaaa 2640  
 cttctggtgt atgcagctgg tcatgacac tcaaatccgt aaccgaagtg tttagaatt 2700  
 atcgtattca cagtcataac ttacggttaa aactttaaac gaaatcgaac taaactccta 2760  
 acagatatac aagctcaatt gtgtaatgtt ttccaatggt ccacaaagtg gcattctatga 2820  
 ccacgtttcg taaaaattgg gtacgtcata cctaccacac gttccctcta tataagaaac 2880  
 actcattcac ctaattgtcta ccatcacttg caattctcta cttacaaaga tcatatacat 2940  
 ttctgttcac ccatattaaa caccatggc ttccttaac 2979

<210> 31  
 <211> 1778  
 <212> DNA  
 <213> lavendula

<400> 31  
 ctagtataaa ttttttaata gtaggcacgc aaaaatcaaga atctatcttc gtgatagcta 60  
 gagagctcac tatagcagcc tcaatctact ttctcatccg ctactttctt tcaagaatca 120  
 tcaccacacat taaccacggc ggcagccacc gactgcgccc agggccgagg ggctttccga 180  
 ttgtcgggtgc acttctcttc ttggggagaca tgcacatgt cgccttagcc aaaaatggca 240  
 aaacttacgg ccccatcata tacctaaaaa tgggtgcacg gggcatggcc gtcgcgtcaa 300  
 cgcctgcctc cgcctgtgcg ttctcaaaa ccttagacac caacttctct gacccgcctc 360  
 cgaatgcggg tgcacacata tttagcctaca acggggaaga tatggtgttc gcccgctatg 420  
 gcccaaagtg gagattgctc agaaaactga ccaatctcaa catgttgggg aatcatgott 480  
 tagatgggtg ggcaagtgtg aggtctcccg agttgggcta catgctccat gcaaggcagc 540  
 acgccacccg tcatggcgag cccgtggtgc tgcagagat gctcatgtac gccgtgggga 600  
 atatgctcgg gcaggtgata ttaagttagc ggaatttctc gaagaaggga aaggaggtga 660  
 atgagttgaa agatattggt gtggagctca tgaactcagc tggatatttc aatattggtg 720  
 attcatccc atggttgcct tggatggatt tgcaggggat agagagtggg atgaagaat 780  
 tgcacaataa gttcgacaag ttgatcgga aatgattga ggatcatttg aaatcagccc 840  
 acatacgcaa ggccaagccg gatcttcttg attgctctt ggcaaatcgt gatagctcgg 900  
 atgcggagaa gctcacctca accaacgtca aggcctttt actgaacttg ttcaccgcag 960  
 ggaccgacac gtoatcaagg ataatagaat gggcattggc cgagatgato aagaatccaa 1020  
 ccacctaaa tagggccac caagagatgg atagagtcgt tggtagaact cgaagggttg 1080

1140 tgaatcga catcccgac ctaccatcac tagagccat atgaaagaa acatccga  
 1200 agatccatc oaatccatc aatccgccc gaatccgctc agagccctgc gtcgtgagc  
 1260 ggtatcat accaaac accgctca gctgaacat atggctatc ggaagagac  
 1320 ccagctg ggaacatc ctgatatc acccgatg atctatc ggaagagac  
 1380 aggatcga tcccgccg gaacatc agctatc agctatc ggaagagac  
 1440 tctgcccga ggcgcgag ggaatgag tctgagatc tctgagatc ggaagagac  
 1500 agcttga atggagatc ccgcccgcgc ccgcccgcgc ccgcccgcgc tgaacatg  
 1560 accagctt tggcctgc atgagagag ctgcccctc ccgcccgc ctacatc  
 1620 ggcgcctc acatgctat gctccatc tctgctca tctatc tctatc  
 1680 tctatcga atgctcac accatgcat tctgctca tctatc tctatc  
 1740 catatg ctgctatc atctatc atctatc atctatc atctatc  
 1778 atatatc taagatc tcatatc aaaaaa

<210> 32  
 <211> 520  
 <212> PRT  
 <213> Lavandula  
 <400> 32

1 Met gin aan gin gin ser ile phe val ile ala arg gin leu thr ile  
 5  
 10

Ala ala ser ile tyr phe leu ile arg tyr phe leu ser arg ile ile  
 20  
 25  
 30

Thr thr ile thr his gly ser his arg leu pro pro gly pro arg  
 35  
 40  
 45

Gly phe pro ile val gly ala leu pro leu leu gly asp met pro his  
 50  
 55  
 60

Val ala leu ala lys met ala lys thr tyr gly pro ile ile tyr leu  
 65  
 70  
 75  
 80

Lys val gly ala trp gly met ala val ala ser thr pro ala ser ala  
 85  
 90  
 95

- 49 -

Arg Ala Phe Leu Lys Thr Leu Asp Thr Asn Phe Ser Asp Arg Pro Pro  
 100 105 110

Asn Ala Gly Ala Thr Ile Leu Ala Tyr Asn Ala Glu Asp Met Val Phe  
 115 120 125

Ala Arg Tyr Gly Pro Lys Trp Arg Leu Leu Arg Lys Leu Thr Asn Leu  
 130 135 140

His Met Leu Gly Asn His Ala Leu Asp Gly Trp Ala Ser Val Arg Ser  
 145 150 155 160

Ser Glu Leu Gly Tyr Met Leu His Ala Arg His Asp Ala Thr Arg His  
 165 170 175

Gly Glu Pro Val Val Leu Pro Glu Met Leu Met Tyr Ala Val Gly Asn  
 180 185 190

Met Leu Gly Gln Val Ile Leu Ser Arg Arg Ile Phe Glu Lys Lys Gly  
 195 200 205

Lys Glu Val Asn Glu Leu Lys Asp Met Val Val Glu Leu Met Thr Ser  
 210 215 220

Ala Gly Tyr Phe Asn Ile Gly Asp Phe Ile Pro Trp Leu Ala Trp Met  
 225 230 235 240

Asp Leu Gln Gly Ile Glu Ser Gly Met Lys Lys Leu His Asn Lys Phe  
 245 250 255

Asp Lys Leu Ile Gly Lys Met Ile Glu Asp His Leu Lys Ser Ala His  
 260 265 270

Ile Arg Lys Ala Lys Pro Asp Leu Leu Asp Cys Leu Leu Ala Asn Arg  
 275 280 285

Asp Ser Ser Asp Ala Glu Lys Leu Thr Ser Thr Asn Val Lys Ala Leu  
 290 295 300

Leu Leu Asn Leu Phe Thr Ala Gly Thr Asp Thr Ser Ser Ser Ile Ile  
 305 310 315 320

Glu Trp Ala Leu Ala Glu Met Ile Lys Asn Pro Thr Ile Leu Asn Arg

325	335	330	335
Ala His Gln Met Asp Arg Val Val Gln Arg Thr Arg Leu Val	340	345	350
Gln Ser Asp Ile Pro Asn Leu Pro Tyr Leu Arg Ala Ile Cys Tyr Gln	355	360	365
Thr Tyr Arg Lys His Pro Ser Thr Pro Leu Asn Leu Pro Arg Ile Ala	370	375	380
Ser Gln Pro Cys Val Val Asp Gln Tyr Tyr Ile Pro Lys Asn Thr Arg	385	390	395
Leu Ser Val Asn Ile Trp Ala Ile Gln Arg Asp Pro Asp Val Trp Gln	405	410	415
Asn Pro Leu Asp Phe Asn Pro Asp Arg Phe Leu Ser Gln Lys Asn Gln	420	425	430
Arg Ile Asp Pro Arg Gln Asn His Phe Gln Leu Ile Pro Phe Gln Ala	435	440	445
Gly Arg Arg Ile Cys Ala Gly Ala Arg Met Gly Met Val Leu Val Gln	450	455	460
Tyr Ile Leu Gly Thr Leu Val His Ala Phe Gln Trp Gln Leu Pro Ala	465	470	475
Gly Ala Gly Ala Gly Thr Ala Gln Leu Asn Met Asp His Val Phe Gly	485	490	495
Leu Ala Leu Gln Lys Ala Val Pro Leu Thr Ala Met Leu Thr Pro Arg	500	505	510
Leu Pro Ser His Cys Tyr Ala Pro	515		



# INTERNATIONAL SEARCH REPORT

International application No.

PCT/AU2003/001111

<b>A. CLASSIFICATION OF SUBJECT MATTER</b>		
Int. Cl. 7: C12N 15/53		
According to International Patent Classification (IPC) or to both national classification and IPC		
<b>B. FIELDS SEARCHED</b>		
Minimum documentation searched (classification system followed by classification symbols) See "electronic data base" box below		
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched See "electronic data base" box below		
Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) CA, WPIDS: flavonoid 3',5' hydroxylase; DGENE (blast) : SEQ ID NO: 5, 10, 12, 14, 16, 18, 21, 27, 30, 32		
<b>C. DOCUMENTS CONSIDERED TO BE RELEVANT</b>		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	EP 0 632 128 A1 (KYOWA HAKKO KOGYO CO., LTD.) 4 January 1995 See Claim 1; pages 43-46 [shares 68% identity with SEQ ID NO: 12; 66% with SEQ ID NO: 14; 65% with SEQ ID NO 16; 58% with SEQ ID NO: 21]	1-86, 89-92
X	EP 0 522 880 B1 (INTERNATIONAL FLOWER DEVELOPMENTS PTY. LTD.) 21 March 2001 See Fig. 10 [shares 69% identity with SEQ ID NO: 12; 66% with SEQ ID NO: 14; 58% with SEQ ID NO: 21, 73% with SEQ ID NO: 27, 64% with SEQ ID NO: 32]; Fig 9 [shares 65% with SEQ ID NO: 16]	1-86, 89-92
X	WO 2000009720 (INTERNATIONAL FLOWER DEVELOPMENTS PTY. LTD.) 24 February 2000 Specification	1-6, 11, 29-32, 35, 40, 45, 49, 57-82, 89-92
<input checked="" type="checkbox"/> Further documents are listed in the continuation of Box C <input checked="" type="checkbox"/> See patent family annex		
* Special categories of cited documents: "A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier application or patent but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art "&" document member of the same patent family	
Date of the actual completion of the international search 8 January 2004		Date of mailing of the international search report 22 JAN 2004
Name and mailing address of the ISA/AU AUSTRALIAN PATENT OFFICE PO BOX 200, WODEN ACT 2606, AUSTRALIA E-mail address: pct@ipaaustralia.gov.au Facsimile No. (02) 6285 3929		Authorized officer  JAMIE TURNER Telephone No : (02) 6283 2071

# INTERNATIONAL SEARCH REPORT

International application No.  
PCT/AU2003/001111

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 1996036716 A1 (INTERNATIONAL FLOWER DEVELOPMENTS PTY. LTD.) 21 November 1996 Specification	1-6, 11, 29-32, 35, 40, 45, 49, 57-82, 89-92
X	WO 1993020206 A1 (INTERNATIONAL FLOWER DEVELOPMENTS PTY. LTD.) 14 October 1993 See Fig 3A-D [shares 68% identity with SEQ ID NO: 12; 67% with SEQ ID NO: 14; 65% with SEQ ID NO: 16; 58% with SEQ ID NO: 21; 72% with SEQ ID NO: 27; 63% with SEQ ID NO: 32]	1-86, 89-92

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No.

PCT/AU2003/001111

This Annex lists the known "A" publication level patent family members relating to the patent documents cited in the above-mentioned international search report. The Australian Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

Patent Document Cited in Search Report				Patent Family Member			
EP	0632128	AU	29560/92	CA	2130800	CA	2365599
		US	6114601	US	6232109	US	2002100072
		WO	9318155				
EP	0522880	AU	19530/92	AU	22733/92	AU	67895/94
		CA	2112373	CA	2163220	CN	1071456
		CN	1127015	EP	0703982	IE	922272
		JP	2000023686	NZ	243500	NZ	266401
		PL	298239	PL	311691	SG	45175
		SG	45187	US	5349125	US	5569832
		US	5861487	US	5948955	WO	9301290
		WO	9428140	ZA	9205180		
WO	0009720	AU	53815/99				
WO	9636716	AU	56396/96	CA	2202668	EP	0873410
		HU	9802555	NZ	307119	US	6080920
WO	9320206	AU	37413/93	CA	2132961	EP	0640136
		NZ	249808	US	5639870		
END OF ANNEX							

